

Oral presentation

Ms Lujan Elena
lujan.e@procarehealth.com - Spain

Sanmartín Salinas Patricia
sanmartin.p@procarehealth.com - Spain

#3728

EFFECTIVENESS OF A MULTI-INGREDIENT CORIOLUS VERSICOLOR-BASED VAGINAL GEL IN HPV+ and HIV+ PATIENTS: A PILOT OBSERVATIONAL STUDY

34 - Sexually transmitted diseases and HIV infection

Sanmartín Salinas P², Joaquín Hijaña Elosegui J¹

¹Servicio de Obstetricia y Ginecología. Complejo Hospitalario de Jaén, Jaén, Spain

²Procure Health Iberia, Barcelona, Spain

Background/Objectives: Immunosuppressed human immunodeficiency virus (HIV) -positive patients are at greater risk of incident, persistent, or recurrent human papillomavirus (HPV) infection. They also have lower clearance rate, higher viral load, and a marked predisposition for being colonized by several serotypes; all leading to more frequent and severe HPV-dependent lesions 1. A Coriolus versicolor-based vaginal gel have shown to repair HPV-dependent low-grade cervical lesions and to increase high-risk HPV clearance in immunocompetent HPV-positive patients 2. The aim is to provide evidence about the effectiveness of a multi-ingredient Coriolus versicolor-based vaginal gel on HPV-dependent cervical alterations and HPV clearance in HIV+ patients

Methods: Pilot, prospective, one-cohort, observational study. 15 HIV-positive patients colonized by HPV in the endocervix region with an anomalous cervicovaginal cytology were included to receive a Coriolus versicolor-based vaginal gel 1 cannula/day for 21 days during first month + 1 cannula/alternate days for 5 months. Analysis of HPV patients with normal cytology and colposcopy image (improved alterations) and patients with HPV cleared (measured using hybrid capture test) is presented. The study was approved by an IRB and informed consent was signed by patients.

Results: The overall HPV clearance and cytological normalization rates were 73.33% and 80%, respectively. Endocervical colonization by HPV also partially cleared in 13.33% of the cases. At the end of the study, the normalization of the colposcopy anomalies associated to HPV was achieved in 55.56%.

Conclusions: Our results suggest that the proposed Coriolus versicolor-based vaginal gel treatment scheme could be an effective therapy in the management of endocervical HPV infection in HIV + patients. Its effects are similar to those obtained in patients without immunosuppression.

References: (1. Liu G, Sharma M, Tan N, Barnabas RV. HIV-positive women have higher risk of human papilloma virus infection, precancerous lesions, and cervical cancer. *AIDS*. 2018 Mar 27;32(6):795-808.) (2. Serrano L, López AC, González SP, Palacios S, Dexeus D, Centeno-Mediavilla C, Coronado P, de la Fuente J, López JA, Vanrell C, Cortés J. Efficacy of a Coriolus versicolor-Based Vaginal Gel in Women With Human Papillomavirus-Dependent Cervical Lesions: The PALOMA Study. *J Low Genit Tract Dis*. 2021 Apr 1;25(2):130-136)

Oral presentation

Prof. Dr Jeremic Igor
jerigor@yahoo.com - Serbia

Igor Jeremic
jerigor@yahoo.com -

#3065

The sync up of colposcopic findings with cervix precancerous changes in cervix with young female patients

25 - Colposcopy

Igor J

Background/Objectives: The study included 120 young female patients age 17-30 from Serbia, Montenegro Bosnia and Hercegovina as well as patients from EU countries. All patients had HPV confirmed diagnosis (DNK type procedure). Predominant types with patients in Serbia were 6,31,33,45,52,56,66 Montenegro: 16,18,44, 35, 33, 51,56,62,68 whereas the patients from Bosnia and Herzegovina and EU countries had a combination of the afore stated ones. The organism of young females is unprepared and the incubation period is shortened from 3 weeks to 3 months.

Methods: A colposcopy procedure has been done to all patients. Depending on the severity of HPV infection, colposcopy findings differed. With younger patients (17-21 years of age) more severe infection resulted in a combination of several pathological images (mosaic, punctation Aw epithelia) whereas with older patients (25-30 years of age) we found out often leuco-plaque and irregular vascular pattern. With these patients we performed the removal of the changes by application of radio wave LOOP excision. The essence of the technique is the combination of radio wave excision and deep vaporization. With excision we remove the change in the width of up to 3, 5 cm and depth of 2,5 cm, and with additional deep vaporization of the bottom of the change we remove dysplastic change, all in local anesthesia in 3 to 5 minutes.

Results: Each female patient had a confirmed diagnosis of HPV infection of the cervix. With 47% H-SIL has been diagnosed, and 21% of patients had HPV cervicitis condylomata. The majority of patients with H-SIL diagnosis were younger than 25 years and after colposcopy procedure the findings were a mixture of pathological images, which appeared immediately upon smearing of the cervix with dilution of vinegar acid (AW epithelia, rough mosaic, and punctuation in the majority of cases). L-SIL was dominant with colposcopy image of rough AW epithelia or mosaic.

Conclusions: Pathological colposcopic findings is a clear indicator of HPV infection of the cervix. In case there is a combination of a number of pathological images, it is necessary to apply an adequate therapeutic procedure in accordance with the patient's age. It is very important to note that combination of HPV viruses at this particular geographical location is entirely present in polyvalent HPV vaccines.

References: . Patented the special technique of radiowave LOOP excision with radiowave vaporization as a method of treatment of choice with difficult cases of dysplasia on PVU (CIN I,CIN II, CIN III) 5. Patented the special technique of removal of condyloms on vagina and mucous membrane of labia (radiowave vaporization) 6. Patented the technique of removal of giant anal and intraanal condyloma, with the use of the mixed technique of radiowave vaporization and excision, without bleeding. 20 years of work only in the field of HPV infections More than 6000 edited RF LOOP excisions.

HPV -RF therapy

Poster

Dr Soerensen Sarah M
sam@cancer.dk - Denmark

Dr Soerensen Sarah M
sam@cancer.dk - Denmark

#3076

INVESTIGATING SURVIVAL AFTER PENILE CANCER: A NATIONWIDE STUDY OF HUMAN PAPILLOMAVIRUS (HPV) AND P16 AS PROGNOSTIC MARKERS IN PENILE CANCER

15 - Molecular markers

Thomsen L, Toft B, Elversang J, Stilling C, Stifter S, Hjulskov S, Kjaer-frifeldt S, Waldstrøm M, Aagaard M, Jakobsen J, Petersen P, Munk C, Kjaer S

Background/Objectives: Human papillomavirus (HPV) positivity and expression of p16 have been suggested as potential prognostic markers in penile cancer. However, the majority of previous studies are limited by small sample sizes, lack of adjustment for other prognostic factors, and shorter follow-up. In this nationwide study, we will investigate the prognostic significance of HPV DNA status and p16 expression for survival after penile cancer in a large cohort of patients.

Methods: In the nationwide Danish Pathology register, we identified all cases of penile squamous cell carcinomas (SCC) in Denmark during 1995-2017. We restricted our retrieval of material to the six largest centers in the study period. Since 2009, treatment of penile cancer has been centralized at two centers (Rigshospitalet, Copenhagen and Aarhus University Hospital). Archived tumor material will be retrieved from all cases, and the diagnosis reviewed by a pathologist. HPV DNA testing of the tumor material will be performed using INNO LiPA HPV Genotyping Extra test, and p16 staining will be performed using the CINtec histology kit at a central laboratory. P16 slides will be evaluated and scored by two pathologists independently, and tumors with >70% tumor staining cells will be considered positive. From nationwide registries, we will obtain information on deaths and recurrences in the study population during up to 26 years. We will estimate the overall, disease-specific and disease-free survival after penile SCC according to HPV status and p16 expression. We will use Cox proportional hazards regression to investigate whether HPV and p16 have independent prognostic value after adjusting for other known prognostic factors, including cancer stage at diagnosis.

Results: Altogether, we identified 936 cases of penile SCC at the included pathology departments during 1995-2017, corresponding to approximately 82% of all cases in Denmark. Cases were diagnosed at pathology departments at: Aarhus University Hospital (N=350), Rigshospitalet, Copenhagen (N=320), Odense University Hospital (N=96), Aalborg University Hospital (N=67), Herlev Hospital (N=73), and Vejle, Lillebælt Hospital (N=30). 21.9% of cases were diagnosed during 1995-2002, 35.8% during 2003-2010, and 42.3% during 2011-2017. According to records, surgical specimens are available in 811 cases (86%), and biopsies in 80 cases (9%). In 45 cases (5%), the type of available tumor material is to be determined, as the tumor material is yet not retrieved from the archives.

Conclusions: Data collection for the study is ongoing. A more detailed description of the study design, final number of included cases and patient characteristics will be presented.

Poster

Prof. Cui Fuqiang
cuifuq@126.com - China

Prof. Cui Fuqiang
cuifuq@126.com - China

#3097

Estimating the prevalence of non-cervical human papillomavirus infection in mainland China: The design of PROGRESS-Plus study

03 - Epidemiology and natural history

Background/Objectives: Evidence indicated that human papillomavirus (HPV) infection may increase risks of anogenital warts, oral warts, and a varying proportion of several cancers among men and women. The prevalence of cervical HPV infection has been extensively reported in women worldwide. Much less has been reported on the disease burden of non-cervical HPV infection in women and men, especially in China. Thus, a study named PROGRESS (PREvalence of Oral hpv infection, a Global aSSessment)-Plus will be conducted in mainland China to estimate the prevalence of HPV oral infection in women, and of HPV oral and anogenital infection in men. This abstract describes the design of the PROGRESS-Plus study.

Methods: PROGRESS-Plus is a national, multi-site, cross-sectional study in China, conducted from 2020-2023.

Results: Approximately 4,742 participants (2,371 women and 2,371 men) aged 18-60 years will be recruited from 6 regions in China through 12 study sites during 2021 Nov-2022 Nov for oral/anogenital sample collection. HPV DNA detection and genotype identification will be performed using the SPF10 LiPA assay at a central laboratory. The HPV infection rates will be reported overall as well as by HPV genotype, age, and geographic region. The risk factors (e.g. demographics and sexual behaviors) associated with oral and anogenital HPV infection will be surveyed among study participants.

Conclusions: Results from the PROGRESS Plus study will fill important evidence gaps in the oral HPV burden in the Chinese general population and the anogenital HPV burden among men. This evidence can raise public awareness of HPV infection and help health care providers, policymakers, public health professionals, and health care consumers make decisions for gender-neutral HPV vaccination. In addition, the approaches established in the PROGRESS Plus study could provide a methodological framework for future epidemiologic research related to sexually transmitted diseases and other health outcomes in mainland China.

References: NA

Oral presentation

Ms Downham Laura
downhaml@iarc.fr - France

Downham Laura
downhaml@students.iarc.fr - France

#3117

Performance of oncoprotein-based tests to detect CIN2+ for cervical cancer screening and triage: a systematic literature review

10 - HPV screening

Downham L, Rol M, Ramirez T, Valls J, Gunter M, Almonte M

Background/Objectives: WHO current guidelines recommend using DNA HPV detection rather than cytology in primary screening and treatment approaches for cervical cancer. HPV testing has high sensitivity but moderate specificity to detect cervical intraepithelial neoplasia lesions 2 or greater (CIN2+). Therefore, accurate triage tests are needed to identify women at high risk of CIN2+. The HPV viral oncoproteins E6 and E7 are necessary for oncogenic transformation and are overexpressed during cervical carcinogenesis. The detection of these proteins are promising candidate markers for screening or triage. Objective: To provide an up-to-date systematic literature review of the clinical performance of oncoprotein-based HPV tests to detect CIN2+ and CIN3+ lesions for cervical cancer screening and triage.

Methods: A systematic literature search of PubMed, Embase and Web of Science was conducted. Studies reporting original data on the sensitivity and specificity of oncoprotein-based HPV tests for cervical cancer primary screening or triage in women aged 18+ were included. Histologically confirmed CIN2+ or CIN3+ were considered as the gold standard. A meta-analysis using mixed effects logistic regression models was performed to obtain heterogeneity-adjusted sensitivity and specificity estimates.

Results: Twenty-three studies were included in the analysis, the majority being observational studies (96%) and conducted in Asia (44%). Seventeen studies (74%) reported the performance of oncoprotein-based tests in screening settings while 6 studies (26%) in triage settings. Eighteen studies (78%) targeted the viral oncoprotein E6 alone and mean age was above 50 years old for most of the studies (74%). Sensitivity for CIN3+ and specificity (excluding CIN2) varied across studies (ranges 37.8%-100.0% and 67.9%-100.0% respectively) in screening settings, resulting in pooled estimates of 58.1% (95%CI 44.1-70.8) for sensitivity and 98.7% (95%CI 98.0-99.2) for specificity. Sensitivity and specificity for CIN2+ were 45.1% (95%CI 29.0-62.2) and 98.7% (95%CI 98.0-99.2) respectively. In a triage setting, the sensitivity for CIN3+ was higher (69.0%, 95% CI 57.6-78.4) and the specificity slightly lower (95.7%, 95% CI 90.8-98.0). The same pattern was observed for CIN2+ with a sensitivity of 58.2% (95%CI 50.5-65.5) and a specificity of 92.3% (95%CI 85.1-96.2).

Conclusions: These preliminary results show that oncoprotein-based HPV tests have better performance in triage than in screening settings, although a high heterogeneity was observed across studies, which could be due to the test target (different HPV types and oncoproteins), the sample collection method (collected by a clinician or self-collected), device (endocervical brush, cervical brush), or medium (Swab or PreservCyt). These factors will be further investigated and a study-quality assessment will be conducted.

Poster

Dr Cho Angela
angela012687@gmail.com - South Korea

Dr Cho Angela
angela012687@gmail.com - South Korea

#3137

Clinical characteristics and risk factors of invasiveness and recurrence in the patients with extramammary Paget's disease of the vulva

27 - Vulvar diseases and neoplasia

Shin E, Lee S, Park J, Kim D, Suh D, Kim J, Kim Y, Kim Y

Background/Objectives: Extramammary Paget's disease (EMPD) of vulva is a rare and slow growing disease. Studies about EMPD had not been established, and especially, studies on Asians are still lacking. We aimed to identify clinical features of patients with EMPD of the vulva in Korea, and evaluate the risk factors of invasiveness and recurrence.

Methods: We conducted retrospective analysis of the patients with EMPD of vulva who underwent surgical treatment in Asan medical center from 2005 to 2020. Clinical characteristics, histopathologic results, surgical complications, and recurrence were reviewed. The differences between the patients' clinical and histopathologic factors were examined using Chi-square and Fisher's exact tests. All statistical analyses were conducted using SPSS 20.0 software (Chicago, USA).

Results: Forty-seven patients with EMPD of vulva were treated with wide excision or radical vulvectomy in Asan medical center. The mean age at the time of initial diagnosis was 64.3 (range 44-80) years. Among them, 17 (36.2%) patients had invasive diseases, and seven (14.9%) patients experienced recurrence. Median duration of recurrence was 67.2 (range 32.8-160.0) months. Vulvar lesion size on physical examination was significantly associated with invasiveness ($p=0.01$). Margin status, adnexal involvement, and lesion sized was not significantly associated with recurrence (all $p>0.05$).

Conclusions: Vulvar lesion size on physical examination was significantly associated with invasive disease in the patients with EMPD of vulva. Margin status during surgery was not significantly associated with recurrence.

Oral presentation

Mr Partanen Veli-matti
veli-matti.partanen@cancer.fi - Finland

Partanen Veli-matti
veli-matti.partanen@cancer.fi - Finland

#3154

Correcting cervical cancer incidence and mortality in the Nordic countries by reallocation of unspecified uterine cancer cases and deaths.

40 - Public health

Partanen V, Heinävaara S, Anttila A, Hakkarainen J, Lönnberg S

Background/Objectives: The incidence of and mortality from cervical cancer is underestimated if the presence of uterine cancers, where the exact topography (site of origin) is not specified, is omitted. In this paper we present the corrected figures on mortality from cervical cancer in the Nordic countries by reallocating unspecified uterine cancer deaths and cases to originate either from corpus uteri or cervix uteri. To further validate the accuracy of reallocation, we also analyzed how well the reallocation captures the changes occurring due to a transition in cause of death coding in Norway that took place in 2005.

Methods: This study uses data available in the NORDCAN database, which contains aggregated cancer data from all the Nordic countries for years 1960-2016. The unspecified uterine cancer cases and deaths were reallocated to either cervix uteri or corpus uteri based on the estimated probability that follows the distribution of cases and deaths with verified topography. The estimated proportion of cases and deaths for both cancers were calculated for each combination of age group, year, and country as a proportion of cases (and deaths respectively) with known topography. Annual age-standardized rates were calculated by direct age-adjustment.

Results: The proportions of unspecified uterine cancers were higher in the mortality data than in incidence data, with mean values for years 1960-2016 ranging between 5.1-26.6% and 0.2-6.8% by country, respectively. In the Nordic countries combined, the reallocation increased the number of cases by 4% and deaths by approximately 20% for both cancers. Finland was the only Nordic country, where the mortality rate did not increase substantially after reallocation.

Conclusions: The reallocation procedure had a significant impact on mortality from cervical cancer for countries where the proportion of cancer deaths coded as uterus, not otherwise specified, is substantial. More effort to validate cause of death data with incidence data from cancer registries is warranted to avoid erroneous conclusions of temporal trends based on uncorrected cancer burden.

Oral presentation

Dr Felsher Marisa
marisa.felsher@merck.com - United States

Felsher Marisa
marisa.felsher@merck.com - United States

#3174

ECONOMIC AND HUMANISTIC BURDEN OF HPV-RELATED DISEASE IN INDONESIA: A QUALITATIVE ANALYSIS

39 - Low resource settings

Felsher M, Setiawan D, Varga S, Della Pia R, Riley D, Newman R, Beveridge A, Oswald C, Kothari S, Postma M

Background/Objectives: The global burden of human papillomavirus (HPV)-related disease(s) is increasing in developing countries, including Indonesia. However, few studies have explored how having a HPV-related disease impacts the lived experiences of adult men and women in Indonesia. Therefore, the objective of this analysis was to qualitatively explore the economic and humanistic burden of HPV-related disease(s) in patients from Indonesia.

Methods: Between June 2021 and September 2021, semi-structured interviews were conducted with patients diagnosed with a HPV-related disease (genital warts or cancer), identified from a range of clinical settings in Jakarta, Indonesia and surrounding regions. Clinical settings included government mandated primary care clinics and hospitals providing specialist care, such as ear nose and throat specialists, urologists and gynecologists. For further insight, health care professionals (HCPs) specialized in treating HPV-related disease(s) were also recruited for a separate semi-structured interview. This research received ethical approval from the ethical committee at Muhammadiyah Purwokerto University. Thematic analysis of qualitative interviews was used to explore the impact of HPV-related disease(s) on finances, social relationships, physical and mental health.

Results: Eighteen patients were recruited (nine men and nine women), aged 30-56 years. Patients had head/neck cancer (n=8), anogenital cancer (anal, penile, or vaginal cancers; n=4), cervical cancer (n=3), or genital warts (n=3). Ten HCPs were also recruited. Patients most frequently emphasized the humanistic burden of their HPV-related disease(s), particularly surrounding the negative impact on their emotional and mental state, and relationships with friends and family. The overall cost of HPV-related disease(s) caused a severe economic burden for patients. Patients reported alterations in physical ability and productivity due to their disease(s), leading to low or reduced income. HCPs mostly emphasized the financial impact of HPV-related disease(s) and consistently understated the emotional and societal burden of disease.

Conclusions: Patients described various ways in which HPV-related disease(s) negatively impacted their lives. There was evidence of a substantial economic burden, but the largest emphasis was on mental and social health. This research highlights the need to identify ways to reduce the negative psychosocial consequences of HPV disease(s) in Indonesia. Furthermore, increased HCP education of the broader humanistic impacts of HPV-related disease(s) may help them best support their patients.

Oral presentation

Ms Wang Wei (vivian)
wei.wang40@merck.com - United States

Wang Wei
wei.wang40@merck.com - United States

#3176

Cervical Cancer Screening Guidelines and Screening Practices in 11 Countries: A Systematic Literature Review

10 - HPV screening

Wang W, Emanuele A, Kristina H, Natalie H, Smita K, Anushua S

Background/Objectives: In 2020, World Health Organization (WHO) released a global strategy to accelerate the elimination of cervical cancer globally. One of the goals by 2030 is to achieve 70% of women screened using a high-performance test. As one of the two cervical cancer prevention measures, namely vaccination against HPV (primary prevention) and screening (secondary prevention), the screening guidelines and practice are critical but also often change to reflect scientific developments. Clinicians and policymakers need to understand the current screening guidelines and practice to support cancer prevention efforts.

Methods: A systematic literature review, complemented by targeted searches, has been conducted in peer-reviewed publications and policy documents (2005/01-2021/01) to describe cervical cancer screening guidelines and clinical practice, in 11 selected countries, namely, Australia, Canada, China, France, Germany, Italy, Japan, Spain, Sweden, the UK, and the US.

Results: A total of 84 peer-reviewed publications and 33 policy documents were included. We found that only 6 out of 11 countries assessed have an organized population-based screening program in place. Considerable differences persist across countries concerning the recommended screening intervals and the type of screening test. Methods of data collection are also heterogeneous, and systematic data collection is often not established. Due to the heterogeneity in data collection, the comparability of screening practices and rates is limited. Moreover, guideline changes are implemented with a substantial time lag in clinical practice: one to several years are needed until screening behavior is in agreement with the updated recommendations. Screening rates outside the target population (e.g. women < 20 years) are considerable and represent an indicator of discrepancies between guidelines and practice.

Conclusions: Cervical cancer screening guidelines, practices, and methods of data collection are heterogeneous globally. Future changes in guidelines and screening practice may occur as the first cohorts of vaccinated women reach screening age.

Oral presentation

Dr Pompeo Giampaolo
g.pompeo@ispro.toscana.it - Italy

Pompeo Giampaolo
g.pompeo@ispro.toscana.it - Italy

#3182

Implementation of a centralized HPV-based cervical cancer screening programme in Tuscany: first round results and comparison with the foregoing Pap-based screening programme

10 - HPV screening

Carozzi F, Burrone E, Confortini M, Pompeo G, Bisanzi S, Cellai F, Paganini I, Mantellini P, Iossa A, Lelli L, Di Pierro C, Matucci M, Cannistrà S, Rosati R, Mongia A, Visioli C, Gorini G, Sani C

Background/Objectives: In Tuscany region HPV-based cervical cancer screening programme is ongoing from 2013, with a strong level of centralization towards the Regional Laboratory for Cancer Prevention of ISPRO institute (Florence). Aims of this study were to evaluate the performance of the first round of HPV-based screening and to compare it with the previous round of Pap-based screening.

Methods: The transition from Pap to HPV-based screening started gradually in 2013 from older women and from 3 out of 12 Tuscany Local Health Units (LHUs). Data from Florence and Grosseto LHUs (about 300,000 women) were analysed and performance screening indicators were estimated.

Results: HPV-based indicators recorded a good performance, with an increased compliance vs. Pap-based programme. We registered a substantial decrease of waiting times from sampling to test reporting, probably related to the centralization strategy. Since the screening protocol was the same and since the samples were analyzed in a unique regional laboratory, we could hypothesize that the different HPV positivity (6.8% in Florence vs 8.4% in Grosseto) was due to real difference in the HPV prevalence among women of the two LHUs. HPV-based programme led to an important increase of colposcopy referral rate (4.3% vs. 1.2%) and to a significant increase of about three-fold of CIN2+ DR (8.3‰ vs. 3.4‰). The difference of the CIN2+ DR between traditional Pap test and HPV+/triage Pap test registered at baseline (respectively, 3.4‰ vs. 5.8‰ for Florence and 3.6‰ vs. 6.2‰ for Grosseto) was due to a better performance of the Pap test of triage in comparison with the Pap test as primary test.

Conclusions: The transition phase carried out in Tuscany region from Pap to HPV-based screening was well planned and the choice to introduce a strong centralization towards a unique laboratory with a long experience in this specific field, determined a higher efficiency of the screening process. Our data confirmed that HPV-based is more effective in detecting high-grade precancerous and cancerous lesions than Pap-based screening and it is characterized by an "anticipatory effect" in the detection of CIN2+ lesions. Referral to colposcopy, as expected in a prevalence round, increased significantly; thus, screening programmes that decided to move from Pap-based to HPV-based screening, should include increased resources dedicated to colposcopy services. Therefore, the management of women with a positive HPV test in successive HPV-based rounds that were negative in the previous HPV round could probably be different, since this population is characterized by a lower risk to develop a CIN2+ lesion because they are women with a recent HPV infection. Thus it is strongly necessary to start thinking how the current screening protocol could be modified in the subsequent screening rounds (incidence round), in order to reduce the risk of unnecessary colposcopy and treatments.

Oral presentation

Dr Tewari Krishnansu
ktewari@uci.edu - United States

Tewari Krishnansu S
ktewari@hs.uci.edu - United States

#3208

EMPOWER-CERVICAL 1/GOG-3016/ENGOT-CX9: RESULTS OF PHASE 3 TRIAL OF CEMIPIMAB VS INVESTIGATOR'S CHOICE CHEMOTHERAPY IN RECURRENT/METASTATIC CERVICAL CARCINOMA

08 - Immunotherapy - Immuno-oncology - New treatments

Tewari K, Monk B, Vergote I, Miller A, De Melo A, Kim H, Kim Y, Samouëlian V, Lorusso D, Damian F, Chang C, Gotovkin E, Takahashi S, Ramone D, Pikiel J, Guerra Alía E, Colombo N, Makarova Y, Li J, Jamil S, Jankovic V, Chen C, Seebach F, Lowy I, Mathias M, Fury M

Background/Objectives: EMPOWER-Cervical 1/GOG-3016/ENGOT-cx9 (NCT03257267) is an open-label, randomised (1:1), multicentre, Phase 3 trial of cemiplimab versus investigator's choice (IC) chemotherapy (chemo) in recurrent/metastatic (R/M) cervical cancer that has progressed after first-line platinum-based treatment.

Methods: Patients were enrolled regardless of programmed cell death-ligand 1 (PD-L1) expression; received cemiplimab 350 mg intravenously every 3 weeks or IC chemo (pemetrexed, vinorelbine, gemcitabine, irinotecan or topotecan), up to 96 weeks; and were stratified by histology (squamous cell carcinoma [SCC] versus adenocarcinoma or adenosquamous carcinoma [AC]). Primary endpoint was overall survival (OS), analysed hierarchically in patients with SCC followed by overall population (SCC + AC). Additional endpoints included progression-free survival (PFS), objective response rate (ORR), quality of life (QoL) and safety. Interim analysis was scheduled when 85% events occurred among patients with SCC pts.

Results: 608 patients were randomised: median age was 51.0 years (range, 22.0–87.0); 477 patients had SCC (77.8%), 131 patients had AC (22.2%) histology; and Eastern Cooperative Oncology Group performance status was 0 (46.5%) or 1 (53.5%). Median cemiplimab exposure was 15.2 weeks (range, 1.4–100.7). At interim analysis, median OS favoured cemiplimab versus chemotherapy (total population: 12.0 vs 8.5 months; hazard ratio [HR], 0.69; 95% confidence interval [CI], 0.56–0.84; one-sided $P < 0.001$; SCC population: 11.1 vs 8.8 months; HR, 0.73; 95% CI, 0.58–0.91; one-sided $P = 0.003$; AC population 13.3 vs 7.0 months; HR, 0.56; 95% CI, 0.36–0.85; nominal one-sided $P < 0.005$ [not adjusted for multiplicity]. PFS, ORR, in overall and SCC populations, and mean change from baseline QoL in the SCC population, favoured cemiplimab versus chemotherapy. The most common treatment-emergent adverse events (AEs) of any grade for cemiplimab versus IC chemo were anaemia (25.0% vs 44.5%), nausea (18.3% vs 33.4%), fatigue (16.7% vs 15.5%), and vomiting (16.0% vs 23.4%). Discontinuation due to AEs occurred in 8.7% (cemiplimab) and 5.2% (IC chemo) of patients.

Conclusions: Cemiplimab significantly improved OS over single-agent chemo for patients with R/M cervical cancer after first-line treatment regardless of histology and despite not having been selected by PD-L1 status. No new safety signals were observed.

Poster

Ms Popova Anna
asya-med@mail.ru - Russia

Ms Popova Anna
asya-med@mail.ru - Russia

#3241

Prevalence of different types of HPV HCR in cervix and anus among HIV-positive women in Moscow region

34 - Sexually transmitted diseases and HIV infection

Popova A, Domonova E, Vinogradova N, Akimkin V

Background/Objectives: HIV-infected women have a higher risk of HPV infection than HIV-negative women, and a higher risk of persistence and malignancy. A number of researchers recommend screening for HPV-associated anus disease in HIV-infected women, regardless of the results of a cervical examination. Unfortunately AIDS Centers in the Russian Federation do not have funding for diagnostics of HPV-associated diseases. Objective: to study the prevalence of human papillomavirus of high carcinogenic risk (HPV HCR) in cervix and anus among HIV-infected women in Moscow region.

Methods: 101 HIV-infected women from Moscow region were examined from August 2018 to July 2019. All women were tested for HPV-test with the determination of 14 types of HPV HCR (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68) in cervix and anus.

Results: Among the 101 HIV-women mostly young people (38.99 ± 6.52 y-s, min-26, max-58, median-58) predominated. In 60.4% cases (95%CL: 50.9 - 69.9) of HIV-infected women were diagnosed with HPV HCR. At the same time the frequency of HPV detection differed in the studied anatomical locus: in 50.8% of cases HPV was detected simultaneously in the cervix and anus, in 21.3% - only cervix and in 27.9% - only anus. All 14 HPV HCR types were diagnosed. Prevalence of different types of HPV HCR among HIV-positive women significantly depending by anatomical locus (tab.1). In cervix, 33 and 52 types of HPV were in the lead (20.5% and 18.2%, respectively). In anus - 16 and 51 types of HPV (31% and 27%, respectively). In 67.7% (95%CL: 51.3 - 84.2) HIV-women with HPV-infection the types of HPV were recorded, which were absent them in cervix during the simultaneous examination of the two studied loci in anus. At the same time, in the anus in 38.7% cases, the number of identified HPV types was greater than that found during examination of the cervical canal.

Conclusions: High incidence of HPV detection was revealed in the group of HIV-infected women from the Moscow region. All 14 HPV HCR types were diagnosed in all studied anatomical locus. The results obtained demonstrate the necessity to introduce screening for HPV-associated diseases at the AIDS centers in the Moscow region of the Russian Federation.

Oral presentation

Dr Zakhirova Nargiza
dr.zahirova@mail.ru - Uzbekistan

Zakhirova Nargiza
dr.zahirova@mail.ru - Russia

#3257

NATIONAL HPV VACCINATION PROGRAM IN THE REPUBLIC OF UZBEKISTAN

06 - HPV prophylactic vaccines

Zakhirova N, Tillyashaykhov M, Akhmedov O, Osmanova E, Saydakhmedova V, Islamova Z, Mamatova M

Background/Objectives: Cervical cancer prevention by introducing HPV vaccination in the Republic of Uzbekistan.

Methods: According to the national immunization schedule, with the support of international organizations GAVI, WHO, UNICEF and the Ministry of Health of the Republic of Uzbekistan, the HPV vaccination program for 9-year old girls was launched in 2019. Used vaccine Gardasil MK (tetravalent) 2 doses 0-6 months. Vaccinations were carried out in medical offices of public and private schools, in family polyclinics at the place of residence.

Results: From October 2019 to June 2021, 2 rounds of HPV vaccinations were carried out. In the first round, 290585 girls out of 297018 were vaccinated throughout the Republic, vaccination coverage was 97.8%. In the second round, 300893 girls were vaccinated out of 304,390 to be vaccinated, vaccination coverage was 98.8%. In total, 591,478 girls of 9 years old were vaccinated with HPV in the republic. Adverse reactions to vaccination were observed in 68 (0.1%) girls in the form of subfebrile fever, no serious complications after vaccination were observed. In November of this year, it was begun the first round of vaccination of girls 12-14 years old in the Republic.

Conclusions: In the Republic of Uzbekistan, the national HPV vaccination program for 9-year old girls was carried out successfully, with a total coverage of 98.3%.

Poster

Dr Zakhirova Nargiza
dr.zahirova@mail.ru - Uzbekistan

Dr Zakhirova Nargiza
dr.zahirova@mail.ru - Uzbekistan

#3258

PILOT SCREENING OF CERVICAL CANCER HPV TESTING IN THE REPUBLIC OF KARAKALPAKISTAN

10 - HPV screening

Akhmedov O, Osmanova E, Saydakhmedova V, Otajonov M, Mamatova M, Islamova Z

Background/Objectives: Detection of HPV associated precancerous diseases of the cervix and cervical cancer in ecologically unfavorable regions of the Republic of Karakalpakstan.

Methods: The first UNFPA-WHO pilot project on HPV-based cervical cancer screening is underway with funding from the Government of Japan, technical assistance from WHO, IARC and UNFPA, and some support from the French Embassy. The pilot screening project was approved by the Ministry of Health of the Republic, covering 10 districts, mainly near the Aral Sea: Khojeyli, Kanlykul, Shumanay, Chimbay, Karauzyak, Kungrad, Kegeili, Muynak, Buzatau, Nukus district. The age of women is 30-55, the number of women is 50,000 (100%).

Results: Interim analysis: from August to October 2021, 17,662 (35%) were tested, 915 (5.6%) of which HPV positive, 15,908 (90.1%) - HPV negative. By randomization 1/1, a triage of VIA was performed - 438 (37.9%) / colposcopy - 364 (47.8%), invalid and erroneous tests - 0.4%, by the end of the project 32,338 (65%) should be tested. 92 (10%) referred to oncologist, pathological conclusion: CIN I - 5 (0.5%), CIN II - 14 (1.5%), CIN III - 6 (0.7%), Cancer in situ - 6 (0.7%).

Conclusions: The completion of the pilot screening is scheduled for late February 2022. Upon successful completion of the project, a national cervical cancer screening program will be developed.

Oral presentation

Dr Acuti Martellucci Cecilia
cecilia.martellucci@gmail.com - Italy

Acuti Martellucci Cecilia
cecilia.martellucci@unife.it - Italy

#3267

EFFECTIVENESS OF VACCINATION RE-INVITATION AT 25 YEARS OF AGE TO INCREASE HPV VACCINE UPTAKE

06 - HPV prophylactic vaccines

Acuti Martellucci C, Morettini M, Massetti L, Flacco M, Manzoli L, Giacomini G, Catalini A, Pasqualini F

Background/Objectives: The HPV vaccine was found to be highly effective against cervical cancer (Lei et al, NEJM 2020). However, coverage of this vaccination is often lower than the 90% target, and several strategies have been implemented to try to increase vaccine uptake (Bruni et al, Preventive Medicine 2021). This study aims to assess whether re-inviting women aged 25 years was effective in increasing HPV vaccine uptake.

Methods: The Province of Ancona, Italy actively offers free HPV vaccination from age 12 years, and started with the 1996 birth cohort in 2009 (males were invited starting in 2017). This cohort was also the first to remain eligible for the vaccine until age 25 years: in 2021, unvaccinated women were actively re-invited in the context of the cervical screening campaign. Women had to send an email requesting the vaccine, and dedicated sessions were organised for the three doses (0, 2, and 6 months schedule). Data were collected from the official registries of the Local Health Agency, and the main outcome was the increase in uptake of at least two vaccine doses following the re-invitation.

Results: Out of the whole cohort of 2074 women, 66.0% were vaccinated with at least two doses in their adolescence, while the remaining (n=705) did not adhere to the vaccination and were thus actively invited for catch-up in 2021. Those requesting an appointment were 125 (17.7% of those invited). Of these, 27 did not show up for their first dose, for a final uptake of 13.9% (n=98). Thus, the overall uptake in this cohort increased from 66.0% to 70.7%.

Conclusions: Re-inviting the women who were not vaccinated up to 25 years of age increased by 5% the overall vaccine uptake, but 86% of the unvaccinated women still did not adhere. If these data are confirmed, the cost effectiveness of repeating the vaccination campaign should be carefully evaluated, as the available resources could be devoted to ensure the maximum immunization coverages among adolescents.

References: Lei J et al. HPV Vaccination and the Risk of Invasive Cervical Cancer. New England Journal of Medicine (2020) 383(14): 1340-1348. Bruni L et al. HPV vaccination introduction worldwide and WHO and UNICEF estimates of national HPV immunization coverage 2010-2019. Preventive Medicine (2021) 144: 106399.

Oral presentation

Ms Tisler Anna
anna.tisler@ut.ee - Estonia

Tisler Anna
anna.tisler@ut.ee - Estonia

#3283

WOMEN'S RIGHT TO LIFE IS NOT AGING, OLDER WOMEN HAVE TO BE SCREENED FOR CERVICAL CANCER

10 - HPV screening

Tisler A, Pärna K, Uusküla A

Background/Objectives: Estonia is a European Union country experiencing high cervical cancer mortality (4.3 / 100 000) and incidence (18.5 / 100 000) rates in 2020 despite having organised screening program from 2006. Inadequate uptake, insufficient quality of the Pap-smear based screening, and stage distribution shifting towards later stages are described before. Importantly, cervical cancer screening is targeting women up to 65 years. In 2020, 31% of incident cervical cancer cases in Estonia were reported among women ≥ 65 . High-risk HPV infection is a necessary cause for cervical cancer. The purpose of the study was to determine a population-based HPV prevalence and to describe sociodemographic and sexual behavioral patterns among women wide age range in Estonia.

Methods: A random age stratified (30-33, 57-60, 67-70) study sample comprised 3065 women representing Estonian female population. All women were invited to the study from October 2020-May 2021 using e-mail or paper invitation. Each participant or those who did not react on invitation was sent a package containing a (self-sampling) vaginal swab collection kit (Eswab, Copan), collection instructions, informed consent and questionnaire on sociodemographic, reproductive history, sexual behaviour and healthcare utilization data. HPV genotyping was made using Luminex xMAP (16, 18, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 66, 68, 82).

Results: Filled out questionnaires and samples were returned from 1347 (44%) targeted sample. Prevalence of any type of HPV infection was 19%. We observed a U-shape prevalence curve - with hrHPV prevalence of 22%, 15%, and 19% among women aged 30-33, 57-60 and 67-70 respectively. The most common HPV types detected in study population were HPV 53 (19%), 56 (16%) and 16 (14%); among older women HPV 68 (33%) and HPV 53(29%). 20% and 34% of women aged 67-70 have not been screened for the last five and three years respectively. High proportion of older women did not have higher education (60%), were having unprotected sex during their first sexual intercourse (70%) and a quarter reported being currently sexually active.

Conclusions: To our knowledge this is the largest assessment of HPV prevalence in Estonian female population. The overall HPV prevalence in our study population was high and documents a high prevalence of hrHPV among older women (aged ≥ 65) currently excluded from screening. However, these older women are at continuous risk for cervical cancer. Discontinuing screening at age 65 is premature and extension of screening age span in older age is warranted.

Oral presentation

Ms Liang Linda
linda.liang@tum.de - Germany

Klug Stefanie J.
stefanie.klug@tum.de - Germany

#3285

REASONS FOR NON-ATTENDANCE OF COLPOSCOPY FOLLOWING AN ABNORMAL CERVICAL CANCER SCREENING RESULT IN GERMANY

25 - Colposcopy

Schauberger G, Fischbeck S, Ikenberg H, Blettner M

Background/Objectives: Following an abnormal screening result, colposcopy is an important step to determine disease status and guide management. Barriers exist for women to return for colposcopy, even in high-resource countries such as Germany. We described the reasons for non-attendance within a real-world screening study.

Methods: The MARZY population-based prospective cohort study collected screening data on women aged 30-65 years in Germany. These analyses were based on screening conducted at baseline (round 1) and follow-up (round 2) screening 3 years later. Screening was conducted by cytology and Human Papillomavirus (HPV) co-testing (Hybrid Capture 2). Study colposcopy referrals included abnormal cytology (atypical squamous cells of undetermined significance or worse; ASC-US+), high-risk HPV results (hrHPV) or both. Female study personnel actively recalled women by telephone who were referred to colposcopy but did not arrange an appointment within 3 months of their screen-positive result. Reasons for non-attendance were gathered during these active recall interviews and were descriptively examined.

Results: Of 2,627 women screened at round 1, 222 were referred to colposcopy and initially 106 did not undergo colposcopy. After 29 women were motivated to make an appointment, 77 (34.7%) still did not attend despite active recall. At round 2, 107 of 2,093 screened were referred and 33 initially did not undergo colposcopy within the recommended time frame. After recall efforts, 23 (21.5%) still did not attend. Across both rounds, 89 did not attend colposcopy (28.9%) and among participants with low-grade cytological abnormalities or worse (LSIL+) and/or hrHPV, 55 did not attend (23.2%) despite recall efforts. 17.4% of participants positive in both tests (LSIL+ threshold) did not attend. Major reasons for non-attendance were lack of time (56.4%), barriers such as accessibility to the clinic (45.6%), and personal advice against study colposcopy by the office-based gynaecologist who conducted screening (43.9%). The majority of these women advised against study colposcopy had hrHPV only and reported having a repeat cytology test done. Almost a third of non-attendees reported that colposcopy clinic choice was limited and a quarter reported they had forgotten the appointment. Fear of the procedure or outcome of the appointment were reported in 15% of non-attendees.

Conclusions: Colposcopy attendance particularly in women with high-grade lesions needs improvement in Germany. Non-attendance can be addressed by improving patient education on screening results, and minimised by a systematic failsafe active recall component within the current screening program.

Prof. De La Fuente Valero Jesus
delavalero@gmail.com - Spain

García Godoy Cristina
cristina.garcia@uriach.com -

#3287

DESIGN AND OBJECTIVE OF GLUCANVIR STUDY: EFFICACY OF AN INTRAVAGINAL TREATMENT WITH CARBOXYMETHYL-B-GLUCAN AND POLYCARBOPHIL IN HIGH RISK-HPV CLEARANCE

08 - Immunotherapy - Immuno-oncology - New treatments

De La Fuente Valero J, Hernández Aguado J, Martínez Lamela E, Aguión Gálvez G, Sánchez Torres D, Sanz Espinosa E, Pérez Quintanilla A, Del Pino P, García Godoy C

Background/Objectives: Human papillomavirus (HPV) infection represents a significant source of morbidity and mortality worldwide. High-risk oncogenic HPVs cause 99.7% of cervical cancers. Most of these infections are transitory and only if they are persistent and caused by a high-risk oncogenic HPV are they an important risk factor for the development of cervical intraepithelial lesions and invasive cervical cancer. Immunity plays a key factor in eliminating HPV infection. HPV evasion of these immune defense mechanisms is critical for the persistence of the infection and leads to the development of preneoplastic lesions and ultimately to cervical cancer. Medical treatments represents a therapeutic strategy to avoid the evasion mechanisms of HPV. One of these most promising treatments is beta-glucans that seem capable of affecting the course of HPV infection.

Methods: monocentric, interventional, longitudinal, prospective, open label trial with a control group with randomized, consecutive recruitment for each group, to evaluate the safety and efficacy of an intravaginal treatment with carboxymethyl- β -glucan and polycarbophil in the clearance of High-risk Human Papillomavirus Infection in women without cytological alterations. n=200. Unvaccinated HPV+ women aged 30-40 with normal cytology were randomized into 2 groups: A) Carboxymethyl- β -glucan (Colpofix®) 1 application / day x 20 days, rest 10. Repeat 20 x 3 cycles. B) Control group: no treatment (usual clinical practice).

Results: There are more than 6000 studies that have described the effects of glucans. Recent studies focusing on their influence on cytotoxic and helper T cells, antigen presenting cells (APCs), inflammatory pathways, and oxidative burst (using reactive oxygen species to kill cells) have revealed that they may also have some anti-cancer properties. Carboxymethyl β -glucan gel treatment has been studied in 4 different studies on affected individuals by CIN1 or/and HPV+. These case-control studies demonstrated that it has an anti-cervical cancer role in CIN1 regression versus wait and see statistically significant. These studies suggested that, in addition to the anticancer effects of beta-glucans, they also have some effects on infection by HPV, the main cause of cervical cancer.

Conclusions: Data from this ongoing study will indicate if β -glucans has a therapeutic effect upon the clearance of HPV infection on the different genotypes. Given a new approach different from wait and see and would represent a significant advancement in the management of HPV positive patients.

References: ClinicalTrials.gov Id.: NCT04781023 Chaichian S, Moazzami B, Sadoughi F, Haddad Kashani H, Zaroudi M, Asemi Z. Functional activities of beta-glucans in the prevention or treatment of cervical cancer. J Ovarian Res. 2020 Mar 5;13(1):24. doi: 10.1186/s13048-020-00626-7. Review. Stentella P, Biamonti A, Carraro C, Inghirami P, Mancino P, Pietrangeli D, Votano S, Lazzari P, DE Medici C. Efficacy of carboxymethyl beta-glucan in cervical intraepithelial neoplasia: a retrospective, case-control study. Minerva Ginecol. 2017 Oct;69(5):425-430. doi: 10.23736/S0026-4784.17.04053-9. Pietrantonio E, Signore F, Berardi G, Donadio F, Donadio C. [Role of beta-glucan in the treatment of recurrent candidiasis and HPV-correlated lesions and reparative process of epidermis]. Minerva Ginecol. 2010 Feb;62(1):1-5. Italian. Scardamaglia P, Carraro C, Mancino P, Stentella P. [Effectiveness of the treatment with beta-glucan in the HPV-CIN 1 lesions]. Minerva Ginecol. 2010 Oct;62(5):389-93. Italian. Lavitola G, Della Corte L, De Rosa N, Nappi C, Bifulco G. Effects on Vaginal Microbiota Restoration and Cervical Epithelialization in Positive HPV Patients Undergoing Vaginal Treatment with Carboxy-Methyl-Beta-Glucan. Biomed Res Int. 2020 Apr 27; 2020:5476389. doi: 10.1155/2020/5476389. PMID: 32420349; PMCID: PMC7201736. Laccetta G, Carrone A, Burratti M, Mancino P. Effect of the treatment with β -glucan in women with cervical cytologic report of atypical squamous cells of undetermined significance (ASCUS) and low-grade squamous intraepithelial lesions (L-SIL). Minerva Ginecol. 2015 Apr;67(2):113-20. Epub 2015 Feb 5. PMID: 25668505

Oral presentation

Dr Wu Elise
chi-tao.wu@merck.com - United States

Wu Elise
chi-tao.wu@merck.com -

#3288

NEXT ERA OF HPV MODELS: A CLOUD-BASED USER-FRIENDLY DYNATIC TRANSMISSION MODEL

36 - Economics and modelling

Wu E, Varga S, Sharma A, Palmer C, Kothari S, Dasbach E

Background/Objectives: Human papillomavirus (HPV) infection is a well-known cause of cervical cancer in woman. In addition, HPV infection is associated with diseases in men and women including anal and head and neck cancers as well as genital warts. HPV vaccination has been available since 2006 and over time has expanded in the breadth of the disease and populations that it covers. Given this, payers and policy decision-makers are interested in better understanding of the projected public health impact and cost-effectiveness (CE) of different HPV vaccination strategies holistically and timely. PRIME model, a static policy analysis tool, represents one of the publicly available models that evaluates the benefit of HPV vaccination in women for cervical cancer. Given its limited scope, several dynamic transmission models (DTM) have been developed capture the full spectrum of HPV vaccine benefit. DTM is complex, requiring calibrations and is time-consuming. Therefore, developing an agile and flexible tool that can execute multiple scenarios in real-time and in a user-friendly fashion is needed. The objective of this exercise is to develop a cloud-based Mathematica execution framework that could run multiple complex computations in parallel and help describe and capture the full value of HPV vaccination for a broad set of stakeholders.

Methods: Three steps were taken to transform a sophisticated standalone model into an agile and user-friendly application. First, A cloud-based epidemiologic and economic tool for nonavalent HPV vaccination was developed which combines DTM of HPV infection in heterosexual men and women with economic modeling to measure the CE of various vaccination strategies. Second, the model is managed by a single secure health economic platform that provides stable execution environments for multiple modeling technologies. It is a Model Hub for model developers and consumers to collaborate around uploading inputs, managing, and accessing information of health economic models and the platform provides several capabilities around storage, advanced search, model execution, parallel processing capabilities and version control for scripted language models. Finally, a user-friendly application is developed which allow user to evaluate various HPV vaccine strategies or scenarios in real-time by updating key parameters.

Results: A cloud-based DTM application was launched in 5 pilot counties. Health economists, policy decision-makers and payers utilized the tools to evaluate HPV vaccine benefit in various scenarios and selected the optimal vaccine strategies. The application is evolving continually in both functionalities and covered countries.

Conclusions: A cloud-based agile DTM is beneficial for the HPV community to demonstrate broad value of HPV vaccines in a timely fashion in discussion with key decision maker. This tool would allow HPV researchers to evaluate the long-term benefit of HPV vaccines in various scenarios to determine the optimal vaccination programs.

Oral presentation

Mr Murphy Brendan
Brendan.murphy@merck.com - Canada

Dr Brown Vivien
vbmd@outlook.com - Canada

#3291

COLLABORATING TO SUPPORT GLOBAL ELIMINATION OF CERVICAL CANCER - THE "CANCER WON'T WAIT" PROGRAM IN CANADA

37 - Advocacy, acceptability and psychology

Palmay C, Murphy B, Mccarles J

Background/Objectives: Background: In 2018 the World Health Organization Called for the Global Elimination of Cervical Cancer (insert footnote) and invited organizations all over the world to join this movement. Feb 4th 2020 Canadian Partnership Against Cancer released the Canadian Action Plan (insert footnote). Federation of Medical Women of Canada (FMWC) approached Merck Canada Inc., Kirkland, QC, Canada to facilitate a collaboration between both organizations to increase awareness and create action to reduce/eliminate HPV related cancers in Canada. The result was the "Cancer Won't Wait" Movement. Objective: To raise awareness and encourage health care providers across all fields to become actively involved in helping Canada to achieve the CPAC goal of eliminating HPV related cancers.

Methods: Methods: FMWC and Merck Canada Inc., Kirkland, QC, Canada developed a collaboration agreement to identify roles and responsibilities. Merck Canada Inc., Kirkland, QC, Canada was responsible to use their field force to invite HCP's to join the "Cancer Won't Wait" movement and foster awareness and ongoing participation. FMWC and Scientific Experts were required to create newsletters highlighting Global, National and LOCAL initiatives that were being done to facilitate the WHO and CPAC objective.

Results: Results: In 2021 6 newsletters were created - all having a particular theme: 1) How to Pandemic Proof your Practice (identifying ways to keep vaccinating, and screening patients during COVID 2) From Local to Global (discussing Elimination initiatives across the world). 3) The Y Factor (We can not forget the males) 4) Vaccine Hesitancy and HPV (very timely with the role out of COVID vaccines) 5) Vaccination is a Team Sport (identifying the role that pharmacists can play). 6) Vaccines require collaboration (identifying the roles of schools, public health and HCP's). Despite all HCP's having COVID as a priority, 420+ HCP's have registered to join the movement with the intent of prioritizing HPV prevention within their practices. Cancer Won't Wait has been featured in the Canadian Pharmacist Journal(1), Immunize Canada Bulletin (2), and the Canadian Partnership Against Cancer(3).

Conclusions: Conclusion: The Cancer Won't Wait movement was a result of a public and private partnership. Continuing these types of collaborations will help the WHO, CPAC and other organizations achieve their Cervical Cancer Elimination Goals in their lifetime. This initiative will continue in 2022 and the goal is to reach 5000 HCP's . We encourage others to implement similar collaborations and initiatives.

References: References: Brown V, Tsuyuki RT. The pharmacist's role in prevention of HPV-related cancers. Can Pharm J (Ott). 2021 Jun 18;154(4):228-231. doi: 10.1177/17151635211019997. PMID: 34345310; PMCID: PMC8282915. Immunize Canada Bulletin - Read Issue 310 | Lire No. 310 (mailchi.mp) Canadian Partnership Against Cancer - Action plan for the elimination of cervical cancer in Canada, 2020-2030 - Canadian Partnership Against Cancer

Poster

Mr Alling Joe
joe.alling@primeglobalpeople.com - United Kingdom

Lorusso Domenica
domenica.lorusso@policlinicogemelli.it - Italy

#3293

PHASE 3 RECURRENT/METASTATIC CERVICAL CARCINOMA TRIAL: SUBGROUP EFFICACY ANALYSIS OF CEMIPIMAB VERSUS INDIVIDUAL INVESTIGATOR'S CHOICE CHEMOTHERAPY

08 - Immunotherapy - Immuno-oncology - New treatments

Lorusso D, Vergote I, Tewari K, De Melo A, Kim H, Kim Y, Lisyanskaya A, Damian F, Chang C, Rischin D, Takahashi S, Ramone D, Pikiel J, Guerra Alía E, Li J, McGinniss J, Jamil S, Mathias M, Fury M, Monk B

Background/Objectives: There is no standard-of-care regimen in the second-line setting for women with recurrent/metastatic (R/M) cervical carcinoma. Cemiplimab was recently shown to significantly improve overall survival (OS) compared with investigator's choice (IC) chemotherapy in patients with R/M cervical cancer after first-line platinum-based chemotherapy (NCT03257267; ESMO-VP-2021). We present a pre-planned exploratory subgroup analysis comparing cemiplimab to individual IC chemotherapy options.

Methods: EMPOWER-Cervical 1/GOG-3016/ENGOT-cx9 is an open-label, randomised (1:1), multi-centre, Phase 3 clinical trial of anti-programmed cell death (PD)-1 cemiplimab vs IC single-agent chemotherapy in R/M cervical cancer that has progressed after first-line platinum-based treatment. The selection of single-agent chemotherapy by the investigator (gemcitabine, pemetrexed, vinorelbine, topotecan or irinotecan) was not protocol-defined, but the regimen had to be chosen prior to randomisation. Adult females (age ≥ 18 years) were enrolled regardless of PD-ligand 1 expression and received cemiplimab 350 mg intravenously every 3 weeks or IC chemotherapy for up to 96 weeks; and were stratified by histology (squamous cell carcinoma/ adenocarcinoma or adenosquamous), geographic region (North America/Asia /rest of world), prior bevacizumab, and ECOG performance status (0/1). Primary endpoint was OS. Additional endpoints included progression-free survival (PFS), objective response rate (ORR), duration of response, quality of life and safety. Data cutoff was 4 January 2021.

Results: A total of 608 patients were randomised: 304 to cemiplimab and 304 to IC chemotherapy (gemcitabine, n=121; pemetrexed, n=111; vinorelbine, n=32; topotecan, n=21; irinotecan, n=19) across geographic regions and histologies. Median duration of study follow-up (range) was 4.8 months (0.0–25.9) for the overall population. At second interim analysis, the trial was stopped early for efficacy. OS demonstrated improvements with cemiplimab vs each IC chemotherapy treatment—HR (95% CI) vs: pemetrexed, 0.71 (0.52–0.98); topotecan, 0.78 (0.31–1.96); irinotecan, 0.69 (0.28–1.70); gemcitabine, 0.76 (0.54–1.06); vinorelbine, 0.77 (0.40–1.48)—similar to those observed with cemiplimab vs pooled IC chemotherapy. Results were similar with PFS—HR (95% CI) vs: pemetrexed, 0.70 (0.52–0.94); topotecan, 0.90 (0.43–1.90); irinotecan, 0.77 (0.34–1.72); gemcitabine, 0.73 (0.54–0.97); vinorelbine, 1.21 (0.70–2.09). ORR were consistently higher with cemiplimab—% vs: pemetrexed, 16.0 vs 6.3; topotecan, 15.0 vs 4.8; irinotecan, 23.1 vs 15.8; gemcitabine, 17.6 vs 5.0; vinorelbine, 9.7 vs 6.3.

Conclusions: Improvements in OS, PFS and ORR with cemiplimab trended consistently with the results for the overall population regardless of IC chemotherapy drug.

Oral presentation

Dr Felsher Marisa
marisa.felsher@merck.com - United States

Chen Ya-ting
yating_chen@merck.com - United States

#3299

Assess the impact of Pharmacy-based HPV vaccination among adult men and women in the United States: A Conceptual Framework for Study Design

06 - HPV prophylactic vaccines

Felsher M, Saxena K, Bluml B, Brock K, Little J, Chen Y

Background/Objectives: Despite the Advisory Committee on Immunization Practices recommendation for HPV vaccinations up to 45 years of age in men and women in the United States, vaccination coverage among adults remain low. Pharmacies are important settings for HPV vaccination among adults due to their population reach and infrastructure for vaccine delivery. During the COVID-19 pandemic pharmacies became routine settings for COVID-19 testing and vaccine delivery, resulting in increased patient perception that pharmacies are accessible options for future vaccinations. To build on the broadening role of pharmacies, we aim to develop a pharmacist-led HPV vaccination program to identify if coadministration of HPV vaccine with influenza and/or COVID-19 vaccine, commonly administered in pharmacy settings, will impact HPV vaccination uptake and series completion among adults.

Methods: This prospective multisite study will be implemented in US pharmacies representing many practice types (e.g., community, chain) and geographic locations, and will be first piloted within five pharmacies prior to scaling up to more pharmacies if deemed acceptable and feasible. The study will have three phases: baseline pre-intervention evaluation and program development, intervention implementation, and post-intervention evaluation. During the pre-intervention phase, participating pharmacies' baseline HPV vaccination rates in the preceding 12 months will be identified through pharmacy database records. Pharmacy staff will be interviewed to inform the design and scalability of the pharmacist-led HPV vaccination program. The intervention phase will consist of educating participating pharmacists on HPV related diseases and cancers, communication strategies for offering HPV vaccine, and ways to identify patient eligibility for HPV vaccine, such as via state immunization information systems (IIS). Over a 12-month period, pharmacists will administer HPV vaccine to consenting patients, document administered vaccines and offer dose reminder services. Patient surveys will be administered to collect socio-demographic characteristics, health attitudes and behaviors. The post-intervention phase will evaluate changes in the number of vaccines administered pre- and post-implementation, as well as patient- and pharmacy-level factors associated with HPV vaccine uptake and series completion. Exit interviews with participating pharmacists will determine acceptability and feasibility of the pilot phase, and the study will be expanded to the remaining sites.

Results: NA

Conclusions: Appropriately designed study to assess pharmacist-led initiatives can facilitate access to HPV immunization services for adult populations.

Poster

Dr Wu Ying Hui
lily.wu@merck.com - Taiwan

Wu Ying-hui Lily
lily.wu@merck.com - Taiwan

#3316

HEALTH IMPACT AND COST-EFFECTIVENESS ANALYSIS OF GENDER-NEUTRAL 9-VALENT HUMAN PAPILLOMAVIRUS VACCINATION IN TAIWAN

36 - Economics and modelling

Chou H, Chu P, Wu Y, Feng C, Wu E, Sukarom I, Pavelyev A

Background/Objectives: Human papillomavirus (HPV) can cause cancers and HPV diseases in both genders. Most prevalent HPV cancer in women is cervical cancer while the most prevalent HPV cancer in men is head and neck (H&N) cancer globally. Many countries have transitioned from female-only vaccination (FOV) to gender-neutral vaccination (GNV), directly protecting men against HPV infection and diseases and possibly further reducing HPV-related diseases in women. Since December 2018, the Taiwanese government has provided the two-dose regimen of the bivalent HPV vaccine (2vHPV) free of charge to 13-year old girls as part of the National Immunization Program (NIP). The 9-valent HPV vaccine (9vHPV) provides additional benefits through direct protection against seven additional HPV types. This study assesses the health impact and cost-effectiveness of 9vHPV GNV vs. 2vHPV FOV strategies in Taiwan.

Methods: Using Taiwanese data, a validated dynamic transmission model was used to simulate the natural history of HPV infections and estimate the cost associated with HPV-related diseases. HPV-associated health outcomes included cervical lesions (CIN-1/2/3), cervical, vaginal, vulvar, H&N, anal, and penile cancers as well as genital warts (GWs) and recurrent respiratory papillomatosis (RRP). Costs, quality-adjusted life-years (QALY), and incremental cost-effectiveness ratio (ICER) were estimated. The model assumed lifelong duration of vaccine protection, herd immunity, a discount rate of 3% for costs and QALYs.

Results: Compared to 2vHPV FOV strategy, 9vHPV GNV strategy would prevent more cases of HPV-related diseases and cancers (15,860 cervical cancer, 271,753 CIN-1/2/3, 132 vaginal, 822 vulvar, 13,885 H&N, 775 anal and 181 penile cancer, 1,708,978 GWs, and 9,873 RRP) and death (8,899 cervical cancer, 54 vaginal, 357 vulvar, 8,799 H&N, 364 anal and 102 penile cancer, and 438 RRP). 9vHPV GNV strategy resulted in a 11% reduction in disease management cost as compared to 2vHPV FOV strategy. The ICER of 9vHPV GNV strategy vs. 2vHPV FOV strategy was NTD 391,360 /QALY.

Conclusions: A 9vHPV GNV strategy for 13-year old girls and boys would have additional public health and economic impact and would be highly cost-effective as compared to the current 2vHPV FOV strategy, relative to per capita GDP, which is estimated at NTD 833,447 for Taiwan. The 9vHPV GNV approach would not only help to accelerate the timeline in achieving cervical cancer elimination goal but also to reduce the incidence HPV related H&N cancer. Since Taiwanese data were used in this analysis, the results can be used as a fundamental reference to better evaluate cancer prevention policy in the future.

Oral presentation

Dr Li Mingzhu
mingzhu1815@163.com - China

Dr Li Mingzhu
mingzhu1815@163.com - China

#3317

The value of Pax1 methylation in predicting pathologic upgrading after cervical conization

18 - Methylation

Zhao C, Zhao Y, Li J, Wei L

Background/Objectives: Cold knife conization (CKC) is one of the main methods to treat high-grade cervical lesions. Unexpected early-stage cervical cancer is sometimes incidentally found after CKC, but there are no good predictive markers. We investigated the ability of PAX1 methylation to predict pathologic upgrading after cervical conization.

Methods: A total of 251 women were enrolled in the study, the mean age was 41.7 ± 10.8 years. Quantitative methylation-specific PCR (QMSP) for PAX1 methylation were collected before cervical conization.

Results: PAX1 methylation increased with the severity of the cervical lesion with CIN2(9.5%), CIN3(38.9%), AIS (23.0%) and early-stage cervical cancer(ESCC) (87.0%). PAX1 hypermethylation had more higher risk (95%CI:81.7-95.7) than cytology of LSIL (95%CI: 60.9-78.6) above and HPV16/18+ (95%CI: 69.9-85.3) in predicting CIN3+.The optimal cutoff value in predicting upgrade pathology after CKC was 7.1 and the area under the curve (AUC) was 0.88 (95% CI: 0.84-0.95, $P < 0.001$). The adjusted odds ratio (OR) of PAX1 methylation with $\Delta Cp \geq 7.1$ was significantly higher (OR24.9, 95% CI 7.6-81.2) than cytology of LSIL and above (OR3.5, 95% CI 1.1-11.2)and HPV16/18 (OR2.3, 95% CI 0.7-7.2) in predicting ESCC. There was no significant difference in PAX1 methylation status with tumor stage, depth, width of invasion, LVSI and margin status of ESCC($P \geq 0.05$).

Conclusions: PAX1 hypermethylation level($\Delta Cp \geq 7.1$) may predict pathologic upgrading after cervical conization.

Oral presentation

Dr Brautigam Karen
karen.braeutigam@uksh.de - Germany

Bräutigam Karen
Karen.Braeutigam@uksh.de - Germany

#3318

FREQUENCY, DISTRIBUTION, AND CORRELATION OF HPV IN DIVERSE ANOGENITAL AND ORAL SAMPLES

09 - HPV testing

Bräutigam K, Meier S, Meneder S, Polack S, Köster F, Rody A, Baum S

Background/Objectives: Human papilloma virus (HPV)-associated cancers increased world-wide in the last years. In addition to cervical cancer (CC), where HPV infections are attributable in >90 % of all cases, also other anogenital and oropharyngeal cancers are related to HPV infection. Furthermore, a high association with HPV was reported for intraepithelial neoplasia (INs) of the different anogenital areas.

Methods: In this study cervical, vaginal, vulvar, anal, and oral samples were collected from 509 women visiting our dysplasia consultation. DNA was isolated and full HPV genotyping (18 high risk [HR] and 12 low risk [LR] HPV types) was performed using EUROArray HPV.

Results: HR HPV (+/- LR HPV) was detected in 61.6% of cervical, 62.5% of vaginal, 64.3% of vulvar, 60.4% of anal, and in 14.6% of oral samples. HPV 16 showed the highest prevalence in all areas (cervix 45.7%, vagina 50.0%, vulva 53.9%, anus 54% and oral 79.7%). Besides HPV 16, HPV 31 was found most frequently (17.3%) followed by HPV 52 (12.0%), HPV 53 (11.3%), and HPV 18 in 9.0% in cervical, HPV 31 (16.0%), HPV 53 (13.8%), HPV 52 (12.6%), and HPV 39 (9.4%) in vaginal, HPV 53 (17.7%), HPV 31 (15.8%), HPV 52 (10.8%), and HPV 51 (8.7%) in vulvar, HPV 53 (18.9%), HPV 31 (11.3%), HPV 73 (10.3%), and HPV 51 (9.6%) in anal as well as HPV 53 (14.9%), HPV 51 (9.5%), and HPV 31 and 73 (4.1% each) in oral samples. A substantial to almost perfect concordance for most of the HPV types was examined in the cervical, vaginal and vulvar samples, and moderate to substantial concordance between those and anal samples. A good correlation was shown for high percentage of HR HPV infection in CC and HSIL (high squamous intraepithelial lesion) (CIN2/3) compared to lower HR HPV infection in LSIL (low SIL) (CIN 1) and CIN 0/normal samples.

Conclusions: As anticipated, HPV 16 showed the highest prevalence in all investigated areas. In cervical and vaginal samples, HPV 31 is frequent, while HPV 18 was detected rarely in all areas. Interestingly, HPV 53 could often be detected in different areas. As expected, the prevalence of HPV and HR-HPV infections were higher in women with precancerous lesions (CIN/VaIN /VIN ≥ 2) and carcinoma compared to women with normal cytology.

Oral presentation

Dr Macdonald Iain
iain.macdonald@mu.ie - Ireland

Macdonald Iain
iain.macdonald@mu.ie - Ireland

#3319

RAISING HPV VACCINE AWARENESS AND ADVOCACY THROUGH STEAM WORKSHOPS

38 - Health education

Macdonald I, Healy C, Malone E, Firth R, Mcdermott A

Background/Objectives: HPV Immunity Community seeks to find new ways to educate and improve Human Papilloma Virus (HPV) vaccination awareness and advocacy amongst school children. The project brought together Irish and Scottish academics from Design, Biology, and Education, and the Irish Cancer Society (ICS), in an interdisciplinary, intersectoral and international collaboration. Our aim was to improve students' understanding and awareness of, and advocacy for, vaccination, and counter misinformation, about the HPV vaccine. HPV vaccines can save lives and eradicate associated cancers, but the uptake of the vaccine among first-year second level schoolgirls in Ireland, to whom it is offered for free, has dropped from 87% in 2020 to 50% in the space of two years (ICS 2021). The focus and prioritisation of the recent COVID-19 vaccination roll-out has contributed to the reduced uptake.

Methods: A qualitative arts-based research methodology was adopted and embedded in a series of workshops, delivered consecutively, over one week. The innovative STEAM approach used drama in education activities, storytelling through artwork and video creation, and focus group interviews. The workshops facilitated a dialogue amongst the participants to co-create localised, culturally inclusive, and scientifically informed stories. Following ethical approval the study involved 16-17 year olds (n=20) from three Irish post-primary schools in County Kildare, Ireland. Using the principles of Storytelling the students developed concept boards, scripts and storyboards through an iterative process of presentations and idea selection in a visual thinking methodology (Avgerinou and Pettersson 2020). COVID restrictions forced a blended delivery with a mix of in-person and parallel online collaborative participation from colleagues in Scotland, and contributors from ICS. We evaluated the success and impact of our programme on 3 key criteria: their knowledge of immunology, their confidence in expressing their knowledge about immunology, and their confidence in advocating for vaccination and countering misinformation.

Results: Analysis of the data drew out 5 key insights: people no longer fear the serious diseases due to the success of the immunisation programmes, a lack of personal research and open discussion results in there being a poor awareness of HPV vaccine, a STEAM approach was successful at changing students' attitudes towards the HPV vaccine from passive to positive, combining creative methods with scientific concepts leads to meaningful change in human behaviour, HPV vaccine advocates were created through the use of a STEAM approach to health education.

Conclusions: The programme has empowered the young people involved to become advocates for HPV and other vaccines in their peer group and community and hence we might anticipate their health will benefit. More importantly the clear positive impact of the programme shows us that these methods hold the promise of being a template for how to positively impact the health trajectory of others across inter-generations within schools, families and as future parents. We are currently working with ICS to get wider attention and notice for our methods in the hope of greatly extending its reach and potential. As we do this we will educate and protect the participants from the cancers prevented by HPV vaccination and also help insulate these communities from the growing tide of health misinformation.

References: Avgerinou, M. D., & Pettersson, R. (2020). Visual literacy theory: Moving forward. In Handbook of Visual Communication (pp. 433-464). London: Routledge. IRC (2021) <https://www.cancer.ie/cancer-information-and-support/cancer-information/about-cancer/causes-of-cancer/hpv>

Oral presentation

Ms Pedersen Birgitte
birgitte.toennes.pedersen@regionh.dk - Denmark

Ms Pedersen Birgitte Tonnes
birgitte.toennes.pedersen@regionh.dk - Denmark

#3322

HPV SELF-SAMPLING IN THE CAPITAL REGION OF DENMARK - NEW OPERATIONAL EXPERIENCES

13 - Self-sampling

Pedersen B, Pedersen H, Serizawa R, Ejegod D, Bonde J

Background/Objectives: The Danish cervical cancer screening program is a free-of-charge cancer prevention program for all Danish resident women between 23 and 65 years. The coverage is 73%, but screening attendance is slowly declining. Notwithstanding, almost half of all newly diagnosed cervical cancers are found among screening non-attenders. To increase screening attendance, the Capital Region of Denmark implemented HPV self-sampling as an alternative offer to women not attending the regular screening offer. Operational experiences from the set-up of the self-sampling offer are presented here.

Methods: This was an opt-in offer to 57,717 screening non-attending women in 2017-2018. The women received an invitation letter and could opt-in by letter, phone, e-mail or webpage. Invitation and return-of-kit reminders were used in the set-up. HPV positive women were recommended to go to their General Practitioner (GP) for a follow-up sample. HPV negative women returned to the ordinary screening program.

Results: Of all invited women, 27% opted in. The purpose designed web-platform was the most frequent used method of response, 63% opted in by the HPV-self sampling webpage. A significant association between age and response methods was found ($p < 0.0001$). Women aged 27-29 years and 30-39 years used the webpage more often compared to women above the age of 40 years, whereas women above the age of 50 years more often used letter for registration compared to the other age groups. Use of invitation and return-of-kit reminders generated 8.6% and 6.1% additional responses and participation, respectively. Overall, 17% returned the HPV self-sampling kit for analysis. In addition, 14% had a regular clinician collected screening sample after receiving the invitation for self-sampling, leading to a total screening of 31% women after invitation. HPV prevalence was 15%, amongst those, 92% adhered to the recommended follow-up.

Conclusions: The frequent use of the HPV self-sample website when responding to the invitation and the diversity in response method dependent on the age of the invited woman emphasizes the benefit of not only novel technology, but also a flexibility in response methods to meet different preferences in the screening population. The additional participation generated by an active reminder strategy underlines the importance of timely communication in screening programs. The findings from this study are useful in the future optimization of cervical cancer screening by HPV self-sampling.

Oral presentation

Dr Scibior-bentkowska Dorota

d.scibior-bentkowska@qmul.ac.uk - United Kingdom

Dr Scibior-bentkowska Dorota

d.scibior-bentkowska@qmul.ac.uk - United Kingdom

#3331

LONGITUDINAL STUDY OF VAGINAL MICROBIOME PRE AND POST TREATMENT IDENTIFIES BIOMARKERS FOR CERVICAL INTRAEPITHELIAL NEOPLASIA 3 (CIN3).

19 - Microbiome

Scibior-bentkowska D, Cristiana B, Belinda N

Background/Objectives: Increasing evidence suggests vaginal dysbiosis is associated with persistence of human papillomavirus (HPV) infection and cervical intraepithelial neoplasia (CIN 1-3) development. In this pilot longitudinal study we aimed to investigate the potential of vaginal microbiome biomarkers to predict CIN3 development in high risk HPV (hr-HPV) positive women.

Methods: A total of 59 women with normal cytology at initial screening and follow up time over six years were enrolled from the ARTISTIC randomised trial. The cohort included 14 HPV negative and 15 hr-HPV positive women through whole follow up time. Additionally 30 hr-HPV positive women, who developed CIN3 at the first follow-up, then surgically treated for the disease were testing HPV-negative were also included. Exfoliated cervical specimens were used as starting material for whole genomic and bacterial DNA extraction. Vaginal microbiota composition was determined by 16S rRNA gene fragments sequencing on Ion Chef Instrument. The S5 methylation classifier assays were performed as previously described (Brentnall et al, 2015).

Results: We identified unique microbial biomarkers associated with CIN3 development and recovery after surgical treatment. Hr-HPV positive women with CIN3 showed a significant overrepresentation of following microbial species: *Sneathia amnii*, *Megasphaera genomosp.*, *Peptostreptococcus anaerobius* and *Achromobacter spanius*. However, *Sneathia amnii* was the only bacteria consistently associated with CIN3 in all group comparisons performed ($p < 0.01$). Conversely, after successful treatment women were hr-HPV negative and exhibited an increased representation of *Lactobacillus* species, especially *Lactobacillus gasseri* ($p < 0.01$). Higher proportions of *Lactobacillus helveticus*, *Lactobacillus suntoryeus* and *Lactobacillus vaginalis* showed a potential protective role against CIN3 development in women with persistent hr-HPV infection.

Conclusions: We confirmed in our study S5 classifier scores are increasing with cervical disease severity. Increasing *Sneathia amnii* abundance was directly proportional to the S5 score increase during cervical disease development. This result might indicate its possible role in modifying the epigenetic landscape of the cervicovaginal space. Further investigations are required to establish a link between the identified potential vaginal microbiome biomarkers and their influence on epigenetic mechanisms.

References: Banila C, Lorincz AT, Scibior-Bentkowska D, Clifford GM, Kumbi B, Beyene D, Wheeler CM, Cuschieri K, Cuzick J, Nedjai B. Clinical performance of methylation as a biomarker for cervical carcinoma in situ and cancer diagnosis: A worldwide study. *Int J Cancer*. 2022 Jan 15;150(2):290-302. Nedjai B, Reuter C, Ahmad A, Banwait R, Warman R, Carton J, Boer S, Cuzick J, Lorincz AT. Molecular progression to cervical precancer, epigenetic switch or sequential model? *Int J Cancer*. 2018 Oct 1;143(7):1720-1730. Cadman L, Reuter C, Jitlal M, Kleeman M, Austin J, Hollingworth T, Parberry AL, Ashdown-Barr L, Patel D, Nedjai B, Lorincz AT, Cuzick J. A Randomized Comparison of Different Vaginal Self-sampling Devices and Urine for Human Papillomavirus Testing-Predictors 5.1. *Cancer Epidemiol Biomarkers Prev*. 2021 Apr;30(4):661-668. Ramírez AT, Sánchez GI, Nedjai B, Agudelo MC, Brentnall AR, Cuschieri K, Castañeda KM, Cuzick J, Lorincz AT; ASC-US-COL Trial Group. Effective methylation triage of HPV positive women with abnormal cytology in a middle-income country. *Int J Cancer*. 2021 Mar 15;148(6):1383-1393. Hernández-López R, Lorincz AT, Torres-Ibarra L, Reuter C, Scibior-Bentkowska D, Warman R, Nedjai B, Mendiola-Pastrana I, León-Maldonado L, Rivera-Paredes B, Ramírez-Palacios P, Lazcano-Ponce E, Cuzick J, Salmerón J; FRIDA Study Group. Methylation estimates the risk of precancer in HPV-infected women with discrepant results between cytology and HPV16/18 genotyping. *Clin Epigenetics*. 2019 Oct 12;11(1):140. Louvanto K, Aro K, Nedjai B, Bützow R, Jakobsson M, Kalliala I, Dillner J, Nieminen P, Lorincz A. Methylation in Predicting Progression of Untreated High-grade Cervical Intraepithelial Neoplasia. *Clin Infect Dis*. 2020 Jun 10;70(12):2582-2590. Brentnall AR, Vasiljevic N, Scibior-Bentkowska D, Cadman L, Austin J, Cuzick J, Lorincz AT. HPV33 DNA methylation measurement improves cervical pre-cancer risk estimation of an H

Poster

Dr Nyanchoka Linda
Linda.Nyanchoka@kreftregisteret.no - Norway

Dr Nyanchoka Linda
Linda.Nyanchoka@kreftregisteret.no - Norway

#3334

UNDERSTANDING FACILITATORS AND BARRIERS FOR FOLLOW-UP AFTER AN ABNORMAL CERVICAL CANCER-SCREENING EXAM AMONG WOMEN LIVING IN REMOTE AREAS OF ROMANIA: A QUALITATIVE STUDY

11 - Screening for women difficult to reach

Turda I, Damian A, Nygård M

Background/Objectives: As of 2020 in Europe, Romania has the second highest incidence and mortality rates of cervical cancer, after Montenegro. In efforts to address cervical cancer in the country, the Romanian government established a national cervical cancer-screening programme in 2012. The programme provides free testing as well as free treatment in the event of cervical pre-cancer diagnosis for women 25-64 years old with health insurance who are referred from a programme-registered general practitioner. Participation in screening, re-testing and follow-up of detected precancerous lesions is suboptimal, thus reducing the overall effectiveness of the programme. The overall aim of the study is to examine facilitators and barriers to cervical cancer screening follow-up after an abnormal cervical cancer-screening exam among underserved women living in remote areas of Romania.

Methods: We will conduct an exploratory qualitative study using semi-structured interviews. Data analysis will be based on the thematic analysis outlined by Braun and Clarke. We will use QSR International's NVivo 12 as the qualitative data analysis software for both data management and analysis.

Results: Preliminary findings to be updated. The initial coding framework for our analysis will be inductively derived from the data. In this sense, our approach will include bottom-up development of analytic categories and themes. We will use QSR International's NVivo 12 as the qualitative data analysis software for both data management and analysis.

Conclusions: Study findings will inform recommendations for the Romania national policy for the cervical cancer prevention programme, with a particular focus on underserved women living in remote areas with limited access to health care services.

Poster

Ms Popova Anna
asya-med@mail.ru - Russia

Dr Domonova Elvira
elvirakuznetsova@mail.ru - Russia

#3336

Self-sampling as a cervical screening among HIV-positive women in Moscow region

13 - Self-sampling

Popova A, Vinogradova N, Akimkin V

Background/Objectives: HIV-infected women are monitored in AIDS centers in the Russian Federation. Unfortunately, the AIDS Centers in the Russian Federation do not diagnose HPV-associated diseases. HIV-infected women are forced to go to general medical facilities for cervical screening, which they often do not do for a number of reasons (stigma, lack of free time, disclosure of the diagnosis). Objective: to study the effectiveness of using self-sampling at the first stage of cervical screening of HIV-infected women in Moscow region.

Methods: 100 HIV-infected women from Moscow region were examined from February 2020 to May 2021. All women were tested for HPV-test with the determination of 14 types of HPV HCR (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68). Test material: smear of epithelial cells of the vagina and cervical canal. All women completed the questionnaire (results were assessed on a ten-point scale, where hard = 0 and easy = 10).

Results: Among the 100 HIV-women mostly young people (37.74 ± 6.3 y-s, Me - 38, min - 19, max - 57, 95%CL: 37.7-38.9). Route of HIV infection: 72.6% - sexual route, 23% - when using intravenous psychoactive substances, 4.4% - could not answer. HIV-women have CD4+ lymphocytes 617.7 ± 258.4 cells / μ l. 96.4% of women received ART. In 43% cases (95%CL: 33.3 - 52.7) of HIV-infected women were diagnosed HPV of using self-sampling. In 36% cases (95%CL: 26.6 - 45.4) were diagnosed HPV in cervix. Results of the questionnaire: ease of taking a smear - 9.4, accessibility of understanding and ease of following instructions - 9.5, comfort of using the kit - 8.9 and only 15% of respondents would prefer a traditional visit to a gynecologist.

Conclusions: self-sampling is easy and comfortable to use for HIV-infected women. self-sampling efficiently detects HPV infection in a situation of limited funding. Using self-sampling at the AIDS center is the only way to increase participation in cervical screening for HIV-infected women in the Moscow region of the Russian Federation.

Poster

Prof. Dr Zivadinovic Radomir
zivadinovicrasa@gmail.com - Serbia

Prof. Dr Zivadinovic Radomir
zivadinovicrasa@gmail.com - Serbia

#3348

Does the prevalence and genotype distribution of HPV oncologic potential change over time?

09 - HPV testing

Zivadinovic R, Zivadinovic A, Petric A, Zivadinovic L, Zivadinovic J

Background/Objectives: The distribution of HPV oncogenic potential is changing due to virus mutations, seroprevalence for a specific virus, or alternations in our immune response to the virus. There are more than 118 detected types of HPV in anogenital region. Clinical experience has shown their different oncogenic potential. So, they are classified as: high-risk 16, 18, 31, 33, 39, 45, 51, 52, 56, 58, 59, 68, 73, 82, probably high-risk 26, 53, 66, and low-risk 6, 11, 40, 42, 43, 44, 54, 61, 70 (1). Within the group of high-risk types, HPV types 16 and 18 are the most oncogenic ones since the risk of progression and cancer development for these two types is twice as high as for other types from this group (2).

Methods: Statistical Hi kvadrat test

Results: Oncogenic HPV potential also depends on geographic region, race or ethnicity, religion The first research on oncogenic potential and geographic tropism in south-eastern Serbia was performed in 2017. Types 56 and 58 were then found to be predominant as specific tropism for this region. Further research conducted until 2000 have shown changes in oncogenic distribution in relation to the year of the research. Each year a new type is identified in the top 10, and then it disappears in the years to come. In our region these types are 56, 54, 59, 39. If we rule out these types, then a regularity in oncogenic potential distribution can be observed. Top 7 HPV types according to their oncogenic potential are as follows: 16, 31, 58, 33, 18, 45 and 51.

Conclusions: Understanding of oncogenic potential distribution is important for: the assessment of oncogenic risk and the need for personalized, more aggressive diagnostic and therapeutic approach; triage of HPV positive tests and cytology-negative HPV + tests: the assessment and selection of the most effective vaccine in one geographic region

References: 1. Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, Snijders PJ, Peto J, Meijer CJ, Muñoz N. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. J Pathol. 1999; 189: 12-19. 2. Thomas C. Evaluation of HPV 16 and 18 genotyping for the triage of women with high risk HPV pos. Cytology neg. 2011; 136(4): 578-86.

Poster

Mr Del Pino Pablo
pablo.delpino@uriach.com - Spain

Mr Del Pino Pablo
pablo.delpino@uriach.com - Spain

#3349

DESIGN AND OBJECTIVE OF GLUCANCIN STUDY: Efficacy of Intravaginal Carboxymethyl- β -glucan and Polycarbophil on Low-grade Cervical Lesions

08 - Immunotherapy - Immuno-oncology - New treatments

Hernández Aguado J, Nozaleda G, Del Pino P

Background/Objectives: Human papillomavirus (HPV) infection represents a significant source of morbidity and mortality worldwide. High-risk oncogenic HPVs cause 99.7% of cervical cancers. Most of these infections are transitory and only if they are persistent and caused by a high-risk oncogenic HPV are they an important risk factor for the development of cervical intraepithelial lesions and invasive cervical cancer. Immunity plays a key factor in eliminating HPV infection. HPV evasion of these immune defense mechanisms is critical for the persistence of the infection and leads to the development of preneoplastic lesions and ultimately to cervical cancer. Medical treatments represent a therapeutic strategy to avoid the evasion mechanisms of HPV. One of these most promising treatments is beta-glucans that seem capable of affecting the course of HPV infection.

Methods: Multicentric, Interventional, longitudinal, prospective, open-label trial with a control group with randomized, consecutive recruitment for each group, to evaluate the safety and efficacy of carboxymethyl-beta-glucan and polycarbophil as a treatment for low-grade CIN associated with HR-HPV infection. n=200. Unvaccinated women aged 30-50 with LSIL/CIN1 histological results on cervical biopsy preceded by HR-HPV+ test were randomized into 2 groups: A) Carboxymethyl- β -glucan (Colpofix®) once/day for 3 months B) Control group: no treatment (usual clinical practice).

Results: There are more than 6000 studies that have described the effects of glucans. Recent studies focusing on their influence on cytotoxic and helper T cells, APCs, inflammatory pathways, and oxidative burst (using reactive oxygen species to kill cells) have revealed that they may also have some anti-cancer properties. Carboxymethyl β -glucan gel treatment has been studied in 4 different studies on affected individuals by CIN1 or/and HPV+. These case-control studies demonstrated that it has an anti-cervical cancer role in CIN1 regression vs wait and see statistically significant. These studies suggested that, in addition to the anticancer effects of beta-glucans, they also have some effects on infection by HPV, the main cause of cervical cancer.

Conclusions: Data from this ongoing study will indicate if β -glucans have a therapeutic effect on the Regression of Low-grade Cervical Intraepithelial Lesions on the different genotypes. Given a new approach different from wait and see and would represent a significant advance in the management of HPV-positive patients.

References: ClinicalTrials.gov Id.: NCT04771546 Chaichian S, Moazzami B, Sadoughi F, Haddad Kashani H, Zaroudi M, Asemi Z. Functional activities of beta-glucans in the prevention or treatment of cervical cancer. J Ovarian Res. 2020 Mar 5;13(1):24. doi: 10.1186/s13048-020-00626-7. Review. Stentella P, Biamonti A, Carraro C, Inghirami P, Mancino P, Pietrangeli D, Votano S, Lazzari P, DE Medici C. Efficacy of carboxymethyl beta-glucan in cervical intraepithelial neoplasia: a retrospective, case-control study. Minerva Ginecol. 2017 Oct;69(5):425-430. doi: 10.23736/S0026-4784.17.04053-9. Pietrantoni E, Signore F, Berardi G, Donadio F, Donadio C. [Role of beta-glucan in the treatment of recurrent candidiasis and HPV-correlated lesions and reparative process of epidermis]. Minerva Ginecol. 2010 Feb;62(1):1-5. Italian. Scardamaglia P, Carraro C, Mancino P, Stentella P. [Effectiveness of the treatment with beta-glucan in the HPV-CIN 1 lesions]. Minerva Ginecol. 2010 Oct;62(5):389-93. Italian. Lavitola G, Della Corte L, De Rosa N, Nappi C, Bifulco G. Effects on Vaginal Microbiota Restoration and Cervical Epithelialization in Positive HPV Patients Undergoing Vaginal Treatment with Carboxy-Methyl-Beta-Glucan. Biomed Res Int. 2020 Apr 27; 2020:5476389. doi: 10.1155/2020/5476389. PMID: 32420349; PMCID: PMC7201736. Laccetta G, Carrone A, Burratti M, Mancino P. Effect of the treatment with β -glucan in women with cervical cytologic report of atypical squamous cells of undetermined significance (ASCUS) and low-grade squamous intraepithelial lesions (L-SIL). Minerva Ginecol. 2015 Apr;67(2):113-20. Epub 2015 Feb 5. PMID: 25668505

Oral presentation

Mr Wei Ye
weiyel02021@163.com - China

Ye Wei
weiyel02021@163.com -

#3372

Single-cell Profiling Analysis Reveals that Targeting CXCR4+ CD8 T cells Promotes anti-PD-1 Immunotherapy of Cervical Cancer

26 - Cervical neoplasia

Canhui C, Ye W, Shitong L, Wenhua Z, Ting P, Chao H, Tian C, Bingham L, Wencheng D, Ding M, Peng W

Background/Objectives: Since anti-programmed cell death 1 (PD-1) immunotherapy has limited effect in cervical cancer, alternative co-target pathways in the tumor immune microenvironment (TIME) are underscoring. CXC chemokine receptor 4 (CXCR4) blockade facilitated immunotherapy in many cancer types. However, the mechanisms of targeting CXCR4 underlying tumor infiltration remain to be determined.

Methods: Single-cell transcriptome data of skin, liver, lung, prostate, colon, cervix, uterus, cervical cancer, endothelial cells, stem cells, pancreatic progenitor cells, nasal airway epithelium, breast epithelium, peripheral blood mononuclear cells (PBMCs), T cells, B cells, CD34+ cells, and CD14+ cells were analyzed to identify the specific expression of CXCR4 in CD8 T cells. In addition, RNA-seq data from the TCGA were integrated to demonstrate the correlation of CXCR4+ CD8 T cells in suppressive TIME of cervical cancer. Furthermore, CXCR4 inhibitors and agonists were used to explore the effects of targeting CXCR4+ CD8 T cells in promoting anti-PD-1 immunotherapy.

Results: Single-cell transcriptome analysis of multi-tissues and multi-cells revealed that CXCR4 was specifically expressed in T cells. Further analysis of single-cell profiling of the cervix and uterus demonstrated that CXCR4 was specifically expressed in CD8 T cells, with a positive correlation with the expression of PD-1 in cervical cancer. In TIME of cervical cancer, CXCR4+ CD8 T cells were associated with lymphocyte activation and tumor immune cell infiltration. By inhibiting the expression of CXCR4 via BL-8040 and AMD3100, the expression of PD-1 was reduced in concentration dependence, with the upregulation of PI3K-AKT-mTOR pathway. In addition, the expression patterns were reversed via the CXCR4 agonist, NUCC-390. Moreover, in vivo assays found that inhibiting CXCR4 promotes anti-PD-1 immunotherapy of cervical cancer via remodeling TIME.

Conclusions: Overall, single-cell profiling analysis reveals that CXCR4 was specifically expressed in CD8 T cells in cervical cancer, and combined CXCR4 and PD-1 blocked may expand the benefit of immunotherapy in cervical cancer.

Oral presentation

Dr Boero Veronica
veronica.boero@policlinico.mi.it - Italy

Boero Veronica
veronica.boero@policlinico.mi.it - Italy

#3376

The "CIV Classification," a New Proposal for the Architectural Grading of Vulvar Lichen Sclerosus

27 - Vulvar diseases and neoplasia

Background/Objectives: The purpose of this cross-sectional study was to prepare a reliable and easy-to-use architectural classification for vulvar lichen sclerosus (VLS) aimed at defining the morphological patterns of this condition.

Methods: An expert panel composed by 7 physicians with expertise in clinical care of vulvar conditions outlined the architectural criteria for the definition of VLS severity (phimosis of the clitoris, involvement of the interlabial sulci, narrowing of the vulvar introitus), identifying 5 grades to build up a classification. Thirteen physicians with 2-30 years expertise in vulvar diseases (nonexpert group) were asked to evaluate 3-5 pictures from 137 patients. Each physician individually assigned a grade to each case, according to the previously mentioned criteria. Interrater reliability was analyzed by means of intraclass correlation coefficient (ICC). The reliability concerning the 2 classifications of each rater was analyzed by means of κ statistic. Intraobserver and interobserver reliability in vivo was analyzed by means of κ index.

Results: This study provides a new classification of VLS, based on defined anatomical criteria and graded into mutually exclusive progressive classes. The ICC analysis showed a substantial interrater reliability of the classification, ICC = 0.89 (0.87-0.91), both in the expert panel and in the nonexpert group (ICC = 0.92 and 0.87, respectively). An "almost perfect" intraobserver and interobserver reliability was achieved among physicians in vivo (κ = 0.93).

Conclusions: Our classification showed a high reliability. It is easy to use, and it can be applied in clinical practice and eventually, in the evaluation of regenerative and cosmetic surgery.

Oral presentation

Reuschenbach Miriam
miriam.reuschenbach@msd.de - Germany

Luzak Agnes
agnes.luzak@msd.de - Germany

#3382

BURDEN OF CONIZATION AND ASSOCIATED HEALTHCARE COSTS IN WOMEN WITH CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN) - AN ANALYSIS OF GERMAN STATUTORY HEALTH INSURANCE CLAIMS DATA FROM 2013-2018

26 - Cervical neoplasia

Background/Objectives: Cervical intraepithelial neoplasia (CIN) is caused by human papillomavirus (HPV) infection. Management of CIN includes watchful surveillance or excision by conization. Data on clinical and economic burden of CIN and its treatment are limited for Germany. This study aimed to assess annual conization rates in women aged 18-45 years and related healthcare costs during a two-year follow-up period from the German statutory health insurance perspective.

Methods: We conducted a retrospective claims data analysis using the InGef Research Database (2013-2018). Cervical conizations in 18-45-year-old women were identified by OPS codes (5-671.0*/5-671.1*). For a subpopulation, subsequent conizations after a 90-day wash-out period and healthcare costs during a 24-month follow-up period (starting with an index conization) were assessed. Costs were compared (1:1:1 matching) in women with CIN diagnosis (ICD-10-GM codes N87.0, N87.1, N87.2 and/or D06) who underwent conization (study cohort 1) and women with CIN who did not undergo conization (study cohort 2) versus women with neither CIN nor conization (control group) as all-cause costs for outpatient care, inpatient care, outpatient pharmaceuticals, aids and remedies, and sick leave costs. Mean imputation was applied for missing aids and remedies (~20%) and sick leave data (~5-23%).

Results: Between 2013-2018, the annual number of 18-45 year old women who underwent conization ranged from 611,380 to 630,284, giving an annual conization rate of 2.4/1,000 in 2013 and 2.1/1,000 in 2018, with over 60% of the observed women with conizations between 31-45 years old. The subpopulation used for the analysis of subsequent conizations and healthcare costs included N=2,749 women. Of these, 2.91% underwent a subsequent conization three months or later (up to 24 months) after the initial conization. Mean total healthcare costs during the 24-months follow-up period were higher in both study cohorts (study cohort 1: €4,446, $p<0.01$; study cohort 2: €3,754, $p=0.09$) compared to the control group (€3,426) and were mainly driven by inpatient care (study cohort 1: €1,849, $p<0.01$; study cohort 2: €1,348, controls: €1,169, $p=0.07$) followed by outpatient care (study cohort 1: €1,418, $p<0.01$; study cohort 2: €1,339, $p<0.01$; controls: €1,170). The highest mean total cost difference was found in women between 36-45 years.

Conclusions: Our results highlight a substantial burden of cervical conizations, especially in women aged 30 and older. This burden leads to €1,020 and €328 higher healthcare costs in women with a CIN diagnosis with and without conization, respectively, compared to women without CIN in a two-year follow-up period.

Oral presentation

Ms Vahteristo Maija
maiya.vahteristo@cancer.fi - Finland

Vahteristo Maija
maiya.vahteristo@cancer.fi - Finland

#3383

Searching for optimal algorithms for primary HPV screening

12 - Triage of HPV positive women

Vahteristo M

Background/Objectives: Primary HPV screening programmes have been implemented in many European countries. Currently many of these screening programmes suffer from overdiagnosis and high colposcopy burden. Our goal was to identify the key characteristics of various algorithms for the management protocols for primary HPV screening with cytology triage, based on the Finnish HPV implementation trial with over 20-year follow-up.

Methods: Using the HPV implementation trial data, we formulated post-hoc 6 different algorithms for HPV screening (HPV 1 - 6) and one for cytology screening. We combined these data with all the CIN II+ cases registered during this trial. Sensitivity, specificity, and colposcopy referral rate for CIN II+ detected during the first and second screening round were calculated for the algorithms.

Results: Results are presented in a picture provided as a supplementary file.

Conclusions: Almost all the HPV algorithms were superior in the CIN II+ sensitivity compared to the cytology algorithm which has a high effectiveness in Finland. The results were consistent in the first and second screening rounds of the trial. In HPV algorithms, the return of HPV positives back to the routine screening at any stage of the screening round will reduce the sensitivity of the screening. To achieve algorithms for primary HPV screening with both low colposcopy burden and good sensitivity, other triage methods than cytology are needed.

Poster

Dr Silva Rita
rita.sccsilva@gmail.com - Portugal

Pacheco Maria Amália
malita57@gmail.com -

#3384

Cervical adenocarcinoma: review of the last 5 years in a tertiary center

26 - Cervical neoplasia

Background/Objectives: Cervical cancer is a leading cause of mortality among women. In 2020, an estimated 604 000 women were diagnosed with cervical cancer worldwide and about 342 000 women died from the disease. Despite the implementation of screening and vaccination programs against HPV, cervical cancer is still considered a public health problem today. The epidermoid cancers are more prevalent, but the incidence of adenocarcinoma has increased over the past few years, especially in young women. This study aimed to describe the cases of cervical adenocarcinoma in the last 5 years in our center.

Methods: Were review 12 women diagnosed with cervical adenocarcinoma from january 2017 to october 2021.

Results: Were analyzed twelve cases of adenocarcinoma: 4 cases of adenocarcinoma in situ (33,3%) and 8 invasive (66,7%). The average age at diagnosis was 47,3 years (min-max 31-75) with an average age at coital age 17 years. The average number of partners was 2.75 however 4 women did not respond. Regarding the age group, 66.7% were of reproductive age, with 75% of them using hormonal contraception. Regarding personal history, the patients were healthy, except one with a history of drug abuse and HCV infection. Smoking habits were described by 1/3 of the patients and only 8% had complied with the HPV vaccination schedule. Seventy-five percent of patients were referred to the center with altered cytology (ASCUS=4; AGC=3; HSIL=2) and twenty five percent were referred due to abnormal uterine hemorrhage/coitorrhagia. Two of the patients with a final diagnosis of adenocarcinoma invasive reported that it was the first cytology ever performed. HPV typing was performed in only seven patients: in six cases (86%) HPV 16 was present and one (14%) was negative for HPV. On colposcopic examination, grade 1 findings were found in 16% (n=2), grade 2 in 66% (n=8), and frank suspicion of invasion in 16% (n=2). The final histological diagnosis was obtained by directed biopsy in 10 patients and 2 by large loop excision of the transformation zone (LLETZ). Sixty-three percent of invasive adenocarcinomas were diagnosed at stage I of FIGO (IA 2:1, IB1:2;IB3:2); 12.5% in stage II (IIB:1), 12.5% in stage III (IIIC2: 1) and 12.5% in stage IV (IVB: 1). Regarding treatment, two women opted, informedly, for conservative treatment (LLETZ), five (42%) after EZT were submitted to hysterectomy (33% radical hysterectomy with bilateral anexectomy and bilateral pelvic lymphadenectomy and 16% total hysterectomy); three patients underwent chemotherapy plus radiotherapy (CRT) and one patient is currently pregnant (diagnosis at 29 weeks) with indication for CRT after cesarean section.

Conclusions: Although the sample was small, we can gain that the incidence of adenocarcinomas in our unit was slightly higher than the reduction in the literature (31% vs 25%) with a strong association with HPV 18. Thus constituting a real challenge for the clinic, being urgente the need to implemente more studies aimed at facilitating the diagnosis and approach of this entity.

Oral presentation

Mr Renner Janis
janis.renner@med.uni-tuebingen.de - Germany

Mr Renner Janis
janis.renner@med.uni-tuebingen.de - Germany

#3394

BRD4S INTERACTS WITH VIRAL E2 PROTEIN TO LIMIT HUMAN PAPILLOMAVIRUS LATE TRANSCRIPTION

02 - Viral and molecular biology

Background/Objectives: Papillomaviruses have coevolved with their host by using cellular factors like Bromodomain containing protein 4 (Brd4) to support genome maintenance, gene transcription and replication. The C-terminal domain (CTD) of Brd4 represents a highly conserved binding site for the N-terminal domain of E2, which is important for its transcription activator function. Recently the PDID (phosphorylation dependent interaction domain) of Brd4 was described as an additional interaction site for E2. In addition to full-length Brd4 a short isoform of Brd4 (Brd4S) is expressed which contains the PDID but lacks the CTD and thus might interact with E2. Therefore, we analyzed the interaction of Brd4S and E2 and explored its functional consequences.

Methods: We used flow cytometry based-FRET analysis and co-immunoprecipitations to analyze protein-protein interactions, chromatin-immunoprecipitation (ChIP) to study interaction with the viral origin, immunofluorescence analysis for co-localization studies. SiRNA knockdown were used to determine viral transcription and replication activities in undifferentiated and differentiated cells

Results: We confirmed that the isolated CTD and PDID of Brd4 interact with E2 by FRET assays. Consistent with this, Brd4S co-immunoprecipitated with HPV31 E2. Immunoblot analyses indicated that Brd4S is present in significant amounts in HPV-negative and -positive keratinocytes. Immunofluorescence analysis revealed a partial co-localization of Brd4S with E2 in the nucleus which was greatly enhanced by the presence of the E1 protein and the viral origin. Consistent with this, ChIP-experiments revealed that BRD4S binds to the viral origin. Knockdown and overexpression studies indicated that Brd4S inhibits E2's transactivation function. In line with this, knockdown of BRD4S induced viral late transcription in undifferentiated cells maintaining replicating HPV31 genomes.

Conclusions: Our study reveals an inhibitory role of BRD4S on HPV transcription, which may serve as an immune escape mechanism by the suppression of L1 transcripts and thus contribute to the establishment of persistent HPV infections.

Poster

Dr Strbac Mirjana
mirjana.strbac@izjzv.org.rs - Serbia

Dr Strbac Mirjana
mirjana.strbac@izjzv.org.rs - Serbia

#3400

MOTIVATION FOR THE HPV VACCINE UPTAKE AMONG PARENTS OF VACCINATED CHILDREN IN NOVI SAD (SERBIA): RESULTS FROM THE INITIAL FINDINGS

40 - Public health

Background/Objectives: Current estimates indicate that Serbia ranks third in the morbidity rate and fourth in the death rate from cervical cancer in Europe. Every year in Serbia around 634 women die from cervical cancer. The HPV vaccine, has been available in Serbia since 2008, recommended, but not "free of charge". To determine motivation for the HPV vaccination of children, a cross-sectional survey was conducted among parents during November, 2021. Within the project in the City of Novi Sad, the HPV vaccine was offered "free of charge" for children aged 12-18 years, regardless of gender.

Methods: The Institute of Public Health of Vojvodina collected data at the vaccination point in collaboration with the Health Care Centre of Novi Sad, using an anonymous survey questionnaire. The questionnaire consisted of parents' sociodemographic data, age and sex of children and 17 motives (positive parental attitude towards HPV vaccination) as possible reasons for deciding to vaccinate children. Chi-squared test or Fisher's exact test was for categorical and Wilcoxon rank-sum test for nonparametric data and significance levels are considered $p < 0.05$.

Results: 136 parents of vaccinated children aged 12-18 participated in the survey. The average age of the parents was 45 years, and 80% were female. The percentage of vaccinated girls in respect to boys was 88%. Out of the total number of vaccinated children, 25% were 15 years old. The strongest motive (29.4%) for vaccination in the entire group was the attitude that "It is better to immunize child than expose them to risk possible infection". Around 26% of parents that were healthcare workers selected answer that strongest motive for vaccination was recommendation from a paediatrician, while parents from other professions selected the affirmation that it is better to immunize child than to expose them to risk of possible HPV infection (24.7%). Also, paediatrician's recommendation was more frequently selected motive among health professionals compared with other professions (78.3% vs 50.7%, $p = 0.02$). Recommendation given from friends or family members was motive for 64.4% of parents from other professions compared with 34.8% of health professionals ($p = 0.01$). Motive related to the fact that vaccine is "free of charge" during the period of this project was selected in similar percentage by both groups of parents ($p = 0.942$).

Conclusions: This project is ongoing. In order to better understand the main reasons for acceptance of HPV vaccine additional research is needed. Until then, it is crucial to raise awareness of infection consequences and establish public dialogs in media as well as among pediatric professionals.

References: Bruni L, Albero G, Serrano B, Mena M, Collado JJ, Gómez D, Muñoz J, Bosch FX, de Sanjosé S. ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and Related Diseases in Serbia. Summary Report 22 October 2021. [Accessed 20 November 2021]

Oral presentation

Mrs De Lepper Marion
marion.de.lepper@msd.de - Germany

De Lepper Marion
marion.de.lepper@msd.de - Germany

#3408

EFFECT OF HUMAN PAPILLOMAVIRUS (HPV) VACCINATION ON OCCURRENCE OF SUBSEQUENT ANOGENITAL WARTS (AGWs) IN PATIENTS WITH CURRENT OR HISTORY OF AGWs WORLDWIDE: A SYSTEMATIC REVIEW AND META-ANALYSIS

06 - HPV prophylactic vaccines

Background/Objectives: Anogenital warts (AGWs) are responsible for a high burden of disease and are caused mainly by the low-risk human papillomavirus (HPV) types 6 and 11. After standard therapy up to approximately 48.5% of patients suffer from re-detection of AGWs. The aim of this systematic review was to evaluate whether prophylactic HPV vaccination reduced the risk of subsequent AGW occurrence in patients with current or history of AGWs.

Methods: A literature search was carried out on June 30, 2021 in PubMed, MEDLINE, EMBASE, Global Health, Scopus, and the Cochrane library. Randomised trials and analytical observational studies on patients with clinically diagnosed history or current AGWs worldwide were included. Studies on patients only diagnosed by laboratory tests were excluded. Studies were included when they used HPV vaccines covering at least HPV type 6 and/or 11. The primary outcome for the randomised trials was the proportion of AGW re-detection after HPV vaccination in patients with previously clinically diagnosed AGWs. The calculated risk ratio (RR) of those studies were summarised in a meta-analysis. A sensitivity analysis was conducted, excluding the study that mostly differed regarding study design and study population. The observational studies were evaluated narratively. Finally, quality assessment of the included studies and assessment for publication bias were conducted.

Results: Totally, six studies were included in the systematic review, four randomised trials and two observational studies. The effect of HPV vaccination on subsequent AGWs among patients with current AGWs which received AGW therapy was investigated in randomised trials, while the association between HPV vaccination and subsequent AGWs among patients with AGW history either cleared naturally or after treatment was investigated in observational studies. The summary RR of the randomised trials showed evidence of a risk reduction in subsequent occurrence of AGWs in patients with HPV vaccination compared to controls among patients with previous AGWs (RR = 0.56, 95% confidence interval (CI): 0.33 - 0.95). A sensitivity analysis resulted in RR = 0.46 (95% CI: 0.33 - 0.64), indicating a robust result. We did not carry out a meta-analysis on included observational studies due to the limited number of similar studies. Most observational studies reported a lower risk of subsequent AGWs among HPV vaccinated people compared to controls among patients with AGWs.

Conclusions: The systematic review showed some evidence that HPV 6 and/or 11 targeted vaccination reduced the risk of subsequent AGW occurrence among patients with previous AGWs. More studies may validate this result.

Poster

Dr Natasa Nikolic
natasa.nikolic@izjzv.org.rs - Serbia

Nikolic Natasa
natasa.nikolic@izjzv.org.rs -

#3413

PREVALENCE AND GENOTYPE DISTRIBUTION OF HIGH-RISK HPV INFECTION IN TEN YEAR PERIOD IN VOJVODINA, SERBIA

14 - Genotyping

Background/Objectives: Human papillomavirus (HPV) infection is the leading risk factor for cervical cancer development. Among women in Serbia, cervical cancer is the 5th most frequent cancer and the 2nd most frequent cancer in women 15-44 years of age. The incidence rate of cervical cancer in Serbia is approximately twice as high as the average incidence rate in Europe (10.7/100,000). This study aimed to determine the prevalence and distribution of different high-risk (HR) HPV genotypes according to age among women in Vojvodina, northern province of Serbia.

Methods: From January 2012 to November 2021, a total of 7974 women from 14 to 80 years of age, with an abnormal Pap test and normal cytology results with suspicious colposcopic findings, were included in the study. Extraction of HPV DNA was performed using SaMag STD DNA Extraction Kit on SaMag-12 Automatic Nucleic Acids Extraction System. Twelve HR HPV types (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, and 59) were determined by the HPV High Risk Typing Real-TM kit on the SaCycler-96 Real Time PCR System.

Results: The overall prevalence of HR HPV was 43.3% (3455/7974). We observed that 62% of women had a single infection, while 38% had multiple HR HPV infections. The results showed that members of the alpha-9 HPV species, particularly HPV 16 (25.2%), HPV 31 (18.9%), and HPV 39 (6.9%), were the most prevalent HPV genotypes in our province. Additionally, 6.9% of the single infections were caused by HPV 51, a member of the alpha-5 HPV species. Other HR HPV genotypes were found in 42.1% of the single infections. HR HPV 16 was the most prevalent genotype overall, 30.2% (1042/3455). The age-specific distribution of single HR HPV infection was as follows: 14-20 (57.4%), 21-30 (55.0%), 31-40 (45.8%), 41-50 (37.6%), 51-60 (34.3%), and the older than 61 years (44.1%).

Conclusions: The high prevalence and presence of vaccine HPV types, particularly type 16, indicates the need for a large-scale immunization program in our province in order to decrease the prevalence of HPV infections and future cervical cancer incidence. Determination of the contribution of HR HPV genotypes causing cervical cancerogenesis is valuable as a starting point for the implementation of a vaccination program and as a critical step in the algorithm of cancer diagnosis and treatment of individual patients.

Oral presentation

Dr Felsher Marisa
marisa.felsher@merck.com - United States

Morais Edith
edith.morais@msd.com -

#3453

THE IMPACT OF COVID-19 ON THE IMPLEMENTATION OF PROGRESS (PRevalence of Oral hpv infection, a Global aSSessment) STUDY

29 - Oral HPV infection

Background/Objectives: Little is known about the natural history of oral HPV in the general population. We designed PROGRESS (PRevalence of Oral hpv infection, a Global aSSessment), a noninterventional study among adults recruited from dental offices in France, Germany, Spain, the United Kingdom and the United States (US) to assess oral HPV prevalence, incidence and persistence. The COVID-19 pandemic has impacted research conducted by dentists as the nature of their work puts them and patients at high risk of COVID-19 infection. We piloted PROGRESS in four dental sites to assess feasibility of study implementation during the global pandemic.

Methods: Between November 2020 and June 2021, four sites were selected for the pilot phase: one in France, one in Spain and two in the US. Participating dentists were instructed to recruit 68 patients presenting for dental examinations over a minimum of 28 and maximum of 40 working days, for 272 participants total. The study was approved by Ethics Committees in accordance with local and national requirements. Participants gave informed consent, completed questionnaires and provided oral rinse/gargle (ORG) specimens. ORG specimens were shipped to reference laboratories for HPV-DNA detection using the SPF10/DEIA/LiPA25 system. Implementation was assessed as follows: total participants enrolled per site; total days between first and last participant in study per site; total ORG samples transported to laboratory; proportion of samples adequate for testing; and total completed participant questionnaires. Interviews were held with dentists to identify hurdles to study implementation.

Results: Three dentists recruited 68/68 participants; the French site recruited 45/68 participants for a total of 249 participants. The mean number of days between first and last participant enrolled was 60 days: 65 days in France, 66 days in Spain, 28 days in US site 1 and 82 days in US site 2. There were 249/249 samples collected and sent to reference laboratories, 249/249 samples processed and classified as adequate after DNA quality control and 249/249 completed questionnaires. Dentists reported that recruitment delays were due to insufficient time and staff to execute fieldwork.

Conclusions: COVID-19 did not impact ORG collection, transportation, or processing nor completion of questionnaires, but did impact the rate at which participants were enrolled in three of four sites, due to insufficient time and staffing. PROGRESS will expand to 108 more dental sites and enroll 7,628 participants between 2021 and 2022. Strategies to support rapid participant recruitment such as weekly reminder calls will be used to ensure timely enrollment.

Oral presentation

Dr Nitkowski Jenna
jnitkowski@mcw.edu - United States

Nitkowski Jenna
jnitkowski@mcw.edu - United States

#3470

EFFECTS OF ENVIRONMENTAL CONDITIONS ON HOME-BASED SELF-SAMPLING KITS FOR ANAL CANCER SCREENING

13 - Self-sampling

Nitkowski J, Giuliano A, Ridolfi T, Chiao E, Fernandez M, Schick V, Swartz M, Smith J, Brzezinski B, Nyitray A

Background/Objectives: The performance of home-based self-sampling kits may be subject to environmental conditions such as extreme temperatures, freeze thaw cycles, or long transport times in transit to laboratories. We aim to determine the environment's effect on specimen adequacy for HPV genotyping of a home-based self-sampling anal cancer screening kit.

Methods: The Prevent Anal Cancer Study in Milwaukee, Wisconsin, USA recruited men who have sex with men and transgender persons 25 years of age or older. One-half of participants were randomized to receive a mailed self-collection kit which included a flocked swab (Copan Italia Spa), a vial of 2 mL of standard transport media (Qiagen), and a temperature recorder. The kits were insulated with foam and temperature was recorded every twenty minutes. Participants mailed the kit back to the lab. Subsequently, the SPF10 LiPA assay was used for HPV genotyping including the detection of human β -globin to determine sample inadequacy by qPCR. Using logistic regression, we assessed the association between specimen inadequacy and mean temperature, lowest temperature, highest temperature, temperature range, presence of a freeze-thaw cycle, and the number of days in an uncontrolled environment. This analysis assessed inadequacy for the first 52 kits.

Results: A total of 9.6% of specimens were inadequate for HPV genotyping. Kits experienced an average of 12.7 days in an uncontrolled environment at a mean temperature of 20.2°C. Kits were subjected to low temperatures ranging from -16.0°C to 21.4°C, and high temperatures ranging from 22.1°C to 39.4°C. On average, kits were subjected to a temperature range of 18.1°C (min=6.1, max=40.2). After adjusting for number of days in an uncontrolled environment, none of the time and temperature variables were associated with specimen inadequacy including low temperature (aOR 0.92, 95% CI 0.82-1.02), high temperature (aOR 0.95, 95% CI 0.72-1.24), and temperature range (aOR 1.12, 95% CI 0.98-1.27). A higher proportion of inadequate specimens experienced a freeze-thaw cycle (40%) as compared to adequate specimens (13%), but this difference did not reach statistical significance. Sensitivity analyses using participant-reported date of swabbing revealed that low temperature and temperature range were significantly associated with specimen inadequacy, although more of these data were missing and subject to recall bias.

Conclusions: Over 90% of home-based self-swabs were adequate. Despite transit during different seasons in Milwaukee, specimen inadequacy was not associated with temperature (low, high, mean, or range), freeze-thaw cycle, or days in an uncontrolled environment.

Oral presentation

Dr Muresu Narcisa
illarisechi@yahoo.it - Italy

Sechi Illari
illarisechi@yahoo.it - Italy

#3491

Comparison of traditional and rapid nucleic acid extraction using a non-alcohol-based medium for vaginal self-samples

09 - HPV testing

Sechi I, Del Rio A, Muresu N, Dettori A, Usai M, Cossu A, Cocuzza C, Muroi M, Castriano S, Piana A

Background/Objectives: Vaginal self-collected sample is a convenient and economical method, as well as a useful strategy to reach women who do not participate to cervical cancer screening. Although the higher sensitivity of the HPV-DNA testing, the accuracy of the test is related to the stability of the swabs during transport and to the elution medium, particularly for vaginal self-collection. Moreover, new approaches in HPV screening, as rapid thermal nucleic acids extraction (RTNAE), could be useful in reducing cost and time of analyses. MSwabTM is a non-alcohol-based medium for the collection, and storage of clinical specimens, suitable for traditional NA extraction or RTNAE. The aim of the present study was to evaluate the analytical performance of both traditional and RTNAE methods using dry self-collected FLOQSwabs[®] eluted in MSwabTM medium (COPAN Italia) for high-risk HPV detection.

Methods: A total of 194 women were enrolled, within regional LILT (Italian League against Tumor) project. Dry self-collected FLOQSwabs[®] samples, were suspended in 2 mL of MSwabTM and processed at the Molecular Epidemiological laboratories of University of Sassari (Sardinia, Italy). Traditional DNA extraction was carried out firstly by an automated system (Seegene AdvanSureTM E3) and analyzed by AnyplexTM II HPV HR (Seegene). All high-risk HPV-positive and some HPV-negative MSwab-samples, stored at room temperature and at -20°C, were processed with a RTNAE at 98°C/5mins and analyzed for high-risk HPV. Diagnostic agreement was assessed with the Cohen's K statistic.

Results: A total of 15.5% (30/194) of samples resulted positive after traditional extraction method. In the 30 HPV-positive and 10 HPV-negative, RTNAE tested after storage at RT (mean days of storage: 177) and storage at -20°C, all negative samples and 26/30 (86.7%) of HPV-positive were confirmed. Overall, an excellent agreement was detected among the two extraction methods and storage conditions: - After RT storage: 90% agreement ($k=0.77$, $p<0.0001$) between Traditional DNA extraction vs RTNAE methods. - After -20°C storage: 90% agreement ($k=0.77$, $p<0.0001$) between Traditional DNA extraction vs RTNAE methods. - RTNAE methods after RT vs -20 °C storage: 95% agreement ($K= 0.89$, $p<0.0001$).

Conclusions: The present study showed as MSwabTM medium was a reliable and stable medium for elution and storage of dry vaginal self-collected FLOQSwabs[®]. These preliminary results showed a comparable high diagnostic accuracy of HPV-DNA between automated and RTNA extraction. MSwabTM direct-rapid thermal extraction, improves results turnaround time and saves costly extraction reagents. A non-alcohol medium for rapid and traditional nucleic acids extraction for detection of HPV in vaginal self-collected samples.

Oral presentation

Ms Sand Freja Laerke
freja@cancer.dk - Denmark

Sand Freja L.
freja@cancer.dk - Denmark

#3492

RISK OF RECURRENT DISEASE AFTER CONIZATION ACCORDING TO TIMING OF HPV VACCINATION IN RELATION TO TREATMENT

06 - HPV prophylactic vaccines

Sand F, Galanakis M, Frederiksen K, Kjaer S

Background/Objectives: The aim of this study was to examine the risk of cervical intraepithelial neoplasia grade 2 or worse (CIN2+) after conization according to timing of HPV vaccination in relation to treatment.

Methods: We conducted a nationwide cohort study using multiple Danish registries. Initially, we identified women diagnosed with CIN3 on the cone (2009-2016) and their HPV vaccination status. Vaccinees were defined as women vaccinated between 3 months before until 1 year after conization. Vaccinees were divided into groups according to timing of vaccination; 0-3 months before conization, 0-2 months after conization and 2-12 months after conization. The women were followed from one year after conization until diagnosis of CIN2+, conization, total hysterectomy, death, emigration or end of follow-up (December 2018). Cox proportional hazard regression was used to estimate hazard ratios (HRs) and corresponding 95% confidence intervals (CIs) of CIN2+ comparing vaccinees with non-vaccinees. The HRs were adjusted for age, histology on cone, education, year of conization, repeat conizations and CIN2+ lesions between conization and start of follow-up.

Results: We included 19301 women diagnosed with CIN3 on the cone of which 4052 women were vaccinated and 15249 women were not vaccinated in relation to conization. During follow-up, 183 and 732 women were diagnosed with CIN2+ among vaccinated and unvaccinated women, respectively. Among the vaccinees, 1131 women were vaccinated 0-3 months before conization, 1350 women were vaccinated 0-2 months after conization and 1571 women were vaccinated 2-12 months after conization. Unvaccinated women had a 4- and 8-year absolute risk of CIN2+ of 3.7 (95%CI: 3.4-4.0) and 5.5 (95%CI: 5.1-6.0), respectively. The corresponding absolute risks of CIN2+ was 2.8 (95%CI: 1.9-3.7) and 4.6 (95%CI: 3.2-6.1) for women vaccinated 0-2 months after conization. Women vaccinated 0-3 months before or 2-12 months after conization had approximately the same absolute risk of CIN2+ as unvaccinated women. Women vaccinated just after conization (after 0-2 months) had a decreased risk of CIN2+ compared to unvaccinated women (HR_{adjusted}=0.7; 95%CI: 0.5-1.0). The adjusted HR of CIN2+ was 0.9 (95%CI: 0.7-1.2) and 0.9 (95%CI: 0.7-1.2) for women vaccinated 0-3 months before and 2-12 months after conization, respectively.

Conclusions: Our results indicate that vaccinating women just after conization might decrease the subsequent risk of CIN2+.

Oral presentation

Mr Dolk Christiaan
christiaan.dolk@merck.com - Netherlands

Dolk Christiaan
christiaan.dolk@merck.com - Netherlands

#3493

THE PUBLIC HEALTH IMPACT AND COST EFFECTIVENESS OF GENDER NEUTRAL 9VHPV VACCINATION IN THE NETHERLANDS

36 - Economics and modelling

Dolk C, Sabale U, Palmer C

Background/Objectives: HPV vaccination targeting 12-year-old-girls is included in the National immunization program (NIP) since 2010, and uptake currently plateaus at approximately 60%. Since introduction, a bivalent human papillomavirus (2vHPV) vaccine against two HPV types 16/18 responsible for 70% of cervical cancer was used. In 2019, the Dutch Health Council recommended lowering the age of the primary vaccination cohort and the inclusion of boys in the NIP. Following this recommendation, starting in 2022, both 9-year-old boys and girls will be eligible for HPV vaccination.¹ This study assesses whether a gender-neutral vaccination with a nonavalent HPV vaccine (9vHPV) against HPV types 6/11/16/18/31/33/45/52/58 is a cost-effective strategy in the Netherlands.

Methods: Using a previously published dynamic transmission model for HPV infections that has been adapted to the Netherlands, we performed a base line analysis assuming the program continues the current strategy of gender-neutral 2vHPV vaccination (with a switch to the 9-year-olds and compared the outcomes with those resulting from a strategy with 9vHPV vaccination in the same cohort.^{2,3} Various sensitivity analyses were performed, including assumptions of vaccination coverage rates (VCRs) and the degree of cross-protection. The analysis is performed from a healthcare perspective over a 100-year time horizon.

Results: Compared to 2vHPV vaccination, vaccination with a 9vHPV vaccine was estimated to significantly decrease the cumulative incidence of HPV 6/11/16/18/31/33/45/52/58-related diseases, averting over 3000 additional HPV related cancer cases, approximately two-thirds of which are averted cervical cancer cases, over the studied horizon, with over 1000 additional lives saved. Additionally, 9vHPV vaccination was estimated to avert over 1 million cases of genital warts and decrease the incidence of genital warts by over 80%. Incidence of both adult- and juvenile-onset of recurrent respiratory papillomatosis were reduced by approximately 95%, with a total of over 4000 averted cases. Compared to a 2vHPV vaccination strategy, 9vHPV vaccination was demonstrated highly cost-effective (ICER under €8000 per quality-adjusted life-years), which is robust according to sensitivity analysis results around VCRs.

Conclusions: 9vHPV vaccination can considerably decrease more HPV-related cancers and genital warts than the current 2vHPV vaccination program and is a highly cost-effective strategy.

References: 1. van Lier, E. A. et al. Immunisation coverage and annual report National Immunisation Programme in the Netherlands 2020. [In Dutch] (2021). 2. Elbasha, E. H., Dasbach, E. J. & Insinga, R. P. Model for assessing human papillomavirus vaccination strategies. *Emerg. Infect. Dis.* 13, 28-41 (2007). 3. Elbasha, E. H. & Dasbach, E. J. Impact of vaccinating boys and men against HPV in the United States. *Vaccine* 28, 6858-6867 (2010).

Oral presentation

Dr Del Rio Arcadia
a.delrio25@studenti.uniss.it - Italy

Dr Del Rio Arcadia
a.delrio25@studenti.uniss.it - Italy

#3498

Vaginal self-sampling as a tool for the prevention of cervical cancer

11 - Screening for women difficult to reach

Dettori A, Sechi I, Muresu N, Usai M, Cossu A, Muroi M, Castriano S, Cocuzza C, Piana A

Background/Objectives: Human papillomavirus (HPV) infection is the major cause of cervical cancer. An effective approach for cervical cancer screening is the use of vaginal self-sampling in HPV-DNA testing, based on its accuracy and sensitivity. Moreover, use of vaginal self-collection device could improve the participation to cervical screening programs. The present study, organized by LILT (Italian League Against Tumors) and University of Sassari (Sardinia, Italy), aimed to increase the adhesion of hard-to-reach women to screening program and identify the potential population target, examining demographic and clinical characteristics of participants.

Methods: From April to July 2021, the participants received a kit containing instruction for use, a vaginal self-sampling device (FLOQSwabs®, Copan Italia) and a questionnaire assessing the ease of use of the collection device and the acceptability of vaginal self-collection. Overall, 371 women returned the questionnaire and a FLOQSwabs®. All dry swabs were processed for DNA extraction and analysed by PCR, using Anyplex™ II HPV HR Detection Kit (Seegene) that detects 14 high-risk HPV genotypes, at the Molecular Epidemiology laboratory of University of Sassari (Sardinia, Italy).

Results: The participating women were 371 (mean age of 42.7 years; SD ± 11.4); 16.1% (60/371) of samples were HPV positive and the most common genotypes were 56, 31, 18 and 51. No invalid results were detected. Co-infection was reported in 33.3% (20/60) of samples. Among the 60 HPV-positive women, 20% (12/60) declared they never received a Pap test. 30.7% and 7.3% of single and married women were HPV-positive, respectively. The highest HPV positivity (27%) was detected in women under 35 years. FLOQSwabs® was found easy to use, self-collection was preferred to clinician collection by ~58% of participants, with variations related to different geographical areas.

Conclusions: The present study confirmed that vaginal self-sampling represents a valuable and well-accepted method in cervical cancer screening. HPV positivity rate and genotypes detected confirmed data previously reported in our setting. The questionnaire highlighted the acceptability of vaginal self-sampling with FLOQSwabs®, and the differences recorded could be associated with the quality and closeness to screening centres in Sardinian region. Self-sampling may increase screening participation, particularly among "non-adherents" women, overcoming practical, economic, and cultural barriers.

Poster

Dr Mitran Mihai
michael_digital@yahoo.com - Romania

Dr Mitran Mihai
michael_digital@yahoo.com - Romania

#3499

HPV INDUCED ALTERATIONS IN VULVAR PATHOLOGY

27 - Vulvar diseases and neoplasia

Olinca M, Poteca A, Bratila E

Background/Objectives: Human Papilloma Viruses have been the subject of numerous studies related to cervical cancer. But the tropism of these viruses for the squamous epithelium does not stop at the cervical level. Histologically similar lesions can be identified in other epithelia. Precursor lesions of squamous cell carcinoma can be divided into two major groups, depending on the association with HPV infection. The new WHO classification (2020) emphasizes the importance of human papillomavirus (HPV) infection in the development of vulvar squamous cell carcinoma, thus reflecting on patient management. Our review studies the correlation between clinical data, HPV status, histological features and molecular markers in vulvar squamous lesions.

Methods: We selected cases diagnosed with vulvar lesions and recorded clinicopathologic variables, as well as immunohistochemical test results. All slides were reviewed and classified according to the new WHO classification.

Results: Of a total of 21 cases, the HPV-associated ones showed distinctive morphology that suggested their nature even before the p16 staining

Conclusions: A clear distinction of the place of origin of the tumor is often difficult to achieve both clinically and by imaging techniques, especially in advanced diffuse lesions, exceeding the limits of anatomical structures. Vulvar squamous lesions represent a heterogenous group of tumours, some with adverse outcome and although their recognition can be difficult, refinement of diagnosis and prognostication based on a systematic evidence review has great value in a better personalized treatment for these patients.

Oral presentation

Dr Ring Linea Landgrebe
lilr@cancer.dk - Denmark

Ring Linea L.
lilr@cancer.dk - Denmark

#3500

PARTICIPATION IN CERVICAL CANCER SCREENING AMONG OVERWEIGHT AND OBESE WOMEN: A SYSTEMATIC REVIEW AND META-ANALYSIS

11 - Screening for women difficult to reach

Background/Objectives: The incidence of cervical cancer has decreased after implementation of cervical cancer screening. However, obese women have higher risk of cervical cancer than women of normal weight. Previous studies have found lower rates of participation in cervical cancer screening among obese compared to non-obese women, but an overview of the literature is lacking. We conducted a systematic review and meta-analysis of studies examining participation in cervical cancer screening among overweight and obese women compared to women with normal body mass index (BMI).

Methods: We conducted this systematic review and meta-analysis in accordance with PRISMA guidelines. The databases PubMed, Embase and the Cochrane Library were searched from 1 January 2000 to 12 July 2021 to identify English language studies reporting participation in cervical cancer screening among overweight or obese women compared to women with normal BMI. We included studies of women from the general female population eligible for cervical cancer screening. Data were extracted on the following variables: author, publication year, country, study design, study population, exclusion criteria, number of included women, age, race, screening interval, method of measuring screening participation and BMI, BMI categories, response rate to survey, statistical method, total number of women and number of women participating in screening in each BMI category, and relative risk estimates. Two reviewers independently extracted data, and inconsistencies were discussed to reach consensus. Furthermore, the quality of studies was scored using an adapted version of the New Castle-Ottawa scale. We plan to use a random effects model to estimate pooled odds ratios of screening participation comparing different BMI categories, e.g. overweight (BMI=25–30), obese class I (BMI=30–35) and obese class II (BMI=35–40), with women with normal BMI (BMI=18.5–25). Where possible, we will pool adjusted and unadjusted relative risk estimates separately.

Results: Our search identified 1676 records, of which we removed 247 duplicates. Titles (N=1429), abstracts (N=287), and full text articles (N=100) were reviewed by two authors independently. Reference lists were reviewed to identify additional relevant studies (N=9). Altogether, we included 57 studies. Data extraction for the study is ongoing. Results on relative risk of cervical cancer screening participation according to BMI will be presented at the conference.

Conclusions: To our knowledge, this is the first systematic review and meta-analysis of cervical cancer screening participation in overweight and obese women including data from a range of different countries.

Poster

Ms Godoy Luani
luanirgodoy@gmail.com - Brazil

Ms Godoy Luani
luanirgodoy@gmail.com - Brazil

#3501

HIGH-RISK HPV TYPES USING NEXT GENERATION SEQUENCE (NGS) IN EUROPEAN AND LATIN WOMEN WITH NEGATIVE AND POSITIVE CYTOLOGY: PRELIMINARY RESULTS FROM ELEVATE STUDY

14 - Genotyping

Guimarães Y, Dias T, Campanella N, Vermandere H, Degomme O, Dos Reis R, Longatto-filho A.

Background/Objectives: The main etiological factor of cervical cancer (CC) and its precursor lesions is a persistent human papillomavirus (HPV) infection. High-risk HPV (hr-HPV)-induced carcinogenesis is mainly related to E6/E7 integration in human DNA that causes transformation of the host cells. This virus can be classified into several levels: genus, species, type, subtype, and variants, which can be divided into lineages and sub-strains according to the percentage of genetic similarity. This project is part of an international collaboration entitled ELEVATE of which the main objective is to develop a new point-of-care screening device based on self-sampling. The new test will be able to detect 14 hr-HPV types and protein biomarker expression.

Methods: In order to develop the HPV DNA test and detect well-conserved target regions, hr-HPV genotyping was performed using Ampliseq manually and Ion Torrent S5 technologies in 1000 samples collected in Brazil, Ecuador, Belgium, and Portugal (300 normal samples, 300 low-grade lesions, and 400 high-grade lesions). Through NGS of the E6/E7 genes, it will be possible to genotype all 14 hr-HPV individually and also HPV co-infections. The genotyping was performed with a workflow developed in the Galaxy platform, in which the alignment was made against the 14 hr-HPV genotypes.

Results: To date, 106, 123, 89, and 68 samples from the Brazilian, Ecuadorian, Belgian, and Portuguese participants respectively have been sequenced. These samples correspond to 218 high-grade lesions, 119 low-grade lesions, and 49 normal samples. All 14 genotypes were present in at least one sample. Considering only positive samples, in all countries, the majority of the samples had HPV coinfections (except for Ecuador: HPV 59 and HPV 16, and co-infections were found in the same number of samples). HPV 16 was most prevalent in Brazil and Portugal, versus HPV 31 in Belgium. Also were analyzed all the HPV types even those the types present in coinfection separately. It is possible to note that some trends persist. We found a high prevalence of HPV 52 in Portugal, which is important since it is not targeted by present vaccines. We also analyzed the types according to cytology results, in samples with high-grade lesions the most common HPV types until now are HPV negative and HPV 16.

Conclusions: It is still necessary to finish the analyses of the 1000 samples to confirm the genotypes frequency tendencies. Also, we will make other bioinformatic analyses like phylogenetic tree and lineages/sub-lineages classification.

Poster

Dr Monti Ermelinda
ermelindamonti@gmail.com - Italy

Dr Monti Ermelinda
ermelindamonti@gmail.com - Italy

#3505

Conservative treatment for cervical adenocarcinoma in situ: long term results

26 - Cervical neoplasia

Background/Objectives: To evaluate the effectiveness of conservative treatment for cervical adenocarcinoma in situ (AIS).

Methods: A retrospective study of Italian women with histologically confirmed AIS on cervical Loop Electrosurgical Excision Procedure (LEEP) specimen who were treated conservatively between 2008 and 2020. The main outcome investigated was the risk of recurrence defined as a subsequent finding of recurrent AIS or invasive adenocarcinoma in a long-term follow-up. All patients underwent colposcopy with endocervical curettage (ECC) and cytology every 6 months for the first two years after initial surgery and then annual cytology.

Results: Thirty women aged 26-51 years with histologically proven AIS on excisional specimen with negative margins, negative apex, and negative endocervical curettage were included. The median follow-up was 5.4 years. One woman had a recurrence of AIS after eight years of follow-up and underwent total hysterectomy. No invasive cervical disease was detected during surveillance.

Conclusions: Women with cervical AIS can be managed conservatively by an excisional procedure, provided that the margins are free and a close and long-term follow-up is guaranteed.

Oral presentation

Mr Kalteis Martin Simon
simon.kalteis@med.uni-heidelberg.de - Germany

Kalteis Martin Simon
simon.kalteis@med.uni-heidelberg.de - Germany

#3506

ESTABLISHMENT AND OPTIMIZATION OF A SOFTWARE-BASED 3D RECONSTRUCTION WORKFLOW FOR ORGANOTYPIC CULTURE MODELS

21 - New technologies

Kalteis M, Altmann J, Köhler R, Stark H, Von Knebel Doeberitz M, Prigge E

Background/Objectives: Three-dimensional (3D) in vitro model systems are increasingly used to study the effects of pharmacological substances in environments that aim to represent in vivo conditions more accurately than traditional 2D cell culture. We recently established a 3D organotypic culture (OTC) model mimicking HPV-induced precancerous lesions in order to evaluate the effects of new therapeutic approaches on lesion size and morphology in response to distinct treatment conditions. In order to characterize changes in tissue model architecture, we developed an open-source-software-based workflow that uses digital image stacks of whole-slide-scanned serial tissue sections to generate 3D renderings for visualization and analysis.

Methods: OTCs were formalin-fixed, paraffin-embedded and serially sectioned into 14 µm thick consecutive tissue sections, which were stained with Meyer's Hematoxylin and Eosin and subsequently scanned on a Hamamatsu S210 whole-slide-scanner imaging system. Tumor cell areas were detected by a custom-trained classifier using QuPath [1] and converted into binary masks. Images and masks were registered, stacked and color-deconvoluted in ImageJ [2,3] and finally fused and rendered as 3D volumes using Fluorender [4]. Tumor lesion volume was approximated by multiplying the segmented lesion area per image by the distance between two consecutive layers and summing the results for the complete stack in a script written for QuPath. Generalizability and adaptability were assessed by applying the workflow to generate 3D reconstructions of different tissue types.

Results: Using the newly established tissue processing and analysis workflow, we were able to reconstruct the 3D morphology of tumor lesions in our organotypic culture model. Different segmentation approaches were evaluated in QuPath and a pixel classifier trained on hand annotated data was selected as the best-performing approach in our dataset. Manual registration of image stacks was satisfactory but laborious and is a potential candidate for further automation. Rendered volumes can be viewed and interacted with in real time and export of images and video sequences is possible. Volumetric computation of tumor cell burden could be successfully performed.

Conclusions: We successfully established a workflow based on serial tissue sections and whole-slide-image stacks that enables 3D reconstruction and visualisation of OTC models. It allows for an in-depth study of the spatial configuration and size of HPV-induced lesions during the course of different treatment approaches. Using scriptable open-source software as building blocks makes our workflow cost-effective, flexible and easily adaptable to different tissue types and segmentation targets.

References: [1] Bankhead, P. et al. (2017). QuPath: Open source software for digital pathology image analysis. doi:10.1038/s41598-017-17204-5 [2] Schindelin, J. et al. (2012). Fiji: an open-source platform for biological-image analysis. doi:10.1038/nmeth.2019 [3] Bogovic, J. A. et al. & Saalfeld, S. (2016, 13-16 April 2016). Robust registration of calcium images by learned contrast synthesis. Paper presented at the 2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI) [4] Wan, Y. et al. & Hansen, C. (2012). FluoRender: An Application of 2D Image Space Methods for 3D and 4D Confocal Microscopy Data Visualization in Neurobiology Research. doi:10.1109/pacificvis.2012.6183592

Oral presentation

Dr Oaknin Ana
aoaknin@vhio.net - Spain

Monk Bradley J
Bradley.Monk@usoncology.com - United States

#3509

IMPACT OF CEMIPIMAB ON QUALITY OF LIFE, FUNCTIONING AND SYMPTOMS IN PATIENTS WITH RECURRENT/METASTATIC CERVICAL CARCINOMA: RESULTS FROM EMPOWER-CERVICAL 1/GOG-3016/ENGOT-CX9

08 - Immunotherapy - Immuno-oncology - New treatments

Background/Objectives: Cemiplimab significantly improved overall survival in patients with recurrent/metastatic (R/M) cervical cancer after first-line (1L) platinum-based chemotherapy (chemo) (NCT03257267; ESMO-VP-2021). We now report patient-reported quality of life (QoL), functioning, and symptoms from the clinical trial.

Methods: Patients with R/M cervical cancer who progressed after 1L chemo were randomised (1:1) to intravenous cemiplimab 350 mg every 3 weeks (N=304) or investigator's choice chemo for up to 96 weeks (N=304). At baseline (BL) and Day 1 of each 6-week treatment cycle (up to 16 cycles), patients were administered the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30. Mixed-effects repeated-measures models estimated overall least-squares (LS) mean change from BL for global health status (GHS)/QoL and physical functioning (PF). Results reported for the squamous cell carcinoma (SCC) and the overall (SCC + adenocarcinoma [AC]) populations were formally defined per statistical hierarchy for multiplicity adjustment. Other analyses for QoL were not adjusted for multiplicity, and nominal P values were reported.

Results: BL scores showed moderate to high functioning and low to moderate symptom burden with minimal differences across groups. For SCC, cemiplimab generally maintained GHS/QoL (LS mean change, 1.2 [95% CI -2.0, 4.3] vs -7.3 [95% CI, -11.5, -3.1]; difference, 8.5; one-sided P=0.00025) and PF (LS mean change, 1.0 [95% CI, -1.9, 3.9] vs -7.3 [95% CI, -11.1, -3.6]; difference, 8.4; one-sided P=0.00008) generally worsened scores with chemo. Findings were consistently better than chemo in the overall population. Significant differences in LS mean changes favoured cemiplimab on other functioning and key symptom scales in the overall and SCC populations (two sided nominal P<0.05). A similar trend was observed in the AC population (formal statistical test was not prespecified).

Conclusions: In patients with R/M cervical cancer after first-line treatment, cemiplimab provided significant benefit versus chemo in GHS/QoL and PF, along with other functioning and key symptom scales.

Poster

Ms Rivers Sydney
sydney.rivers@microbix.com - Canada

Zhelev Pavel
pavel.zhelev@microbix.com - Canada

#3512

PERFORMANCE OF FULL AND EXTENDED GENOTYPING ASSAYS USING A NEW HIGH-RISK HPV MULTIPLEX PANEL

09 - HPV testing

Rivers S, Vaughan L, Galbraith L, Rous C, Shah M, Zhelev P, Alagic A, Gerbaba T, Li J, Auluck J, Mikhael R, Luscher M, Hughes K

Background/Objectives: The development of extended (beyond HPV16/18) and full genotyping HPV assays has resulted in a demand for external control panels that can confirm assay performance across multiple high-risk genotypes. To address this need, we have designed and developed third-party HPV multiplex panels for use with HPV genotyping assays. These panels are designed to facilitate in-house assay verification, provide known positive material for personnel training and assist with routine Quality Control Assurance. Our controls are developed using full-length viral genomes grown in cell culture, which provide 100% clinical sample equivalence and are inactivated to meet Clinical Lab Biosafety requirements. The objective of this study is to confirm prototype panel performance and sample usability using extended and full genotyping assay workflows.

Methods: Microbix designed HPV16/18/45, HPV39/51/52, HPV31/33/66 whole-genome, inactivated HPV liquid multiplex samples formulated in PreservCyt®, which mimic diagnostic targets found in infected patient specimens. All samples were tested using the ELITe Panel (EliTech Group), Onclarity™ HPV assay (BD), and the Anyplex™ II HPV28 Detection Assay (Seegene) to ensure sample compatibility with partial, extended, and full genotyping assays, respectively.

Results: All samples showed acceptable performance when tested on the ELITe Panel, Onclarity™ HPV Assay, and Anyplex™ II HPV28 Detection Assay.

Conclusions: We have successfully designed high-risk HPV multiplex panels to support assay verification/validation, personnel training, and quality assurance. The samples exhibited acceptable performance using multiple Original Equipment Manufacturer platforms, demonstrating their potential use as cross-platform-compatible quality controls, verification panels and External Quality Assessment samples. Finally, our unique manufacturing process, which utilizes full-length viral genomes, provides a universal total nucleic acid (DNA/RNA) control compatible with any molecular assay target region.

Poster

Dr Leroy Charlotte
charlotte.leroy@gstt.nhs.uk - United Kingdom

Dr Leroy Charlotte
charlotte.leroy@gstt.nhs.uk - United Kingdom

#3519

Trans-oral robotic surgery for oropharyngeal squamous cell carcinoma, Guy's hospital institutional experience.

30 - HPV and oropharynx / Head and neck cancer

Faulkner J, Williams O, Sandison A, Jeannon J, Rovira A, Arora A

Background/Objectives: Trans-oral robotic surgery (TORS) has become established in the treatment of early oropharyngeal squamous cell carcinomas (SCC) and may play a key role in treatment de-intensification strategies in HPV related malignancy. Our centre instituted TORS as part of the OPSCC treatment paradigm in 2018 and report our experience.

Methods: This is a prospective case series of all patients who underwent trans-oral robotic surgery for primary treatment of head and neck malignancy between January 2018 and October 2021 at Guy's Hospital (London UK). All patients underwent comprehensive evaluation, staging (AJCC 7th edition) and multidisciplinary meeting (MDM) review. Primary TORS was offered to patients following MDM discussion and adjuvant radiotherapy (RT) or chemoradiotherapy (CRT) was given according to protocol. Specimen margins were analysed as per PATHOS protocol. Patients treated as part of ongoing clinical trials and "off trial" were included for analysis. Functional swallow outcomes were assessed via the MD Anderson Dysphagia Index (MDADI).

Results: 33 patients were treated from March 2018 to October 2021. The cancer location was 51.5% (17 patients) of tonsil, 27.3% of tongue base (9 patients), 15.2% of glosso-tonsillar sulcus and 6% of posterior pharyngeal wall. For the majority of patient (91%), the tumor was HPV positive. 82% of patients had an adjuvant treatment in which 22 patients received radiotherapy and 5 patients had chemoradiotherapy. The 6 other patients were treated only surgically. 6 patients (18%) had a post-operative bleeding but only two of them had to be managed surgically. The Mean MDADI functional outcome post-operative score at least 6 weeks after surgery was 73 with a minimum of 33 and a maximum of 100. 2 patient remained long term gastrostomy dependant, one had RT adjuvant treatment and the other one was a salvage case with free flap reconstruction.

Conclusions: Transoral robotic surgery is a safe and effective treatment method for oropharyngeal squamous cell carcinoma. To date oncological outcomes in our institution are promising. TORS for oropharyngeal SCC has good functional outcomes with an acceptable risk level but this study is limited by a small population. Further investigations with a comparison of functional outcomes in patient treated with standard chemoradiotherapy should be evaluated and the oncological outcomes reported after sufficient follow up.

Oral presentation

Ms Milbourn Hannah
hannah.milbourn@phe.gov.uk - United Kingdom

Milbourn Hannah
hanah.milbourn@phe.gov.uk - United Kingdom

#3521

EVIDENCE OF CROSS PROTECTION FROM THE BIVALENT HPV VACCINE IN YOUNG FEMALES IN ENGLAND

03 - Epidemiology and natural history

Background/Objectives: In 2008 England introduced a human papillomavirus (HPV) vaccination programme using the bivalent vaccine (Cervarix®) protecting against HPV16 and HPV18, changing to the quadrivalent vaccine (Gardasil®) in 2012. We present type-specific HPV DNA results for non-vaccine types from young women undergoing chlamydia screening and women attending for cervical screening who were offered the bivalent vaccine.

Methods: Residual vulvovaginal swab (VVS) specimens were collected from 16-24-year-olds attending for opportunistic chlamydia screening between 2010-2018. Residual Liquid Based Cytology (LBC) samples were collected from cervical screening attendances, at age 25-29, as part of the National Health Service Cervical Screening Programme (NHSCSP), from 2016-2019. Anonymised specimens were sent to the UK Health Security Agency's Virus Reference Department and underwent type-specific HPV DNA testing using an in-house multiplex PCR and Luminex®-based genotyping test. Analyses were restricted to women who would have been offered the bivalent, Cervarix, vaccine.

Results: 17,909 VVS specimens were collected from young women attending for chlamydia screening. The prevalence of HPV31, 33 and/or 45 infection, decreased between 2010-2018, from 6.5% to 3.1% (p value for trend <0.001) for 16-18 year olds and 8.6% to 2.8% (p value for trend <0.001) for 19-21 year olds. Prevalence of five high-risk (HR) HPV types (HPV31/33/45/52/58) declined from 19.9% to 7.3% (p value for trend <0.001) in 16-18 year olds and 21.6% to 9.1% (p value for trend <0.001) in 19-21 year olds. Significant declines were seen for each of these five types individually. 1,331 LBC samples were collected from women testing positive for any HR-HPV infection at cervical screening. Prevalence of HPV31/33/45 was slightly lower among individuals eligible for vaccination compared to those that were not eligible (20.2% (17.4 - 23.1) vs 24.8% (19.9 - 30.3)), but similar for the five HR-HPV types (45.1% (40.9 - 52.8) vs 46.8% (41.6 - 48.6)).

Conclusions: Following 10 years of high coverage vaccination, results from national surveillance of women eligible to receive the bivalent vaccine have shown declines in prevalence of HPV-16/18-related types (HPV31/33/45) and high risk types present in Gardasil 9 (HPV52/58), suggesting continuing evidence of cross-protection consistent with clinical trial evidence. Monitoring of changes in type-specific HPV prevalence is ongoing and remains relevant to inform and evaluate vaccine choice for this programme. Surveillance of type-specific HPV in cervical cancers is in progress - to provide the final evidence of cross-protection from the bivalent vaccine use in England.

Oral presentation

Mr Kusters Johannes
jeroen.kusters@rivm.nl - Netherlands

Kusters Johannes
jeroen.kusters@rivm.nl - Netherlands

#3523

TYPE-SPECIFIC CONCORDANCE OF HPV INFECTION BETWEEN THE GENITAL AND ANAL SITES FOR YOUNG WOMEN AND MSM

03 - Epidemiology and natural history

Background/Objectives: Concordant human papillomavirus (HPV) infection between genital and anal sites is common among women, and may be common among men who have sex with men (MSM). It is unclear what the risk factors are for type-specific anogenital concordance, but two potential risk factors could be receptive anal intercourse (RAI) and autoinoculation. Therefore, this study aimed to investigate the occurrence of type-specific anogenital concordance among young women and MSM visiting sexual health clinics in the Netherlands, and to identify risk factors for type-specific anogenital concordance.

Methods: Data from the PASSYON study, a biennial repeated cross-sectional study in the Netherlands, between 2009 and 2019 were used. Genital and anal HPV infections were identified for 25 HPV genotypes (SPF10-LiPA25). Cohen's kappa was used to assess the degree of type-specific concordance. Provided that at least moderate concordance ($\kappa > 0.4$) was found, risk factors for type-specific concordance were identified using logistic regression with generalised estimated equations. In this model, genital-only infections were compared to type-specific concordant infections. Receptive anal intercourse (RAI) was forced into the multivariable model to investigate its impact on concordance. All analyses were done for women and MSM separately.

Results: The study population consisted of 1,492 women and 614 MSM. Among women, type-specific anogenital concordance was common; 607 women (40.7%) had at least one type-specific concordant infection, and kappa was above 0.4 for 20 genotypes. Anogenital concordance was found most often for HPV-51 (10.7%); the highest kappa was found for HPV-6 ($\kappa = 0.60$). Among MSM type-specific anogenital concordance was not common; 64 men (10.4%) had at least one concordant infection, and kappa was ≤ 0.4 for all genotypes. Risk factors were determined for concordance among women. The multivariable model was based on 2,677 genital infections (genital-only: $n=1,622$; concordant: $n=1,055$), and the only significant risk factor for type-specific concordance among women was genital chlamydia (adjusted OR 1.5, 95% CI: 1.2 - 1.9). RAI was not associated with concordant infection (OR 1.1, 95% CI: 0.9 - 1.3).

Conclusions: Type-specific concordant HPV infection between the genital and anal sites was common among young women, and uncommon among MSM. The only significantly associated risk factor for concordance among women was genital chlamydia infection. As no other observed sexual or demographic risk factor was associated, autoinoculation seems a likely explanation for concordant HPV infection between genital and anal sites for women.

Oral presentation

Dr Van Keer Severien
Severien.VanKeer@uantwerpen.be - Belgium

Van Keer Severien
severien.vankeer@uantwerpen.be - Belgium

#3524

CLINICAL EVALUATION OF HPV-DNA TESTING IN HOME-COLLECTED FIRST-VOID URINE COMPARED TO PAIRED CERVICAL SAMPLES: UPDATE ON VALHUDES

13 - Self-sampling

Van Keer S, Latsuzbaia A, Vanden Broeck D, De Sutter P, Donders G, Doyen J, Tjalma W, Weyers S, Van Den Borst E

Background/Objectives: Urine collection is a non-invasive self-sampling method offering new perspectives in light of reaching women un(der)-screened for cervical cancer. Little evidence is available regarding the clinical accuracy for high-risk (hr)HPV testing on urine. VALHUDES was designed as a diagnostic test accuracy study aiming to compare the clinical accuracy to detect cervical intraepithelial neoplasia grade two or worse (CIN2+) of HPV assays on self-collected samples, including first-void (FV) urine, compared to HPV testing on paired clinician-collected cervical samples.

Methods: 523 women attending colposcopy due to a previous abnormal screen test were enrolled at five Belgian colposcopy clinics (NCT03064087). Women collected a FV urine sample (Colli-Pee with Urine Conservation Medium; Novosanis) at home the day before colposcopy. A cervical sample was taken by a gynaecologist prior to colposcopy (Cervex-Brush, Rovers in Hologic PreservCyt Solution). Disease outcome was assessed by colposcopy and histological assessment of biopsies if indicated. FV urine (index test) outcomes were available for Abbott Alinity m HR HPV (n=494), Roche Cobas 4800 (n=485), and Roche Cobas 6800 (n=491) with paired results using the same HPV test on the cervical samples (comparator). Manufacturer's instructions including assay cut-offs used for cervical samples were applied for FV urine as well. Differences in accuracy were assessed by relative sensitivity and specificity (FV urine vs cervical) with McNemar 95% CI.

Results: Paired samples were available from women aged 19 to 72 years (median age 40). HrHPV testing in FV urine with Abbott Alinity m HR HPV was similarly sensitive for CIN2+ (ratio 0.92; 95% CI: 0.84-1.02) and specific (ratio 1.03; 95% CI: 0.95-1.11) compared to testing cervical samples. Likewise, similar clinical accuracy was found in FV urine compared to cervical samples using both Roche Cobas 4800 (ratio [sensitivity for CIN2+] 0.98; 95% CI: 0.93-1.02, ratio [specificity <CIN2] 0.99; 95% CI: 0.91-1.09) and Roche Cobas 6800 (ratio [sensitivity for CIN2+] 0.96; 95% CI: 0.91-1.02, ratio [specificity <CIN2] 1.01; 95% CI: 0.93-1.09).

Conclusions: These results demonstrate that HPV-DNA based PCR testing is similarly sensitive (CIN2+) and specific (<CIN2) on home-collected FV urine compared to HPV testing on paired cervical specimens taken by a clinician. In agreement with previous clinical evaluations of validated HPV-DNA based PCR tests within VALHUDES, this offers new perspectives in light of reaching un(der)-screened women in organized cervical screening programs. As now, FV urine sampling is not only well-accepted by women, but has a proven similar clinical accuracy compared to cervical samples too using high-throughput HPV-DNA based PCR assays used in routine cervical screening.

Poster

Mr Irzaldy Abyan
abyan.irzaldy@gmail.com - Netherlands

Mr Irzaldy Abyan
abyan.irzaldy@gmail.com - Netherlands

#3531

FERTILITY IN WOMEN WITH CERVICAL CANCER AND PRECANCEROUS LESIONS IN RELATION TO CERVICAL SCREENING: A SWEDISH NATIONWIDE MATCHED-COHORT STUDY

41 - Fertility and HPV

Lei J, Andrae B, Wang J

Background/Objectives: Previous studies have shown that HPV infections and cervical treatment are associated with poorer obstetric and neonatal outcomes. It implies that the reproductivity of women with cervical cancer and precancerous lesions might be impaired. However, on the other hand, cervical screening might have potential positive impact on women's fertility through early detection enabling less-aggressive treatments. Using live birth as an indicator of fertility, this study aims to investigate the potential impact of cervical screening on women's fertility.

Methods: We performed a matched-cohort study using the Swedish national population and healthcare registers. Women with cervical precancerous lesions (n=32,614), screen-detected invasive cervical cancer (n=863), and symptom-detected invasive cervical cancer (n=488) were each matched to 5 women without these diseases at the diagnostic date, who have the same birth year and parity (n=169,816). They were followed until they give a live birth or other censored events. Kaplan-Meier and Cox regression analyses were performed.

Results: After adjustment of country of birth and educational level, the hazard ratios (HRs) of giving live birth for women with cervical precancerous lesions, screen-detected cancer, and symptom-detected cancer were 1.14 (95% CI, 1.11 - 1.16), 0.62 (95% CI, 0.52 - 0.73), and 0.16 (95% CI, 0.11 - 0.24), respectively, compared to women without known disease. Among nulliparous women, the corresponding adjusted HRs of giving live birth were 1.24 (95% CI, 1.21 - 1.28), 0.87 (95% CI, 0.69 - 1.10), and 0.20 (95% CI, 0.12 - 0.33), respectively.

Conclusions: Being diagnosed with cervical precancerous lesions through screening and most likely treated is not associated with lower incidence of giving live birth compared to the healthy population. Being diagnosed with invasive cervical cancer through screening therefore undergone less-aggressive treatment is associated with higher incidence of giving live birth compared to being diagnosed symptomatically. These indicate the potential positive impact of cervical screening on women's fertility.

Poster

Mrs Macios Anna
macios.anna@gmail.com - Poland

Mrs Macios Anna
macios.anna@gmail.com - Poland

#3533

THE EFFECT OF IMPLEMENTATION OF AN ORGANISED CERVICAL CANCER SCREENING PROGRAMME ON CERVICAL CANCER INCIDENCE AND MORTALITY IN POLAND

40 - Public health

Klimiuk A, Komerska K, Glinska P, Nowakowski A

Background/Objectives: The Organised Cervical Cancer Screening Programme (OCCSP) was launched in 2006/2007 in Poland and provides cytology sampling once in a 3 years interval for each insured Polish women between 25 and 59 years old. We have aimed to evaluate cervical cancer (CC) incidence and mortality trends to assess the potential effect of OCCSP implementation.

Methods: Data on number of CC cases and deaths in 1980-2018 were retrieved from the Polish National Cancer Registry and data on female population was generated from the Central Statistical Office of Poland. Standardized incidence rates (SIRs) and standardized mortality rates (SMRs) per 100,000 women were calculated using the world standard population. To identify the influence of screening on specific age groups three trend analyses were performed: for women aged 0+ (whole population), 25-59 (screening age) and 60+ (post-screening age). Joinpoint regression analysis was performed with the maximum number of 3 joinpoints to search for the point in time when trends had changed. Significance level of <0.05 was set as indicating statistically significant difference.

Results: Trends for SIR (Figure1): The SIRs were highest throughout the study period for women 60+. The annual percentage change (APC) was 1.2% in 1980-1985 ($p=.324$) and -2.0% in the subsequent period ($p<.001$). The significant downward trend was observed in cohort aged 25-59 years in 1991-2007 ($APC=-1.7\%$, $p<.001$) and was accelerated after 2007 ($APC=-5.6\%$, $p<.001$). Similar result was obtained for the whole female population, where the trend identified in the period of 1987-2007 ($APC=-1.7\%$, $p<.001$) was hastened afterwards ($APC=-4.4\%$, $p<.001$). Trends for SMR (Figure 2): Insignificant SMR trend among women aged 60+ in 1980-1991 ($APC=-0.4\%$, $p=.247$) was declining in subsequent period and slopes differed for 1991-2004 and 2004-2018 ($APC=-2.8\%$, $p<.001$ and $APC=-0.8\%$, $p=.002$, respectively). For women aged 25-59 the downward trend noticed in 1980-1993 ($APC=-0.9\%$, $p=.001$) was accelerated afterwards ($APC=-1.9\%$, $p<.001$ and $APC=-5.2\%$, $p<.001$ in 1993-2007 and 2007-2018, respectively). Similar pattern can be observed for women of all ages.

Conclusions: The implementation of the OCCSP in Poland seemed to have an impact on accelerating the downward trend for CC incidence and mortality among women in screening age of 25-59 years and in entire female population. However, the modelled trend on CC incidence in elderly women (60+) was not influenced by the OCCSP commencement and trend on CC mortality in this group was slowed down after the start of screening programme. The prolongation of screening age should be discussed in Poland to protect women 60+ who are at higher risk of CC incidence and death.

Oral presentation

Mr Boers Joachim
j.b.boers@erasmusmc.nl - Netherlands

Joachim Boers
j.b.boers@erasmusmc.nl - Netherlands

#3534

THE PROOF-OF-PRINCIPLE OF MARKER DISCOVERY FOR DIFFERENT GYNECOLOGICAL CANCERS BY A NOVEL METHOD FOR GENOME-WIDE DNA METHYLATION PROFILING (MED-SEQ)

18 - Methylation

Joachim B., Ruben B., Wilfred V., Henk V., Joost G., Wim Q

Background/Objectives: DNA methylation serves as an important marker for mis-regulation of gene expression in cancer, is applied to classify tumors and can predict disease outcome and treatment options.

Methods: We developed a novel method that facilitates genome-wide methylation marker discovery allowing successful identification of methylation changes associated with pre-cancer and cancer at very low cost. The assay involves isolation and purification of DNA from formalin-fixed paraffine-embedded (FFPE) or fresh biopsies (only 10-50ng DNA is needed). A DNA methylation dependent restriction enzyme digestion releases 32 base pair DNA methylated fragments that are sequenced by next generation sequencing. This Methylated DNA sequencing (MeD-seq) assay is very robust, allowing detection of DNA methylation at more than 50% of the 30 million CpGs present in our genome. With respect to costs and sequencing depth MeD-seq is superior to all available technologies and requires no DNA bisulphite treatment. MeD-seq is compatible with low amounts of DNA derived from solid tumor tissue enriched by laser capture microdissection (LCM) and liquid biopsies.

Results: We compared MeD-seq profiles of different types of cancers from vulva, cervix, endometrium, fallopian tube and ovary between cancers vs controls and cancers vs other cancers. Identification of Differentially Methylated regions (DMR) was achieved by comparing MeD-seq profiles using genome wide statistical testing using a sliding window approach, visualized through the Integrative GenomicsViewer (IGV) and subsequent identification of primer and probe regions for quantitative Methylation-specific PCRs (qMSP) to detect tumor-specific or general-tumor markers. In addition to DNA methylation data, MeD-seq generates sequencing data which enables the detection of Human Papilloma Virus (HPV) DNA incorporated in the genome of HPV-infected cells. Around half of the gynecological cancer types in our study are HPV-associated and we were able to detect HPV genomic integration and call HPV subtypes based on MeD-seq data.

Conclusions: MeD-seq is a reliable low-cost technology to establish genome-wide DNA methylation profiles of FFPE treated laser dissected material of cancer and controls and can be used to call DMRs for development of PCR-based assays. In addition to interrogating genome wide DNA methylation, MeD-seq is able to detect HPV integration in host genomes and able to call specific HPV subtypes.

Oral presentation

Dr Polonia-valente Rita
ritapoloniavalente@gmail.com - Portugal

Polónia-valente Rita
ritapoloniavalente@gmail.com - Portugal

#3538

VAGINAL INTRAEPITHELIAL NEOPLASIA: IS IT REALLY A RARE CONDITION? - A RETROSPECTIVE STUDY AMONG A COHORT OF A PORTUGUESE ONCOLOGY CENTER

27 - Vulvar diseases and neoplasia

Polónia-valente R, Vitorino A, Ferreira S, Nabais H

Background/Objectives: Vaginal intraepithelial neoplasia (VaIN) is considered the rarest form of intraepithelial neoplasia with an estimated incidence of 0.3 per 100,000 women in the United States, but likely its incidence is underestimated^{1,2,3}. These facts contribute for a challenging management of this lesion without international nor national guidelines. The aim of our study was to describe a cohort of VaIN cases in terms of management and outcomes to aware colposcopists for this condition due to its potential as a premalignant lesion to invasive vaginal cancer.

Methods: We analyzed a cohort of 28 cases with pathological diagnosis of VaIN at the Champalimaud Foundation Center, in Lisbon, Portugal, between 2014-2021 at the Colposcopy Unit of Gynecology Department. Since the diagnosis to follow up visits, we describe the cohort, the treatment/management selected and outcomes.

Results: Our cohort included 28 cases of VaIN diagnosed by pathological examination of vaginal biopsies with suspected HPV infection by colposcopy examination: 19 cases of VAIN1, 5 cases of VAIN2 and 4 cases of VAIN3, within this group there were 2 cases of vaginal invasive cancer. The median age was 50.9 ± 11.6 years old, 14 (50.0%) women were at menopause, 21 (75.0%) were HPV HR positive in cytology sample, only 7 (25.0%) had had a previous hysterectomy - 3 of them for malignant or premalignant cervix lesions and 4 women had concomitantly VIN lesions. Treatment selected for VaIN1 was expectant management 5 in 19 cases, laser Co2 ablation in 11 of 19 cases and 2 cases were treated with imiquimod intravaginally application followed by laser Co2 ablation therapy. In VaIN2 lesions - all five cases were submitted to laser Co2 ablation/excision therapy. In VaIN3 lesions - there were 4 cases: two treated with laser Co2 ablation/excision therapy and two treated with partial vaginectomy. In terms of treatment response of the 28 cases, 6 cases lesions recurred (3 VaIN3, 2 VaIN1, 1 Vain2) after first treatment - with a median time of 39 months (IQR [14-77]) until recurrence.

Conclusions: VaIN is an uncommon pre-malignant condition of the lower genital tract with a risk of progression to invasive cancer of up to 10% (4). Our study found 2 cases of invasive vaginal cancer concomitantly in 28 cases of VaIN (7%). VaIN is mostly associated to women an abnormal Pap smear who are posthysterectomy. Nonetheless we should not disregard this entity in cases of cervical abnormalities with normal colposcopy, as it could be not that uncommon.

References: 1- Sillman FH, Fruchter RG, Chen YS, et al. Vaginal intraepithelial neoplasia: risk factors for persistence, recurrence, and invasion and its management. *Am J Obstet Gynecol* 1997;176:93-9.; 2- Gurumurthy M, Cruickshank ME. 2012. Management of vaginal intraepithelial neoplasia. *Journal of Lower Genital Tract Disease* 16:306-312.) 3- Mahalakshmi Gurumurthy, Simon Leeson, John Tidy & Margaret E. Cruickshank (2019): UK national survey of the management of vaginal intraepithelial neoplasia, *Journal of Obstetrics and Gynaecology*, DOI: 10.1080/01443615.2019.1652887 4- Sopracordevole et al. " High-grade vaginal intraepithelial neoplasia and risk of progression to vaginal cancer: a multicentre study of the Italian Society of Colposcopy and Cervico-Vaginal Pathology (SICPCV)", *Eur Rev Med Pharmacol Sci* 2016 Mar;20(5):818-24

Poster

Mr Trimis Georgios
georgios.trimis@merck.com - Greece

Athanasopoulos Charalambos
charalambos.athanasopoulos@merck.com - Greece

#3539

GEOGRAPHIC DISTRIBUTION AND HPV TESTING METHODOLOGIES OF STUDIES REPORTING HPV ATTRIBUTABLE FRACTION IN ADVANCED STAGE HEAD AND NECK CANCERS: RESULTS OF THE 'ALARM' SYSTEMATIC LITERATURE REVIEW

30 - HPV and oropharynx / Head and neck cancer

Athanasopoulos C, Bencina G, Sabale U, Morais E, Georgilis A, Lazaridou M, Dimitriadis I, Poughias L

Background/Objectives: Human papilloma virus (HPV) is a potentially preventable causative factor of head and neck cancer (HNC) particularly in the oropharynx. In light of the challenges in the management of advanced stage [locally-regionally advanced (LA) and recurrent/metastatic (RM)] HNC and lack of information on HNC attributable to HPV, we systematically reviewed available literature to provide updated information on the proportion of HPV+ cases (i.e., HPV attributable fraction; HPV AF) in advanced stage HNC worldwide. Here, we present exploratory outcomes of regional diversities in data availability and distribution of HPV detection methods.

Methods: We searched on clinicaltrials.gov for interventional studies (IS; i.e., phase I-III trials) as well as MEDLINE (via Pubmed) and Embase databases for non-interventional studies (NIS) of advanced stage HNC published between January 1, 2010 and December 31, 2020. NCT numbers retrieved for IS were further searched on MEDLINE and Embase databases and ASCO/ESMO journals of congresses for abstracts/articles with available results. Studies were selected if initiated from January 1, 2010 onwards and had available HPV prevalence data among patients with LA and/or RM HNC, including oropharynx among head and neck subsites.

Results: A total of 81 studies (62 IS and 19 NIS) were included, contributing data from 53 countries for 9,607 cases of advanced stage HNC. Overall, 61 studies were single-country studies conducted in Northern America, Europe and Eastern Asia, five were multiple-country single-continent studies and the rest were multiple-country multiple-continent studies. In total, 44 studies included data from the USA, 16 from Germany, 15 from France, 13 from Spain, 12 from Italy, and 11 each from Belgium, Canada and the United Kingdom. HPV status was reported to be assessed solely based on p16 expression in 58.0% of the studies, another method in 8.0% and multiple methods in 8.6%, while the method of assessment was not specified in the remaining 24.7% of the studies. The distribution of HPV detection methods was similar when examined separately for IS and NIS.

Conclusions: Our results indicate there is considerable inconsistency in the availability of data on HPV prevalence in the last decade across countries and continents, suggesting further efforts for a better representation of all regions are needed. Additionally, though HPV detection was primarily based on p16 expression, methodological heterogeneity as well as literature gaps were observed, highlighting the necessity for implementing a consistent testing methodology to accurately and consistently assess HPV burden.

Oral presentation

Dr Vaughan Laurence
laurence.vaughan@bd.com - United States

Vaughan Laurence
laurence.vaughan@bd.com - United States

#3542

RISK STRATIFICATION OF HPV-POSITIVE WOMEN USING EXTENDED GENOTYPING AND CYTOLOGY - RESULTS FROM THE BASELINE ONCLARITY TRIAL

12 - Triage of HPV positive women

Vaughan L, Gary D, Parvu V, Andrews J

Background/Objectives: HPV screening is more sensitive than cytology but lacks specificity. Most cervical cancer screening programs triage women using HPV and cytology. The risk profiles of high-risk HPV types are very different, with seven 9-valent vaccine high-risk types contributing approximately 90% of all disease. Here we evaluate the ability of extended genotyping (beyond HPV 16/18) in combination with cytology to refer women to colposcopy.

Methods: We analyzed baseline data from the previously reported BD Onclarity PMA trial (1). Individual genotypes HPV16/HPV18/HPV31/HPV45/HPV51 and three grouped types HPV33_58 / HPV35_39_68 / HPV56_59_66) were combined with cytology classifications (NILM, ASC-US/LSIL, High-grade) to generate hierarchical risk strata for CIN2+ and CIN3+ disease and to refer women to colposcopy. Baseline HPV, cytology and histology information was used to generate Receiving-Operator-Characteristic (ROC) curves of Sensitivity versus 1-Specificity, and CIN2+ and CIN3+ risk profiles. Different colposcopy referral strategies were then compared using tiered risk profiles generated by HPV genotype and or cytology category. We also compared performance to that of the similarly designed dual stain [CINtec PLUS / dual stain (DS)] Roche IMPACT PMA trial (2)

Results: Tiered hierarchical groupings (in descending risk order) of HPV 16, HPV 18/31, HPV 52/33_58, and HPV45/HPV51/HPV35_39_68/HPV56_59_66 resulted in stepwise reduction in both CIN2+ and CIN3+ baseline risk across High-grade, ASC-US/LSIL, and normal cytology categories. CIN2+ and CIN3+ ROC curves of xGT + cytology triage options were similar. Cytology performance in both the Onclarity and IMPACT trials was less sensitive and less specific than extended genotyping options. Previously published DS triage with HPV16/18 genotyping performance mapped closely to both CIN2+ and CIN3+ ROC curves and fell between ASC-US/LSIL HPV33_58/HPV45/HPV52 and NILM/HPV33_58/HPV45/HPV52 extended genotyping options (Table 1).

Conclusions: Hierarchical extended genotyping risk groupings effectively triage patient risk, independent of cytology classification, and facilitates clinical action thresholds for colposcopy referral. When combined with cytology, extended genotyping offers the potential for improving triage of HPV+ women. Data from the BD Onclarity trial suggests that it outperforms cytology triage and may perform similarly to CINtec PLUS/dual stain triage using HPV16/18 genotyping. The implications for national screening programs are discussed. These findings warrant confirmation in head-to-head studies.

References: 1. Stoler MH, Wright TC, Jr., Parvu V, Vaughan L, Yanson K, et al. 2018. The Onclarity Human Papillomavirus Trial: Design, methods, and baseline results. *Gynecol. Oncol.* 149:498-505 2. Wright TC Jr, Stoler MH, Ranger-Moore J, et al. Clinical validation of p16/Ki-67 dual-stained cytology triage of HPV-positive women: Results from the IMPACT trial. *Int. J. Cancer.* 2021;1-11. doi:10.1002/ijc.33812 WRIGHT ET AL. 11

laurence vaughan

Oral presentation

Mrs Castriciano Santina
santina.castriciano@copangroup.com - Canada

Santina Castriciano
santina.castriciano@copangroup.com - Italy

#3544

VALIDATION OF MSWABTM MEDIUM FOR THE ELUTION OF FLOQSWABS[®] FOR HUMAN PAPILLOMAVIRUS (HPV) DETECTION ON SIX COMMERCIAL PCR-BASED HPV ASSAYS

09 - HPV testing

Santina C, Marco H, Ivy N, David H

Background/Objectives: Vaginal self-collection has been advocated to improve women's participation to Human Papillomavirus (HPV) based cervical cancer screening programs. Currently, self-collected swabs are eluted in alcohol-based media, which is flammable, toxic, and costly to transport, and the use of different elution volumes can result in sensitivity variability. A medium to elute self-collected swabs that support HPV stability at different temperatures and is compatible with clinically validated commercial PCR-based HPV assays is necessary. MSwabTM is a non-alcohol-based medium validated for viral nucleic acid amplification assays. Current evidence demonstrates that self-collected specimens are non-inferior to clinical collected specimens for the detection of HPV. The aim of this study was to evaluate the performance MSwabTM medium for the detection of HPV with six commercial clinical validated PCR-based HPV assays using the PROCEED^xTM FLOQ[®] Swab controls.

Methods: In this study PROCEED^xTM FLOQ[®] Swab controls (HPV16, HPV18, HPV45, and Negative), produced by MicroBix Biosystems (Mississauga, Ontario, Canada), were used to simulate the elution of self-collected FLOQSwabs[®] in MSwabTM Medium (COPAN Italia, Brescia, Italy). Five of each PROCEED^xTM FLOQ[®] Swab controls were eluted in 3 ml MSwabTM medium by swirling for 20 seconds before removing the swab. All samples were assessed for the presence of HPV genotypes using the Abbott Alinity m on the Alinity m, Abbott Realtime on the m2000, Qiagen NeuMoDx on the NeuMoDx 96, Roche cobas 4800 on the cobas 4800, Roche cobas on the cobas 6800, and Seegene HPV28 on the STARlet/CFX systems, assays that are clinically validated for HPV detection. Testing was undertaken at the VCS Pathology Molecular Microbiology Laboratory.

Results: HPV16, HPV18, HPV45, and Negative PROCEED^xTM FLOQ[®] Swab controls in MSwabTM medium obtained a 100% sensitivity, specificity, negative and positive predictive values on the Alinity m, Realtime, NeuMoDx, cobas 4800, cobas, and HPV28 assays.

Conclusions: Data obtained in this evaluation, using MicroBix HPV and Negative PROCEED^xTM FLOQ[®] Swab controls eluted in COPAN MSwabTM medium, demonstrated 100% concordant performance using the Abbott Alinity m, Abbott Realtime, Qiagen NeuMoDx, Roche cobas 4800, Roche cobas, and the Seegene HPV28 assays for HPV detection. These data support the use of FLOQSwabs[®] in MSwabTM medium for detection of HPV using commercial clinically validated PCR-based assays.

Oral presentation

Prof. Kldiashvili Ekaterina
e.kldiashvili@tma.edu.ge - Georgia

Iordanishvili Saba
research@tma.edu.ge - Georgia

#3547

rs2232365 Polymorphism of FOXP3 Gene as a Risk Factor of Cervical Cancer Development

02 - Viral and molecular biology

Kldiashvili E, Metreveli T, Iordanishvili S, Al-rustum S

Background/Objectives: Cervical cancer is the second common malignancy in women. This is an immunogenic malignancy, and high-risk human papilloma virus (HPV) subtypes may cause its development. Persistent infection of high-risk HPV subtypes may significantly facilitate the development of cervical intraepithelial neoplasia, which has been confirmed as a major risk factor of cervical cancer. Regulatory T cells (Treg cells) are a group of mature T cells generated in the thymus following the induction of peripheral naïve T cells. They are essential for the inhibition of immune overreaction induced damage, but over-production of Treg cells may block the protective immune response to infection and tumors. FOXP3 (Forkhead box protein P3) is a key transcription factor in regulatory T cells (Tregs), and has important roles in the immunosuppressive functions in Tregs. The role of FOXP3 gene polymorphisms in cancer patients is not determined till now.

Methods: The allele-specific polymerase chain reaction (AS-PCR) has been used for determination of FOXP3 gene rs2232365A/G polymorphism in collected plasma samples. A total 100 plasma samples have been collected from the patients of the Research Institute of Clinical Medicine (Tbilisi, Georgia) in May-August 2019. 90 samples have been collected from the patients with diagnosed cervical cancer (34 plasma samples) and cervical intraepithelial neoplasia (CIN, 56 plasma samples). 10 plasma samples have been collected from the patients negative for intraepithelial lesion or malignancy; these samples have been used as a control group. The communication with patients, collection and labelling of the plasma samples has been performed by the responsible medical staff of the Research Institute of Clinical Medicine. The plasma samples have been provided to the research group anonymously; determination of sample category (i.e., patient with diagnosed cervical cancer / patient with diagnosed CIN /control group) by labeling was impossible.

Results: The present study aimed determination of FOXP3 gene rs2232365A/G polymorphism in collected plasma samples of patients with diagnosed cervical cancer and CIN in comparison with control group. rs2232365A/G polymorphism has been detected in 75.33% of cervical cancer (25 cases, $p = 0.03$) and 64.29% of CIN (36 cases, $p = 0.02$). FOXP3 gene rs2232365A/G polymorphism has not been detected in control group ($p = 0.01$). Treg cells have the key role in the process of immune escape. Their development requires continued expression of Foxp3; attenuated Foxp3 expression results in its functional deficiency. By analysis of the obtained results, we assume that polymorphisms of the Foxp3 gene may contribute to the cervical cancer development.

Conclusions: The present study revealed the link of rs2232365A/G polymorphism of FOXP3 gene with cervical cancer development.

Oral presentation

Głab Grzegorz
grzegorz.glab@invitrogmw.pl - Poland

Dr Głab Grzegorz
gregglab@gmail.com - Poland

#3549

Clinical evaluation of the Mobile ODT AVE (AI) application VisualCheck 1.5 versus 2.0: comparison with human colposcopic assessment

22 - Artificial Intelligence

Background/Objectives: The use of artificial intelligence (AI) algorithms to analyze images of the cervix made with the EVA System Mobile ODT device creates new possibilities in the diagnosis of precancerous and cervical cancer in everyday practice. It can be used wherever a gynecological examination with cervical visualization can be performed and where there is access to Internet networks. Indeed, the system can be used not only by doctors skilled in colposcopy, but also by trained medical personnel with much less experience. The use of VisualCheck® (Mobile ODT, Tel Aviv) application allows for remote online "consultation" of captured images of the cervix after the application of 3-5% acetic acid. The algorithm result, based on hundreds of thousands of images collected in a cloud-based database has been compared to the assessments of experienced colposcopists, and correlation has been shown to be verified. The obtained result allows for the initial identification of the examined women as owners of physiological changes (not suspected to be cancerous) and those with suspected lesions, for whom further in-depth diagnosis is advised (colposcopy after biopsy, HPV HR DNA genotyping, CINTec, or LBC).

Methods: The aim of the study was to compare the diagnostic performance parameters of two versions of the VisualCheck 1.5 application from 2020 and the next, improved version of VisualCheck 2.0 from 2021. The algorithm result for both versions of randomized images from daily practice was compared to assessment of experienced colposcopists. All women identified by AI as suspected and false negative by colposcopist were verified by histopathologic biopsy and PCR DNA HPV HR 14 type genotyping. The sensitivity, diagnostic specificity and the value of Youden's index were calculated for two groups of patients similar in size.

Results: Results: In 2020, 204 results of the study evaluated by the VisualCheck 1.5 versions were assessed, achieving a sensitivity of 89,6%, specificity of 69,2% and a Youden ratio of 58,9%. In 2021 another 269 results of the analysis were analyzed by the modified version of the VisualCheck 2.0 application, obtaining a sensitivity of 92,2%, specificity of 79,4% and the value of Youden index of 77%.

Conclusions: - Both the VisualCheck 1.5 and 2.0 applications have a high diagnostic sensitivity of approx. 90%, which allows the method to be used as an effective primary screening tool. - The VisualCheck 2.0 application has a specificity, which significantly reduces the numbers of false positives compared to version 1.5. The increase in the value of the Youden's index from 58,9% to 77,7% confirms the greater usefulness of 2.0 version as a support for screening tests with the use of EVA System Mobile ODT in everyday practice. - The obtained results confirm the importance of further research on larger populations in other countries, and with biopsy serving as the reference standard in order to further improve and validate AI technology for cervical assessment with the EVA System.

Oral presentation

Mrs Rebolj Matejka
matejka.rebolj@kcl.ac.uk - United Kingdom

Rebolj Matejka
matejka.rebolj@kcl.ac.uk - United Kingdom

#3554

CYTOLOGY INTERPRETATION AFTER A CHANGE TO HUMAN PAPILLOMAVIRUS TESTING IN PRIMARY CERVICAL SCREENING: EVIDENCE FROM THE ENGLISH PILOT

10 - HPV screening

Rebolj M, Mathews C, Denton K

Background/Objectives: Overcalling of abnormalities and excess colposcopy referral of women without high-grade cervical intraepithelial neoplasia (CIN2+) have been major concerns for using cytology triage in women with positive high-risk human papillomavirus (HPV) tests. Previous studies comparing cytology evaluation with and without the knowledge of a positive HPV test tended to be small and used archived material, whereby any cytological abnormalities detected after a positive HPV test had been revealed did not affect the women's clinical management. We investigated the effect of revealing HPV positivity on cytological interpretation in the English HPV pilot study.

Methods: The pilot compared the routine implementation of primary cervical screening based on cytology, where HPV test results were not available prior to cytology reporting, with that based on HPV testing, where cytology was only required after positive HPV tests. Women were managed according to the results of the relevant screening and triage tests. We studied the proportions of women aged 24-64 referred to colposcopy directly (after abnormal cytology in the screening sample) vs. after early recalls at 12 and 24 months (after negative cytology in the screening sample but persistent HPV positivity), the detection of CIN2+, and the positive predictive value (PPV) of colposcopy for CIN2+.

Results: We included 931,539 women screened with cytology and 403,269 screened with HPV testing. Revealed HPV positivity was associated with a higher direct referral to colposcopy after any abnormality, ORadj: 1.16 (95% CI: 1.14-1.18), and was higher for women with borderline changes (ORadj: 1.35, 95% CI: 1.30-1.40). Laboratories with higher direct referral referred fewer persistently HPV-positive women after the early recall. The detection of CIN2+ after direct referral increased with an ORadj of 1.17 (95% CI: 1.13-1.20) for informed vs. uninformed cytology; the detection after borderline changes increased with an ORadj of 1.39 (95% CI: 1.28-1.51). For most age groups and grades of cytological abnormality, the PPV for CIN2+ remained comparable under both conditions of interpreting cytology. In women aged 50-64 with high-grade dyskaryosis, however, the PPV increased from 71% to 83% after revealing HPV positivity, ORadj: 2.05 (95% CI: 1.43-2.93).

Conclusions: Quality-controlled cervical screening programmes can avoid inappropriate overgrading of HPV-positive cytology.

Oral presentation

Dr Siegenbeek Van Heukelom Matthijs
m.l.vanheukelom@amc.uva.nl - Netherlands

Prof. Dr Schim Van Der Loeff Maarten F.
mschim@ggd.amsterdam.nl - Netherlands

#3559

HPV TYPE SPECIFIC E6, E7 ANTIBODIES IN HIV-POSITIVE MSM WITH ANAL SQUAMOUS CELL CARCINOMA UP TO 20 YEARS PRECEDING DIAGNOSIS: A CASE-CONTROL STUDY

20 - Serology

Siegenbeek Van Heukelom M, Jongen V, Waterboer T, Van Der Zee R, Quint K, Prins J, Schim Van Der Loeff M

Background/Objectives: Anal human papillomavirus (HPV) infection may lead to anal squamous cell carcinoma (aSCC). aSCC is rare, but is much more common among HIV-positive men-who-have-sex-with-men (MSM). Earlier studies have shown that a substantial proportion of oropharyngeal squamous cell cancer and aSCC cases are seropositive for early antigen 6 (E6) of HPV-16, often years prior to diagnosis. In HIV+ MSM, aSCC is frequently caused by other HPV-types than HPV16. We aimed to assess whether aSCC cases were more often seropositive for E6 and E7 of the causative HPV-type than people living with HIV (PLWH) without aSCC.

Methods: Eligible for inclusion were HIV-positive MSM diagnosed with aSCC, of whom stored serum samples were available at the date of diagnosis and at least one year prior to diagnosis. Serum samples from each calendar year up to 20 years prior to diagnosis were selected for testing. We selected controls (3:1 ratio) among PLWH without aSCC diagnosis, of whom at least 2 stored serum samples were available 10 years apart. All samples were tested for antibodies against E6 and E7 of 10 HPV-types (6,11,16,18,31,33,35,45,52,58). We identified the causative HPV-type for each aSCC by Laser Capture Microdissection of Formalin-Fixed Paraffin-Embedded specimens using SPF10-PCR/LiPA 25. We assessed the association of aSCC with E6/E7 seropositivity using logistic regression. We also analysed time of seroconversion in both cases and controls.

Results: We included 20 cases with aSCC, of which 11 were caused by HPV-16 (55%), two by HPV-18 (10%), two by HPV-52 (10%) and five by other types (25%). Two of 11 HPV-16 aSCC cases (18%) were ever seropositive for E6 and one for E7 (9%); no other cause-specific E6 and/or E7 seropositivity among cases was found. Seropositivity for E6 among the 60 controls varied from n=0 (HPV-6,18,33,52) to n=2 (HPV-16,31,45) (0-3%), and for E7 between n=0 (HPV-18,35,45) and n=1, except for HPV-31 (n=6). The Odds Ratio (OR) for E6 seropositivity detected at least once among cases compared to controls was 8.0 (95%CI 1.6-39.6); for E7 this was 2.4 (95%CI 0.5-10.9). ORs for seropositivity at time of diagnosis were 9.8 (95%CI 1.9-49.0) for E6, and 2.7 (95%CI 0.6-12.6) for E7. The two E6 seropositive HPV-16 cases seroconverted 2-5 years prior to diagnosis; the E7-seropositive HPV-16 case seroconverted 6 years prior to aSCC diagnosis.

Conclusions: HPV-type-specific E6 and E7 seropositivity prior to aSCC diagnosis in HIV-positive MSM was rare and lower than in other studies, but E6 seropositivity was significantly associated with aSCC. Seroconversion to E6 seropositivity occurred 2-5 years prior to diagnosis.

Oral presentation

Dr Reddin Ian
i.g.reddin@soton.ac.uk - United Kingdom

Reddin Ian
I.G.Reddin@soton.ac.uk - United Kingdom

#3565

INTEGRATED ANALYSIS OF CERVICAL SQUAMOUS CELL CARCINOMA COHORTS FROM THREE CONTINENTS REVEALS CONSERVED SUBTYPES OF PROGNOSTIC SIGNIFICANCE

02 - Viral and molecular biology

Chakravarthy A, Reddin I, Henderson S, Dong C, Kirkwood N, Jeyakumar M, Rothschild Rodriguez D, Gonzalez Martinez N, Mcdermott J, Su X, Egawa N, Fjeldbo C, Skingen V, Halle M, Krakstad C, Soleiman A, Sprung S, Ellis P, Wass M, Michaelis M, Lyng H, Fiegl H, Salvesen H, Thomas G, Doorbar J, Chester K, Feber A, Fenton T

Background/Objectives: Human papillomavirus (HPV)-associated cervical cancer represents one of the leading causes of cancer death worldwide¹. Although low-middle income countries are disproportionately affected, our knowledge of the disease predominantly originates from populations in high-income countries. The relationship, if any, between HPV type and cervical cancer prognosis remains unclear². Using data from 643 cervical squamous cell carcinoma (CSCC) tumours from USA, Europe and Sub-Saharan Africa, we set out to test the hypothesis that we could identify a set of transcriptional and epigenetic features associated with prognosis within CSCC and to establish whether it is also present in independent patient cohorts representing different geographical locations.

Methods: Unsupervised clustering of a discovery cohort from the USA (n=236) using transcriptional data was performed to identify possible subtypes of CSCC. A methylation signature derived from resulting subgroups was used to develop a support vector machine (SVM) classification model that allocated European (n=313) and Sub-Saharan (n=94) validation cohorts to a particular subtype. Prognostic differences between subgroups were calculated using Cox regression multivariate analysis controlling for stage, age, HPV type and treatment.

Results: We identified two CSCC subtypes (C1 and C2) with differing prognosis, independent of HPV type. The poor prognosis associated with C2 tumours in the discovery cohort from the USA was also seen in the European validation cohort. The hazard ratio for C2 tumour 5-year survival obtained from multivariate analysis was 2.56 (p = 0.002), consistent with that seen in the discovery cohort (HR = 2.44, p = 0.02). C1 tumours were largely HPV16-driven, displayed increased cytotoxic T-lymphocyte infiltration and frequently harboured PIK3CA and EP300 mutations. C2 tumours were frequently driven by HPVs from the HPV18-containing alpha-7 clade, harboured alterations in the Hippo signalling pathway and increased expression of immune checkpoint genes, B7-H3 (also known as CD276), NT5E (also known as CD73) and PD-L2 (also known as PDCD1LG2).

Conclusions: In conclusion, we identify two novel, therapy-relevant CSCC subtypes. The key characteristics of these subtypes are conserved across the three cohorts, from the USA, Europe and Sub-Saharan Africa, thus our findings are likely of broad significance.

References: 1. de Martel, C., Plummer, M., Vignat, J. & Franceschi, S. Worldwide burden of cancer attributable to HPV by site, country and HPV type. *International journal of cancer* 141, 664-670 (2017). 2. Galic, V. et al. Prognostic significance of adenocarcinoma histology in women with cervical cancer. *Gynecologic Oncology* 125, 287-291 (2012).

Oral presentation

Ms Suominen Helmi
helmi.suominen@tuni.fi - Finland

Suominen Helmi
helmi.suominen@tuni.fi - Finland

#3567

CHARACTERIZATION OF PERIPHERAL BLOOD T-LYMPHOCYTE SUBSETS AMONG MOTHERS WITH PERSISTENT GENITAL OR ORAL HPV16 INFECTION AND THEIR CHILDREN

05 - Immunology

Suominen H, Paaso A, Koskimaa H, Grenman S, Syrjänen K, Syrjänen S, Louvanto K

Background/Objectives: Cell-mediated immune (CMI) response appears to play a considerable role in influencing the outcome of an HPV infection and the failure to develop sufficient CMI can result in persistency and progression of the infection. CD4+ lymphocytes participate in initiating and maintaining immune response, but the role of CD8+ lymphocytes in HPV infection is less clear. Characterization of various T-cell subsets within in CD4+ and CD8+ populations according to immune activation and differentiation could provide more knowledge on the different outcomes of HPV infection.

Methods: A subset of 42 mothers and their children (n=28), from the Finnish Family HPV study, was included in these evaluations. The participants were divided in four groups based on the mothers' genital or oral HPV16 infection status during the six-year follow-up: persistent HPV16 oral and/or genital vs always HPV negative. Peripheral blood mononuclear cells (PBMCs) from previously cryopreserved venous samples were immunophenotyped using 23 markers analyzed in flow cytometry (BD LSRFortessa™ Cell Analyzer). T-cell subpopulations were defined by presence or absence of different cell surface markers to either early or late activated cells, memory or naïve cells and as differentiated or undifferentiated cells. Proportion of the CD4+ or CD8+ lymphocytes by their immunophenotyped subsets were compared between the groups: mothers, children, and mother-child dyads.

Results: The mean rank distribution of CD8+ memory cells (CD45RO+CCR7-CD8+ cell population) was significantly higher among mothers with persistent genital HPV16 infection than in those who were genital HPV16 negative ($p=0.048$). The median levels of both the antigen presenting CD4+ cells (HLADR+CD3+CD4+) and activated CD8+ cells (CD38+HLADR+CD3+CD8+) were significantly lower in the mothers with the persistent oral HPV16 infection than in those being always oral HPV16 negative ones ($p=0.038$ and $p=0.033$). Interestingly, among children significant higher mean levels of activated CD4+, CD8+ cells and circulating CD8+ memory cells were only seen among those whose mother had persistent oral HPV16 ($p=0.038$, $p=0.008$ and $p=0.0033$, respectively). The mother-child comparisons revealed several statistically significant differences both in CD4+ and CD8+ T-cell subsets and also as related to the mother's HPV status indicating that there is disparity in immunophenotype activation between children and their mothers.

Conclusions: Our results provide tentative evidence that the mother's persistent HPV infection can be associated with changes in peripheral blood T-cell distribution. Furthermore, the mother's HPV infection might result in immune alterations in her offspring, especially the oral infection.

Oral presentation

Ms Wähner Cornelia
cornelia.waehner@msd.de - Germany

Wähner Cornelia
cornelia.waehner@msd.de - Germany

#3569

HPV VACCINATION UPTAKE IN BOYS AFTER INTRODUCTION OF GENDER-NEUTRAL HPV VACCINATION IN GERMANY INCLUDING IMPACT OF COVID-19 PANDEMIC - A RETROSPECTIVE DATABASE ANALYSIS

06 - HPV prophylactic vaccines

Wähner C, Hübner J, Meisel D, Schelling J, Zingel R, Wölle R, Mihm S, Reuschenbach M

Background/Objectives: Since 2007 HPV vaccination in Germany has been recommended and funded for girls. In June 2018 a gender-neutral recommendation for adolescents 9-14 years (2 doses) with catch up to 17 years (3 doses) was published by the German Standing Committee on Vaccination (STIKO) and is part of mandatory funding since January 2019. Vaccination is provided by office-based physicians. The objective of this study was to monitor the uptake of HPV vaccination in boys in Germany. The data reported are an update with the most recent results, including impact of the COVID-19 pandemic.

Methods: Data were used from the IMS® Vaccine Analyzer database between January 2018 and August 2021. This database contains anonymized electronic medical vaccination records from a panel of office-based physicians (pediatricians, general practitioners, gynecologists) and was used for nation-wide projections of administered doses per month to vaccine-eligible adolescents of age 9-17 in Germany. Vaccine coverage rates for the eligible birth cohorts of the male population were estimated using German census data (DESTATIS).

Results: The number of boys (9-17 years) that received their first dose per month increased to the level in girls within four months after mandatory funding and reached a 2019 maximum of 53,139 in July. With the onset of the COVID-19 pandemic, a first drop in numbers for first doses in March 2020 was seen for boys and girls that however recovered quickly until May 2020 and numbers settled between around 33,000 and 55,000 until March 2021. Since April 2021 we have observed a continuous steep drop in administered doses per month with a minimum of 13,178 first doses for boys and 18,261 for girls in August 2021, corresponding to a reduction of 70.1% (for boys) and 50.0% (for girls) compared to the same month in 2019, respectively. Estimated vaccine coverage rates by the end of 2020 for 15- and 18 years old boys were 33.9% and 19.5%, respectively for the first doses and 16.9% and 7.6%, respectively for completed series.

Conclusions: The monthly uptake and annual coverage in boys strongly increased after mandatory funding in 2019 but vaccine coverage rates for boys were still low at the end of 2020. The numbers of administered doses of HPV vaccine decreased substantially among German adolescents during the COVID-19 pandemic, with largest impact in 2021 with a sharp drop by up to 70%. Due to the delay in data delivery, the most recent data is from August 2021 but the study is ongoing until December 2021 and updated results will be presented. To reverse the deficits on long-term health and economic consequences a sustained increase in HPV vaccination rates is required.

Oral presentation

Prof. Carozzi Francesca
frakaro@gmail.com - Italy

Carozzi Francesca Maria
frakaro@gmail.com - Italy

#3574

QUALITY ASSURANCE IN CERVICAL CANCER SCREENING : EXTERNAL QUALITY ASSURANCE PROGRAM (EQA) FOR HPV SCREENING WITH CYTOLOGY TRIAGE AND PAP TEST SCREENING

24 - Risk management

Carozzi F, Avveduto G, Burroni E, Cannistrà S, Matucci M, Paganini I, Sani C, Terreni A

Background/Objectives: Human papillomavirus (HPV) DNA testing has become routine in many diagnostic laboratories, particularly with changes from cervical cytology to HPV DNA as primary screening test. However, cytology continues to play a key role in HPV screening because all HPV-positive women undergo a Pap test, which in this context, is referred to as a Triage Pap test. At the same time in young women (under 30 years of age) the Pap test remains the optimal screening test. Therefore, there are currently two screening pathways differentiated by age. According to the European guidelines for quality assurance in cervical cancer screening, external quality assurance (EQA) is essential for the evaluation of laboratory performance to ensure the quality of both HPV testing and Pap testing as a triage test as well as a screening test.

Methods: In collaboration with the Tuscan Regional Center for laboratory quality assurance (Certified Accredia ISO 17043), we developed two EQA programs: one specific for screening with HPV including evaluation for HPV Test and Triage cytology and a specific program for liquid vaginal cytology only. The HPV program has been active since 2016 and the liquid phase Pap program since 2019. For HPV, the 9 samples are sent in 3 different shippings, and can be requested in either Preservcyt or Surepath but only laboratories using validated tests for screening can participate (C. Meijer et al 2009). The corresponding Triage cytology digitalized slide, and a selection of meaningful images are made available when a positive HPV results is provided. The liquid Pap test EQA program consists of 12 digitalized cytological (vaginal??) smears (??)slides accessible on a dedicated facility.

Results: Results from both EQA programs are made available to participants within 20 days e reports are sent via e-mail. For HPV EQA, in addition to the qualitative data, semiquantitative data of HPV as well as gene housekeeping, when available in the kit used, are also analyzed. The statistical indicators are calculated as mean, standard deviation (DS) and coefficient of variation (CV) of all results arrived and according to the method used by the laboratory. Triage cytology and screening cytology are classified according to Bethesda 2014 and for each exercise it is reported: the consensus cytology interpretation, the descriptive result sent by the laboratory, the number and percentage of results obtained from other laboratories. In addition, for screening cytology a score, based on a weighed concordance is also provided. The basic idea is to replicate the weighted Cohen's Kappa statistic principle (Cohen, 1968), i.e., underlying that some discrepancies are more severe than others. The number the participating laboratories in HPV & PAP Triage EQA is increased from n. 14 in 2016 to n. 54 in 2021. In 2021, laboratories participating in the liquid phase PAP VEQ were 47. Detailed results from both EQA programs will be presented and discussed

Conclusions: The EQA programs that we implemented for screening programs referral laboratories are open to all laboratories working in this field and focus on molecular and morphological tests used in screening for cervical cancer. To our knowledge this is the only experience in Europe and worldwide of a systematic EQA for HPV & triage cytology and screening cytology that laboratories can exploit to accomplish ISO 15189 accreditation.

Oral presentation

Mr Hesselink Bart
at.hesselink@self-screen.nl - Netherlands

Bart Hesselink
at.hesselink@self-screen.nl - Netherlands

#3576

VALIDATION OF THE CLINICAL PERFORMANCE AND REPRODUCIBILITY OF THE NEUMODX HPV ASSAY SELF-SAMPLE WORKFLOW

13 - Self-sampling

Background/Objectives: HPV testing on self-collected (cervico-)vaginal specimens (HPV self-sampling) has similar clinical accuracy for CIN2/3+ compared to clinician-collected samples when performed with a clinically validated PCR-based assay. Therefore HPV self-sampling is a suitable alternative primary screening method for routine cervical screening. However, validation of the HPV self-sampling workflow (i.e., collection device, sample preparation protocol, and HPV test) remains necessary, because use of an HPV test that is clinically validated for clinician-collected samples may not automatically result in high clinical accuracy when applied to self-collected specimens (Belinson et al, 2012). The NeuMoDx™ HPV Assay targets the E7 region of 15 (probably) high risk HPV (hrHPV) types (HPV16, -18, -31, -33, -35, -39, -45, -51, -52, -56, -58, -59, -66, -67, and -68) and confers partial genotyping by individually reporting HPV16 and HPV18, and concurrently detects the 13 other common high-risk types as a pool. The NeuMoDx™ HPV Assay has been clinically validated for use in cervical screening on clinician-collected cervical samples collected in both PreservCyt and SurePath collection medium using the international guidelines for HPV DNA test requirements for primary cervical screening. Recently, the NeuMoDx™ HPV Assay self-sample workflow has been developed. Performance data of the NeuMoDx HPV Assay self-sample workflow will be presented.

Methods:

Results:

Conclusions:

Oral presentation

Dr Tanaka Luana
luana.tanaka@tum.de - Germany

Tanaka Luana F.
luana.tanaka@tum.de - Germany

#3577

SMEAR HISTORY OF WOMEN WITH CERVICAL CANCER: AN AUDIT OF ROUTINELY COLLECTED SMEARS IN THE TEQAZ STUDY

03 - Epidemiology and natural history

Tanaka L, Ikenberg H, Klug S

Background/Objectives: From 1971, Germany offered an opportunistic cytology-based cervical cancer screening for women aged 20 years and older. In 2020, Germany switched to an organised cervical cancer screening programme, with women aged 20 to 34 years receiving cytology-based screening and those aged 35 years and older undergoing HPV-cotesting. Non-participation in screening is a major risk factor for cervical cancer, but some women who frequently participated in cytology-based cervical cancer screening still develop cervical cancer, which could indicate quality issues in cytology. The aim of this study was to analyse the smears history of women who developed cervical cancer screening (cases) in the TeQaZ case-control study.

Methods: TeQaZ is a case-control study on cervical cancer and participation in screening and quality of cytology. We recruited women with newly diagnosed cervical cancer (ICD-10 C53) and population-based controls between 2012 and 2016. We retrieved the smears collected and their results (routine assessment) in the ten years before diagnosis (cases) or study entry (controls) and sent them for an audit performed by an independent cytologist (first review). A second independent cytologist reviewed the smear in case of disagreement between the routine assessment and the first review. Independent cytologists were blinded to the routine assessment. For the present analysis, we selected cases for whom at least three smears were retrieved and reviewed.

Results: Thirty-two cases had at least three smears collected within the ten years preceding diagnosis, contributing with more than 150 smears for the audit. Most were diagnosed with squamous cell carcinoma and small tumours (T1). The median age at study recruitment was about 41 years. Almost half of women had received at least one false negative result, with some women having multiple smears incorrectly classified. A considerable proportion of women had only normal smears in the years preceding cervical cancer diagnosis, with some only receiving the first positive test, which led to cervical cancer diagnosis.

Conclusions: Our analyses of routinely collected smears of women who developed cervical cancer indicate issues with a high proportion of cases who had received at least one false negative result and with a considerable proportion of cases with no history of abnormal smears preceding a cervical cancer diagnosis. Considering that cytology will remain an important part of the screening and remains the method of choice for women aged 20 to 34 years living in Germany, strategies to improve cytology quality should be implemented.

Oral presentation

Ms Doshi Mishelben
mishelben.doshi@affinity-cmc.com - United States

Lily Wu
lily.wu@merck.com - China

#3580

IMMUNOGENICITY AND SAFETY STUDY OF THE 9vHPV VACCINE IN 9-45-YEAR-OLD CHINESE FEMALES

06 - HPV prophylactic vaccines

Liao X, Chen Z, Lv H, Wang S, Liang Z, Yan C, Yu W, Fu R, Shou Q, Chen Y, Song Y, Zheng M, Luxembourg A, Lily W

Background/Objectives: The 9-valent human papillomavirus (9vHPV; HPV6/11/16/18/31/33/45/52/58) vaccine was conditionally approved for Chinese women (16-26 years) in 2018 based on global data. This is one of two post-marketing commitment studies of 9vHPV vaccine in Chinese females. Dual-primary objectives of this study (NCT03903562) were to demonstrate non-inferiority of anti-HPV geometric mean titers (GMTs, 9-19 vs 20-26 years) and seroconversion (27-45 vs 20-26 years).

Methods: Participants (N=1990; 9-19 years: n=690; 20-26 years: n=650; 27-45 years: n=650) were enrolled to receive three doses of 9vHPV vaccine (Months 0, 2, 6). Serum samples were collected (Day 1, 1 month post-Dose 3) and tested by competitive Luminex immunoassay. For each HPV type, non-inferiority criterion for GMT required the lower bound of the two-sided 97.5% confidence interval (CI) of GMT ratio (9-19 vs 20-26-year olds) to be >0.67 , and for seroconversion, the lower bound of the two-sided 97.5% CI in differences (27-45 minus 20-26-year olds) was to be $>-5\%$. Immunogenicity analyses were based on the per-protocol population. Adverse events (AEs) were recorded.

Results: Robust anti-HPV GMTs were observed at 1 month post-Dose 3 in study participants. Nearly all ($\geq 99.8\%$) participants seroconverted to all vaccine HPV types. Prespecified statistical criteria for GMTs/seroconversion rates were met for all vaccine HPV types, demonstrating that antibody responses in 9-19-year olds and 27-45-year olds were non-inferior to those in 20-26-year olds (all P values <0.0001). Injection-site AEs and systemic AEs were reported by 43.3% and 50.9% of 9-19-year olds, 50.5% and 57.1% of 20-26-year olds, 43.8% and 43.4% of 27-45-year olds, respectively. No vaccine-related serious AEs and no deaths were reported.

Conclusions: 9vHPV vaccine-induced antibody responses were non-inferior in Chinese females 9-19 and 27-45 years of age versus 20-26 years of age. The vaccine was generally well tolerated in these populations.

Poster

Ms Doshi Mishelben
mishelben.doshi@affinity-cmc.com - United States

Dr Tu Yingmei
ying.mei.tu@merck.com - United States

#3581

QUADRIVALENT AND 9-VALENT HUMAN PAPILLOMAVIRUS VACCINE IMMUNOGENICITY AND SAFETY IN INDIAN STUDIES

06 - HPV prophylactic vaccines

Walia A, Luxembourg A

Background/Objectives: Efficacy, immunogenicity, and safety of quadrivalent human papillomavirus (qHPV; HPV6/11/16/18) vaccine and 9-valent HPV (9vHPV; HPV6/11/16/18/31/33/45/52/58) vaccine have been demonstrated. We report outcomes in Indian participants from three vaccine clinical trials. Objectives: Evaluate qHPV vaccine immunogenicity and safety in Indian girls (9-15 years) from an open-label, single-arm study (V501-029 [NCT00380367]; N=110), qHPV vaccine safety in Indian females (9-45 years) from a post-marketing safety surveillance study (V501-125; N=188), and 9vHPV vaccine immunogenicity and safety in a subgroup (n=225) of Indian girls (9-15 years), boys (9-15 years), and women (16-26 years) from a global study (V503-002 [NCT00943722]).

Methods: In V501-029 and V503-002, HPV vaccines were administered as three doses (Day 1, Month 2, Month 6). Vaccine HPV-type serum antibodies (Day 1, Month 7) were tested using competitive Luminex immunoassay; immunogenicity was analyzed in the per-protocol population. Injection-site adverse events (AEs), systemic AEs, and serious AEs (SAEs) were assessed. In V501-125, participants received qHPV vaccine during routine care and were actively surveilled for SAEs occurring within 30 days post-qHPV vaccination.

Results: Robust anti-HPV 6, 11, 16, and 18 responses were induced in V501-029; $\geq 97\%$ of participants seroconverted at Month 7 for each HPV type. In V503-002, anti-HPV 6, 11, 16, 18, 31, 33, 45, 52, and 58 responses were robust; $\geq 98\%$ of Indian participants seroconverted at Month 7 for each HPV type. In the V501-029 study, 58% of participants reported AEs (injection-site, 46%; systemic, 32%; mostly mild/moderate in intensity). In the V503-002 study, 70%, 58%, and 56% of Indian girls, boys, and women, respectively, reported AEs (injection-site, 56%-66%; systemic, 13%-28%; mostly mild/moderate in intensity). Across these two studies, no deaths, vaccine-related SAEs, or discontinuations due to AEs were reported. In V501-125, no SAE was reported.

Conclusions: In Indian populations, qHPV and 9vHPV vaccines elicited robust antibody responses and were generally well tolerated.

Oral presentation

Ms Doshi Mishelben
mishelben.doshi@affinity-cmc.com - United States

Luxembourg Alain
alain_luxembourg@merck.com - United States

#3584

IMMUNOGENICITY OF 1, 2, AND 3 DOSES OF 9-VALENT HUMAN PAPILLOMAVIRUS (9vHPV) VACCINE IN GIRLS 9-14 YEARS OF AGE

06 - HPV prophylactic vaccines

Luxembourg A, Bautista O, Saah A

Background/Objectives: In an open-label, randomized clinical trial, 2 doses of 9vHPV vaccine, given 6 or 12 months apart, induced non-inferior antibody responses in 9- to 14-year-olds compared with young women 16-26 years who received 3 doses (i.e., the age group and dose regimen in which vaccine efficacy was established). Thus, efficacy of 2 doses of the 9vHPV vaccine in 9- to 14-year-olds was inferred, which supported licensure of a 2-dose regimen in this population. We report results of exploratory immunogenicity analyses of this trial from girls 9-14 years who received 1, 2, or 3 vaccine doses.

Methods: Girls 9-14 years (N=753) were randomized in a 2:1:2 ratio to 0,6 cohort (2 doses given at 0 and 6 months), 0,12 cohort (2 doses given at 0 and 12 months), and 0,2,6 cohort (3 doses given at 0, 2, and 6 months). Young women 16-26 years (N=314) were enrolled to receive 3 doses. Blood was collected before dose 2 at Months 1 (0,6 cohort subset: n=50) and 6 (0,6 cohort), and Month 12 (0,12 cohort), and after the final dose at Months 7, 12, 24, and 36 (0,6 and 0,2,6 cohorts) or Months 13, 24, and 36 (0,12 cohort). HPV antibody responses to the 9 vaccine HPV types were assessed by competitive Luminex immunoassay (cLIA).

Results: Most (>99%) girls receiving 2 or 3 doses seroconverted for the 9 HPV vaccine types at 1 month post last dose. Anti-HPV geometric mean titers (GMTs) were highest at 1 month post last dose, decreased sharply during the subsequent 12 months, and then decreased more slowly. GMTs at 1 month post last dose in girls receiving 2 or 3 doses were similar to or greater than GMTs in young women receiving 3 doses. This trend was still observed through 24-30 months post last dose. A single dose of vaccine resulted in partial seroconversion (41% to 98% at 6 months after a single dose, depending on the HPV type, vs 97% to 100% at 6 months after completion of the 0,6 regimen) and lower GMTs than after 2 or 3 doses (6- to 14-fold lower than after the 0,6 regimen at 6 months post last dose). Percent seropositivity and GMTs after a single dose declined over time during the observation period (i.e., through 12 months post dose).

Conclusions: Immunogenicity assessment of 2- and 3-dose schedules of the 9vHPV vaccine suggests that 2 properly spaced doses elicit protective antibody response and durable protection. Antibody responses after a single dose were less robust than after 2 or 3 doses. It is unknown whether a 1-dose schedule elicits long-term protective antibody responses.

Oral presentation

Dr Vaernesbranden Magdalena R.
magvae@so-hf.no - Norway

Værnesbranden Magdalena R
magvae@so-hf.no - Norway

#3590

HUMAN PAPILLOMAVIRUS AT MID-GESTATION AND ADVERSE PREGNANCY OUTCOMES: A PROSPECTIVE COHORT STUDY FROM NORWAY AND SWEDEN

40 - Public health

Værnesbranden M, Staff A, Sjøborg K, Aaneland H, Granum B, Haugen G, Hedlin G, Hilde K, Lødrup Carlsen K, Nordlund B, Reh binder E, Skjerven H, Sundet B, Söderhäll C, Vettukattil R, Wiik J, Jonassen C

Background/Objectives: Human papillomavirus (HPV) is a common genital infection, with the highest prevalence during the reproductive years. The association between high-risk HPV (HR-HPV) infections and premalignant and malignant lesions of the cervix, as well as high HPV prevalence during pregnancy is well established. However, the association between HPV infections and adverse pregnancy outcomes is less studied and with conflicting results. This study aimed to investigate the association between HR-HPV infections at mid-gestation and pregnancy outcomes related to placental dysfunction.

Methods: We included 950 women with first-void urine samples collected at the time of enrollment at gestational age 16-22 weeks from the prospective multicenter cohort study of a general population in Norway and Sweden (PreventADALL study, ClinicalTrials.gov NCT02449850). HPV detection and genotyping were performed using the Seegene Anyplex II HPV28 detection assay. This real-time PCR tests presence of genotypes 12 HR-HPV, 8 probably/possibly HR-HPV, and 8 low-risk HPV (LR-HPV). Sociodemographic and pregnancy outcome data were collected using electronic questionnaires and medical records. Adverse pregnancy outcomes were identified as 1. Hypertensive disease of pregnancy (HDP); 2. Gestational diabetes (GDM); 3. Intrauterine growth restriction (IUGR). HDP outcomes included preeclampsia, gestational hypertension, superimposed preeclampsia, eclampsia and hemolysis, elevated liver enzymes and low platelets (HELLP) syndrome. IUGR was defined as $\leq 10\%$ birth weight percentile. We also constructed a composite variable for placental dysfunction related adverse outcomes (HDP, GDM, IUGR and preterm delivery).

Results: Mean age of the women included was 32 years, median pre-pregnant BMI was 25 kg/m². Prevalence of HR-HPV at mid-gestation was 231/950 (24%). The prevalence of HDP, GDM, IUGR and composite outcome variable was 83/950 (9%), 40/950 (4%), 67/950 (7%) and 168/950 (18%), respectively. We found no association between HR-HPV infections and any of these outcomes, HDP; (OR 1.10, 95% CI 0.64-1.86), GDM; (OR 1.30, 95% CI 0.59-2.86), IUGR (OR 1.70, 95% CI 0.88-3.30), or the composite outcome variable (OR 1.08, 95% CI 0.73-1.60).

Conclusions: In this general population cohort with low prevalence of adverse pregnancy outcomes, but high HR-HPV prevalence, we found no association between HR-HPV infections at mid-gestation and placental dysfunction related adverse pregnancy outcomes at delivery.

Oral presentation

Dr Hillen Christoph
c.hillen@uke.de - Germany

Hillen Christoph
c.hillen@uke.de - Germany

#3593

PREVALENCE OF GENITAL HPV INFECTIONS AMONG FEMALE TRANSPLANT RECIPIENTS

01 - HPV disease and COVID-19

Hillen C, Sachs C, Jaeger A, Prieske K, Kaufmann A, Lütgehetmann M, Schmalfeldt B, Sterneck M, Woelber L

Background/Objectives: Immunosuppressive therapy in organ transplant recipients is associated with a higher risk of developing anogenital (pre)malignancies related to Human Papillomavirus (HPV) infections. Previous studies on high-risk (hr) HPV prevalence among female transplant recipients have been inconsistent. Larger studies with accurate patient assessment are needed to assess hrHPV prevalence.

Methods: Female kidney or liver transplant recipients were HPV tested at the uterine cervix with Cobas® HPV Test (Roche Diagnostics) after at least one year of treatment. Further genotyping was performed by Anyplex™ II HPV28 (Seegene). All participants completed a questionnaire regarding medication, sexual behavior, transplant-related and medical history.

Results: 201 patients (median age 52; range 18-78 years) were included. 98 women had received a kidney transplant, 93 women a liver transplant, and 10 women simultaneously liver and kidney transplants. The median duration of immunosuppressive therapy was 9 years. 186/201 (93%) women had attended cervical cancer screening by PAP smear at least once a year before inclusion in the study and 25/201 (12.4%) were HPV vaccinated. To date, 144 cervical smears have been evaluated and 57 cervical smears and anal HPV test results will be available at data presentation. In 14.6% (21/144) of evaluated participants, the cervical HPV test was positive. HPV prevalence was 16.7% (12/72) in kidney and 12.5% (8/64) in liver transplant recipients. The most frequently detected genotype was HPV16 (43%). HPV positively tested women had an average of 6-lifetime sexual partners compared to 3 in HPV negative patients ($p=0.013$). 20/21 patients participated in the follow-up. Cervical hrHPV infection was confirmed in 17/20 (85%) patients, and 11/17 had anal hrHPV co-infection. Biopsies were obtained in 10/21 patients and three patients were confirmed to have high-grade dysplasia (1x cervical intraepithelial neoplasia (CIN) II, 1x CIN III, 1x vaginal intraepithelial neoplasia III).

Conclusions: Data indicate that the prevalence of cervical hrHPV in female transplant recipients is more than twice of that of the general population (6.2% in women above 30 years (Luyten et al., 2014)). Most women (93%) participate in regular cervical cancer screening, however, additional colposcopy might be beneficial to screen for (pre)invasive lesions of portio, vulva and vagina in HPV+ transplant recipients. Final results including anal swab will be available for a presentation showing hrHPV prevalence in the transplant group and the rate of anal co-infection.

References: [Luyten, A., Buttmann-Schweiger, N., Luyten, K., Mauritz, C., Reinecke-Lüthge, A., Pietralla, M., Meijer, C.J.L.M., Petry, K.U., 2014. Early detection of CIN3 and cervical cancer during long-term follow-up using HPV/Pap smear co-testing and risk-adapted follow-up in a locally organised screening programme. *Int. J. Cancer* 135, 1408-1416. <https://doi.org/10.1002/ijc.28783>]

Poster

Dr Pruski Dominik
dominik.pruski@icloud.com - Poland

Millert-kalinska Sonja
millertsonja@gmail.com - Poland

#3598

Incidence of high-risk HPV genotypes in the Wielkopolska region

14 - Genotyping

Pruski D., Millert-kalinska S., Musialowicz V., Lagiedo-Żelazowska M., Sikora J., Jach R., Przybylski M.

Background/Objectives: Human papillomavirus infection is one of the most common sexually transmitted infections. Persistent infection with high-risk HPV (hrHPV) is the direct cause of most cervical intraepithelial neoplasia and invasive cervical cancers. Temporary guidelines of the Polish Society of Gynecologists and Obstetricians recommend the usage of exclusively validated liquid media, tests for 14 hrHPV and the p16 / Ki67 test. Furthermore, current data indicate that genotypes 16 and 18 are assumed to be responsible for about 70% of cervical cancer cases. We aim to assess the most common hrHPV genotypes in the population of women from the Wielkopolska region.

Methods: We obtained material from 1200 women registered to Specialist Medical Practice in 2008 - 2021. The probe for the molecular test was collected with a combi brush and passed to the independent, standardized laboratory. HPV detection was done using PCR followed by DNA enzyme immunoassay and genotyping with a reverse hybridization line probe assay. Sequence analysis was performed to characterize HPV - positive samples with unknown HPV genotypes. The molecular test detected DNA of 41 HPV genotypes.

Results: The age range of the study group is from 18 to 85. We found that 52.4% of patients received HPV - positive test results. We detected that the following high-risk HPV types are the most common: 16, 52, 31, 51, 56 and 18. Of all HPV-positive patients, one-third have genotype 16. The frequency of genotypes 52 and 31 was the same and amounted to 11%. The frequency of genotypes 51, 56 and 18 accounted for 8.4%, 8.4% and 7.8%, respectively.

Conclusions: To our knowledge, this study is the most extensive assessment of high-risk HPV genotypes in Poland. Our results suggest that type-specific high-risk HPV DNA - based screening should focus on HPV types 16, 52, 31, 51, 56 and 18. It coincides with the Polish Society of Colposcopy and Cervical Pathophysiology recommendations. Moreover, such an analysis may indicate the potential groups with the highest risk of precancerous lesions and determine a new vaccination trend against specific HPV genotypes.

Poster

Dr Pruski Dominik
dominik.pruski@icloud.com - Poland

Dr Pruski Dominik
dominik.pruski@icloud.com - Poland

#3599

Level of L1 HPV serum antibodies after vaccination against HPV- a pilot study

05 - Immunology

Pruski D, Lagiedo-Żelazowska M, Millert-kalinska S, Musialowicz V, Sikora J, Jach R, Przybylski M

Background/Objectives: Cervical intraepithelial neoplasia (CIN) and cervical cancer are significant health problems globally. Vaccinations against HPV, which belong to the primary prevention of the development of precancerous lesions, significantly reduced the incidence of HPV-related changes in Australia and the United States. After treating CIN lesions or condylomata, some patients decide to vaccinate against HPV to develop L1 HPV antibodies for protection against re-infection.

Methods: We provide a pilot study in which we obtained material from women registered in Specialist Medical Practice in 2021. The experimental group constitutes patients who received three doses of vaccination against HPV after a diagnosis of either HPV infection or squamous intraepithelial lesions (SIL). In the Department of Immunology, we collected blood to the serum collected tubes (S-Monovette, Sarstedt), put centrifuge at 2.000 rpm for 20 minutes. ELISA microtiter plate was coated with recombinant VLP derived from HPV Type: 6, 11, 16 and 18. After stopping the reaction, we read the optical density (OD) at 450 and 620 nm. We set the seropositive cut points determined by calculated formulation from the manufacture at $OD > 0,303$.

Results: There were nine times more women with positive HPV tests in the experimental group than in the control 1 group. In contrast, no significant dependency between group and HPV test results was found in the experimental group vs control 2 group. Antibody level was higher in the experimental group than in both control groups. Antibody level divided by cut-off value (0.303) was also significantly higher in experimental group than both in control 1 group ($p < 0.001$) and control 2 group ($p < 0.001$). Women with LSIL diagnosis and women with HSIL or cancer diagnosis did not differ significantly in antibody level and antibody level divided by cut-off value.

Conclusions: The study aims to assess the level of anti-HPV (L1 HPV) antibodies in healthy patients and with diagnosed cervix pathology after vaccination. Further research will be necessary to enlarge the research group. The introduction of tests detecting L1 HPV antibodies may facilitate the evaluation of the effectiveness of vaccine programs. The analysis of specific types of immune disorders in patients diagnosed with HPV-related changes will facilitate the identification of groups of women with the highest risk of developing HSIL and, consequently, cervical cancer.

Oral presentation

Dr Thomsen Louise T.
litt@cancer.dk - Denmark

Thomsen Louise T.
litt@cancer.dk - Denmark

#3600

HPV PREVALENCE ACCORDING TO SOCIOECONOMIC CHARACTERISTICS IN A DANISH SCREENING POPULATION

03 - Epidemiology and natural history

Thomsen L, Sand F, Oernskov D, Munk C, Waldstroem M, Kjaer S

Background/Objectives: The Danish health care system is based on a principle of free-of-charge access for all residents, and nationwide cervical cancer prevention programs have been implemented. Nevertheless, social inequality is seen in cervical cancer incidence and mortality, with higher rates in women who have basic education, are unemployed or have low income. It is unknown whether there are also socioeconomic differences in the prevalence of HPV. The aim of this study was to investigate the prevalence of high-risk (hr) HPV according to socioeconomic characteristics in a Danish screening population.

Methods: We used data from a large pilot implementation of HPV-based cervical cancer screening. During May 2017-December 2020, women aged 30-59 years screened in the uptake area of the Dept. of Pathology, Vejle Hospital, Region of Southern Denmark, were offered primary HPV testing or primary cytology depending on municipality of residence. In the HPV group, ThinPrep liquid-based cytology samples were tested with the Cobas® HPV test. Using each woman's personal identification number, we obtained registry information on HPV vaccination and socioeconomic characteristics, including educational level, employment and country of birth. We calculated the hrHPV prevalence according to age and socioeconomic characteristics, and used log-binomial regression to estimate the relative prevalence (RP) of hrHPV infection, adjusted and stratified for age.

Results: We included 36,419 women with a valid HPV result. The majority (85%) were HPV unvaccinated, and those vaccinated had received the quadrivalent vaccine as adults (median age at vaccination: 26 years, range 17-57). The hrHPV prevalence decreased with age (30-39 years: 11.2%; 40-49 years: 8.0%; 50-59 years: 6.2%). Among women aged 30-39, the hrHPV prevalence was higher in those with basic compared with long education (12.8% vs 10.3%; RPage-adjusted=1.24; 95% CI, 1.06-1.45) and in women who were unemployed compared to employed (12.6% vs 10.8%; RPage-adjusted=1.13; 95% CI, 1.00-1.28). Immigrants from non-Western countries had lower hrHPV prevalence than Danish-born women (8.7% vs 11.5%; RPage-adjusted=0.76; 95% CI, 0.63-0.90). Similar patterns were seen in women aged 40-49 and 50-59 years.

Conclusions: In this largely unvaccinated Danish population, we found socioeconomic disparities in hrHPV prevalence, with higher prevalence in women with basic education and unemployment, and lower prevalence in non-Western immigrants. This may be caused by different sexual behavior patterns. Our findings support the need for promoting HPV vaccination and cervical cancer screening in socioeconomically challenged groups.

Oral presentation

Mr Munk Christian
munk@cancer.dk - Denmark

Munk Christian
munk@cancer.dk - Denmark

#3601

GIRLS-ONLY HPV VACCINATION - IMPACT ON THE HPV TYPE DISTRIBUTION IN YOUNG MEN

40 - Public health

Munk C, Reinholdt K, Kjaer A, Hemmingsen C, Oernskov D, Waldstroem M, Kjaer S

Background/Objectives: In Denmark, a girls-only HPV vaccination program has been in place since 2008-9. Subsequent catch-up programs for older girls/women mean that the HPV vaccination coverage in women born from 1985 until 2008 is around 70% across all birth cohorts. Since July 2019, HPV vaccination has been offered to boys aged 12 years. In Denmark, HPV vaccination is offered free-of-charge and in a call-recall system (invitation and reminder to those not responding). The objective of this study was to assess the overall HPV prevalence and the type distribution in younger men before the direct effect of HPV vaccination of younger boys becomes apparent.

Methods: We invited men who attended information days regarding military service at two barracks. In Denmark, all men are obliged by law to attend this information shortly after they turn 18 years. At random days from December 2019 to December 2020, 280 men were included in the study. They answered a shorter questionnaire with questions related to risk factors for HPV infection and a penile swab for subsequent HPV testing was collected. In conjunction with the penile swab, a physician assessed circumcision status and presence of genital warts. The penile swabs were stored at -80 degrees C. HPV testing was done with the Inno-Lipa Extra II (Fujirebio) which can detect 32 HPV types.

Results: The median age was 18 years with 94% being 18 to 20 years old. In all, 21 (7.5%) were circumcised and no one presented with genital warts. Forty men (14.3%) had not had their first sexual intercourse. The median number of sexual partners in the last 6 months was one partner with 76 (27.1%) having had two or more. Finally, 32 men (11.4%) reported to have been HPV vaccinated outside the program at their own expense. Altogether 129 men (46.3%) were HPV positive. Overall, 24.3% of the men had a high-risk HPV type, 23.9% had a low-risk type and 7.1% had an unidentifiable HPV type (HPV X). No infections with HPV types 6, 11, 16, 18, 31, and 45 were detected. The most frequent type was HPV 51 which was detected in 31 men (11.1%) followed by type 59 (8.2%) and type 66 (6.8%). Analyses related to e.g. vaccination status and sexual variables will be presented at the EUROGIN meeting.

Conclusions: The girls-only vaccination program in Denmark seems to a large degree having protected young men against the HPV types included in the licensed vaccines. The overall HPV prevalence is still high but consists largely of less carcinogenic HPV types.

Poster

Dr Di Costanzo Stefania
stefania.dicostanzo@copangroup.com - Italy

Remo Andrea
andrea.remo@aulss9.veneto.it - Italy

#3603

INTRODUCTION OF SELF-SAMPLING IN CERVICAL CANCER PREVENTION SCREENING PROGRAM BASED ON PAPILLOMAVIRUS (HPV) TEST IN THE ITALIAN VENETO REGION (ULSS 9 SCALIGERA): RESILIENCE DURING SARS-COV-2 PANDEMIC.

13 - Self-sampling

Remo A, Dalfior D, Gazzani D, Ciocchetta E, Martello G, Zanella C, Lucchi I, Zoboli E, Grego K, Zambelli L, Narne E, Colombari R

Background/Objectives: Vaginal self-sampling is a recently introduced approach to screen Papilloma virus (HPV) infection for cervical cancer prevention. This method is more convenient for patients compared to the established PAP test, since women can independently collect the sample at home. Several studies show that vaginal self-sampling is widely accepted by women across different cultural, religious, and socio-economic backgrounds and numerous countries are progressively introducing it in their screening programs. Veneto Region (Azienda Zero) developed a project to test vaginal self-sampling as a screening method for cervical cancer screening to increase participation in the non-respondent population. This study evaluates the outcomes of the first round of screening, considering the impact of the concurrent SARS-CoV-2 pandemic.

Methods: 9939 women between 30 and 64 years old in AULSS 9 Scaligera (including the province of Verona) were invited between January 10th 2021 and July 31st 2021 to a home self-sampling-based screening. The target population included respondent, non-respondent and women never invited for the cervical cancer screening. The self-sampling kit was sent to participants' addresses. It included a Copan Self Vaginal FLOQSwabs® REF#552C.80, instructions for use specifically designed for self-sampling and an acceptability questionnaire. After collection, women returned the sample to a local hospital or pharmacy. From there, samples were sent to the laboratory for analysis (Fracastoro Hospital, San Bonifacio). Dry swabs were eluted in 5mL of PreservCyt media and processed with Cobas 4800 (Roche diagnostics).

Results: 28% of invited women from ULSS 9 participated to the screening. Among these, 31% participated to a cervical cancer screening (HPV test or PAP test) for the first time and 8.5% tested HPV positive. From the questionnaire it emerged that 85% of participants prefer self-sampling at home compared to point-of care (POC) sampling and 77% scored self-sampling 4/5 on a 1 to 5 scale. Among women who refused to participate to the screening but filled the questionnaire, 40% would adhere to the program if offered self-sampling in POC.

Conclusions: Self-sampling-based HPV screening in ULSS 9 showed a good rate of participation, increasing the number of first-time participants to cervical cancer screening programs and demonstrating women's preference toward self-sampling at home. While SARS-CoV-2 pandemic halted most cervical cancer screening programs in Italy, this study demonstrates how self-sampling promotes cervical cancer screening resilience, maintaining a good participation rate despite lack of healthcare personnel and logistical difficulties.

Oral presentation

Dr Schmitz Martina
martina.schmitz@oncgnostics.com - Germany

Hansel Alfred
alfred.hansel@oncgnostics.com - Germany

#3604

DNA methylation markers for the detection of VIN, VAIN and vulva or vagina carcinoma

27 - Vulvar diseases and neoplasia

Wunsch K, Hansel A, Petzold A, Dürst M, Schmitz M, Runnebaum I

Background/Objectives: Vulvar and vaginal cancers together accounted for 1.3% of the global cancer incidence among women in 2020. The majority of vaginal cancers and a lesser proportion of vulvar cancers are associated with HPV. Rising incidence rates of vulvar cancer are observed especially in developed countries, and these seem to be mainly attributable to high-risk HPV-positive cases in younger women. Currently, no organized screening is performed in any country, so early detection is often an incidental finding during routine visit at the gynecologist. We analyzed the DNA methylation marker panel comprising GynTect, a CE IVD-marked assay for use in cervical cancer diagnostics, on vulvar and vaginal scrapes from women with vulvar and vaginal disease.

Methods: Scrapes from 43 patients, mean age 64, with vulvar (n=37) and vaginal (n=6) carcinoma as well as 25 vulvar intraepithelial neoplasias (VIN) and 8 vaginal intraepithelial neoplasias (VAIN) of different severity grade (17 VIN3, 5 VIN2, 3 VIN1 and 4 VAIN3, 3 VAIN2 and 1 VAIN1, mean age 58) were analyzed using GynTect, a DNA methylation marker assay comprising CpG islands in the promoter/5' regions of the human genes ASTN1, DLX1, ITGA4, RXFP3, SOX17 and ZNF671. Of the 25 VIN, 21 were HPV positive and 4 were HPV negative. All 8 VAINs were HPV-positive. Of the 37 vulvar carcinoma, 21 were HPV negative, whereas 16 were HPV-positive. All 6 vaginal cancers were HPV positive.

Results: Of the 33 VIN/VAINs, 21 (63.6%) were above the score defined for a GynTect positive. The 4 HPV-negative VINs were also GynTect-negative. All samples with a positive GynTect score were among the HPV positive samples, overall GynTect detection was 72.4% (21 of 29 samples) with a higher proportion of positively tested samples in the VIN group (17 of 21 samples; 80.9%). All 6 vaginal carcinoma cases (all HPV-positive) were GynTect-positive. Of the overall 37 vulvar cancers 28 showed a positive GynTect score. Of the 16 HPV-positive vulvar cancer scrapes, 15 (93.8%) were detected with GynTect, and of the 21 HPV negative cases, 13 (61.9%) showed a positive GynTect score.

Conclusions: As an increasing proportion of vulvar and vaginal cancer and their precancerous lesions are associated with high-risk HPV infection, we tested whether the six DNA methylation markers comprising the cervical cancer diagnostic GynTect, may be useful in diagnostics of these diseases as well. Interestingly, a high proportion of the samples were positive for GynTect, partially irrespective of the HPV status. Thus, in combination with HPV diagnostics, DNA methylation testing with these markers might be a promising tool for early detection of malignant vulvar or vaginal disease.

Oral presentation

Dr Dhillon Sharonjit Kaur
sharon5602@hotmail.com - Belgium

Dhillon Sharonjit Kaur
sharon.dhillon@sciensano.be - Belgium

#3606

CLINICAL VALIDATION OF HPV OncoPredict® SCR AND QT ASSAYS USING THE VALGENT-2 FRAMEWORK

09 - HPV testing

Dhillon S, Giubbi C, Cuschieri K, Cocuzza C

Background/Objectives: Background: Assessment of emerging HPV assays for their use in cervical cancer screening is vital. HPV OncoPredict® SCR and QT (Hiantis Srl) are two independent multiplex real-time PCR assays targeting HR-HPV E6/E7 DNA comprising a "partial" genotyping screening assay (SCR) and a "full" genotyping, normalized viral load (QT) assay. OncoPredict® SCR identifies HPV16 and HPV18 separately and 11 other hrHPV types in aggregate while OncoPredict® QT detects 12 high risk (HR) HPV types independently. The assays can be used either alone or in combination, as a two-step reflex testing allowing HR-HPV genotype-specific viral load determination in screen-positive samples. Quality controls for sample adequacy, the efficiency of nucleic acid extraction and PCR inhibition are included in the testing. Objectives: To evaluate the clinical performance of the HPV OncoPredict® SCR and QT using the international validation of HPV Genotyping Tests (VALGENT-2) framework.

Methods: Methods: The VALGENT-2 panel consists of 1,300 cervical liquid-based cytology (LBC) samples from women aged 20-60 attending the Scottish cervical cancer screening programme (1000 consecutive samples from routine screening enriched with 300 cytological abnormal samples). Disease was defined as histologically confirmed CIN2+ (n= 95 [denominator for sensitivity]), whereas two consecutive negative cytology results were accepted as a proxy for non-disease (n=721 [denominator for specificity]). Performance relative to GP5+/6+- PCR-EIA (standard comparator test) was assessed by a non-inferiority test.

Results: Results: The relative sensitivity and specificity for CIN2+ of OncoPredict® SCR vs GP5+/6+-PCR-EIA were 1.01 (95% CI, 0.99-1.03) and 1.02 (95% CI, 1.00-1.04). The results for OncoPredict® QT vs GP5+/6+-PCR-EIA were similar with a relative sensitivity of 1.01 (95% CI, 0.99-1.03) and relative specificity of 1.03 (95% CI, 1.0-1.06). The p-value for all the tests of non-inferiority was $p \leq 0.001$. Both assays also showed non-inferior accuracy when restricted to women aged 30 years or older.

Conclusions: Conclusions: HPV OncoPredict® SCR and QT assays fulfil the clinical accuracy criteria for use in cervical cancer screening. Acknowledgements: HPV OncoPredict® assays were developed as part of a European Commission funded SME Instrument Phase 2 Project (Grant agreement ID: 806551).

Oral presentation

Mrs Inturrisi Federica
f.inturrisi@amsterdamumc.nl - Netherlands

Inturrisi Federica
f.inturrisi@amsterdamumc.nl - Netherlands

#3607

FOURTEEN-YEAR RISK OF CERVICAL PRECANCER IN HPV-NEGATIVE WOMEN, STRATIFIED FOR PREVIOUS HPV AND CYTOLOGY RESULTS

24 - Risk management

Inturrisi F, Rozendaal L, Heideman D, Meijer C, Berkhof J

Background/Objectives: HPV-based screening programs still use one-size-fits-all protocols but efficiency and efficacy of programs may be improved by stratifying women based on previous screening results.

Methods: We collected 19 years of follow-up of women aged 29-61 from the POBASCAM HPV trial who received cytology and HPV co-testing at enrolment and after 5 years (blind HPV at enrolment in the control group). We selected 18,448 women who tested HPV-negative at 5 years and stratified them according to the preceding HPV and cytology results. We excluded women with a CIN2+ or hysterectomy between enrolment and 5 years. For each subgroup, we calculated 14-year CIN3+ and CIN2+ risk by Kaplan Meier method. Relative risks (RR) were calculated with a HPV-negative test as reference category.

Results: HPV-negative women had a 14-year CIN3+ risk of 0.48% (95% confidence interval (CI) 0.37-0.62). The CIN3+ risk among HPV-negative women was decreased in women who in the previous round had a negative HPV test (relative risk 0.90, 95% CI 0.81-0.99) or a negative co-test (0.90, 0.80-1.00) and markedly increased in women who in the previous round had a positive HPV test (4.92, 2.62-9.21) or a positive co-test (3.50, 1.93-6.36). The CIN3+ risk was not influenced by the previous cytology result. The CIN3+ risk among HPV-negative women was increased after both a previous HPV16-positive test (8.13, 3.15-20.96) and a previous HPV-positive/HPV16-negative test (3.98, 1.67-9.49). For end-point CIN2+, findings were similar except that the CIN2+ risk was increased after previous abnormal cytology (3.47, 2.01-5.99).

Conclusions: Long-term risk of CIN3+ among HPV-negative women were very low when there was no positive HPV test result in the previous screening round 5 years earlier, indicating that HPV results from multiple screens could be used for risk-stratification.

Figure 1. Cumulative risk of CIN3+ and CIN2+ in HPV-negative women per preceding test results after three screening rounds (14 years).

Oral presentation

Dr Jentschke Matthias
Jentschke.Matthias@mh-hannover.de - Germany

Jentschke Matthias
Jentschke.Matthias@mh-hannover.de - Germany

#3608

HaSCo Study: First results of a pilot study for systematic HPV self-sampling for non-responders to the cervical cancer screening program

13 - Self-sampling

Jentschke M, Gross S, Steinkasserer L, Neuhaus B, Geyer S, Koch A, Hillemanns P

Background/Objectives: In this pilot study we evaluate a systematic approach towards human papillomavirus (HPV) self-sampling for non-responders to the German cervical cancer screening program. In 2020, the renewed organized screening program with co-testing (HPV test + cytology) in 3-yearly intervals was started for all women 35 years and older. In the past, the participation rate was around 70% in a time frame of three years. The non-participants are at higher risk to develop invasive cervical cancer. The willingness to participate can be improved by offering self-sampling devices for home-based use. Based on this information, the Hannover Self-Collection study was started to examine the response rate and practicability of a systematic self-sampling approach.

Methods: 20.000 women aged 30 to 65 years living in the city and region of Hannover, Lower Saxony were randomly included. 10.000 women directly received a self-sampling kit, the other 10.000 a letter of information and option to participate in the study (opt-out vs. opt-in strategy). Stratifications were made by age (7 cohorts) and area of living (city vs rural). Women tested positive for high-risk HPV (PCR-based HPV assay) were prompted to get a cytological smear by their gynecologist. Women with normal cytology will be re-checked after 6 months. Suspicious cytology results lead to an immediate colposcopy.

Results: The first stage of the study is executed between January and March 2022. First results will be presented at the congress.

Conclusions: To get hold of non-responders to cervical cancer screening programs, self-sampling for HPV is a promising option. Aim of this study is to generate an overall recommendation to improve cervical cancer screening in Germany, especially for non-responders. This study is supported by Deutsche Krebshilfe.

Oral presentation

Ms Pedersen Helle
Helle.Pedersen.03@regionh.dk - Denmark

Pedersen Helle
Helle.Pedersen.03@regionh.dk - Denmark

#3610

CAN HPV GENOTYPING AND CYTOLOGY TRIAGE FACILITATE HPV SCREENING OF YOUNG WOMEN 23-29 YEARS OLD?

10 - HPV screening

Pedersen H, Tønnes Pedersen B, Korsgaard Andreasen E, Bonde J

Background/Objectives: Primary HPV screening of young women have been advised against on the premise that the high HPV prevalence in this group would lead to unnecessary referrals and treatment. The advent of the childhood HPV vaccinated women in the screening program combined with new molecular approaches to HPV screening could change this dynamic in favor of HPV screening over the current cytology screening offer. Here we present HPV prevalence and genotype frequencies in women age 23-29 attending the organized cervical cancer screening program in Denmark. The data is used to model referral outcomes using HPV screening with HPV genotype and cytology triage as currently used in Denmark for women age 30-59 year.

Methods: 1000 SurePath cervical samples from women age 23-29 attending the cervical cancer screening program in the Capital Region of Denmark were randomly collected in Q3, 2021, tested by the BD Onclarity HPV assay, and combined with routine cytology outcomes retrieved from the National Pathology Database. Referral rates for the 23-29 years old were modelled upon the Danish algorithm for primary HPV screening for women 30-59 years old. Here, HPV positive w. HSIL/AGC/ASC-H/AIS and HPV positive w. ASCUS/LSIL and either of HPV16, 18, 31, 33, 52 are referred to colposcopy. HPV positive w. ASCUS/LSIL and either of HPV35, 39, 45, 51, 56, 58, 66 or 68 as well as HPV positive w. normal cytology independent of genotype are referred to re-test in 12 months. All HPV negative are returned to next screening round. The study is a quality development study approved by AHH-Hvidovre Hospital.

Results: Overall HPV positivity was 25% (N=252) for women age 23-29. Genotypes ranked by frequency was HPV18 (0.8%), HPV16 (2%), HPV31 (5%), HPV45 (6%), HPV33/58 (15%), HPV51 (16%), HPV52 (25%), HPV35/39/68 (32%), and HPV56/59/66 (35%). Overall, 3.5 % were HPV positive≥HSIL (N=16) or ASCUS/LSIL and either HPV16, 18, 31, 33, 52 (N=19); 1.5% were HPV positive w. ASCUS/LSIL and either HPV35, 39, 45, 51, 56, 66, 68 (N=15), and 20% (N=201) had normal triage cytology independent of genotype. One sample was invalid. Applying the described algorithm, direct referral rate to colposcopy amongst the 23-29 years would be 3.5% compared to 1.9% with the current cytology algorithm for this age group. Overall, 4.3% of all tested would be referred to a re-test in 6 months with the cytology algorithm.

Conclusions: HPV16 and 18 is close to be eradicated amongst 23-29 year old women, and the majority of HPV positive screening findings are one or more of HPV 35,39,56,59,66 and 68, generally considered so low in oncogenic potential that they do not merit immediate follow-up. Compared to current cytology screening, we conclude that difference in referral rate is limited when utilizing a HPV screening with genotyping and cytology triage and thus could make primary HPV screening feasible for women age 23-29 year offering the general benefit of improved CIN2/3 detection by HPV screening over cytology also for this age group.

Oral presentation

Ms Checchi Marta
marta.checchi@phe.gov.uk - United Kingdom

Checchi Marta
marta.checchi@phe.gov.uk - United Kingdom

#3611

POST-IMMUNISATION PREVALENCE OF HPV INFECTION IN GAY, BISEXUAL AND OTHER MEN WHO HAVE SEX WITH MEN ATTENDING SPECIALIST SEXUAL HEALTH SERVICES IN ENGLAND

03 - Epidemiology and natural history

Checchi M, Panwar K, French C, Beddows S, Hickman M, Soldan K

Background/Objectives: Human papillomavirus (HPV) vaccination for gay, bisexual and other men who have sex with men (MSM) aged up to 45 years attending specialist sexual health services (SHSs) and HIV clinics began in England as a pilot in June 2016, with national roll-out from April 2018. Reported first dose vaccine uptake from pilot start to end December 2019 was 40.8% at specialist SHSs and 36.2% at HIV clinics, though under-reporting is likely, with a further (second dose) recorded for 56.5% and 62.2% of these MSM, in specialist SHSs and HIV clinics respectively. Multi-site surveillance of type-specific HPV DNA prevalence in MSM attending SHSs in England was initially conducted between 2016-19 as a baseline survey. A post-immunisation survey is currently being conducted to evaluate the impact of the national HPV vaccination programme in MSM.

Methods: Anonymised residual rectal swab specimens from MSM aged ≤ 45 years undergoing chlamydia and/or gonorrhoea screening at a selection of specialist SHSs are currently being collected. Specimens will be linked to data reported to the GUMCAD STI Surveillance System (demographics, prior STI and HIV diagnosis, and HPV vaccination record). Specimens will be tested for type-specific HPV DNA (including 13 HR types as defined by IARC) using an in-house multiplex PCR and Luminex-based genotyping test. We will compare the post-immunisation prevalence of HPV16/18/6/11 to the baseline prevalence in vaccinated versus unvaccinated MSM. Results will be stratified by demographic characteristics and prior STI and HIV diagnosis.

Results: (Not yet available at abstract submission - full results will be presented at EUROGIN) Baseline survey findings indicate that while prevalence of HR-HPV types was high at 50.9% (95% CI 49.3-52.6; n=1858), most MSM eligible for MSM vaccination were not currently infected with an HPV vaccine type (baseline HPV16/18/6/11 prevalence of 35.9% and HPV16/18 prevalence of 19.5%). Based on an expectation of at least a 20% reduction in HPV vaccine types in vaccinated MSM, a minimum of 3,000 rectal swab specimens are being collected from laboratories both in and outside of London. Type-specific HPV DNA testing for all HPV types is expected to be complete by April 2022.

Conclusions: Our data on the post-immunisation prevalence of HPV in MSM will provide the first evidence of the impact of national HPV vaccination in MSM in England and be an early indicator of the programme's success, and possibly identify disparities in the benefit from this selective vaccination programme. Expected reductions in infection prevalence predict that vaccination has the potential to trigger declines in HPV-related disease among MSM, including genital warts as an early disease outcome and anal cancers in future.

Oral presentation

Mr Forslund Ola
ola.forslund@med.lu.se - Sweden

Mr Forslund Ola
ola.forslund@med.lu.se - Sweden

#3612

Co-testing in cervical screening among 40-42 year old women is unreasonable

10 - HPV screening

Forslund O, Borgfeldt C, Leksell A

Background/Objectives: The screening program for cervical cancer in Sweden, recommends use of primary HPV screening for women > 30 years to 65 years. Co-testing with both HPV-analysis and cytology is recommended at the first screening after the age of 40. In order to fulfil co-testing, all screening women within the Region of Skåne at the age of 40-42 years were co-tested. The aim of the audit was to investigate the proportion of severe dysplasia as diagnosed by cytology and histological follow up among women with Aptima HPV negative tests. Also, the cost of the added co-testing cytology to the HPV primary screening program was calculated.

Methods: The local cytology registry was used to identify 40-42 year old women who attended screening and were co-tested during 4 years from January 2017 to December 2020. The Aptima HPV mRNA assay detects 14 HPV types and for Aptima HPV negative women with high-grade cytology or histological HSIL we performed extended HPV typing, for 40 HPV types, by the use of PCR using modified GP5+/6+ (MGP) primers followed by a Luminex assay. To estimate the added cost of cytology to identify each histology confirmed cervical HSIL case among Aptima HPV negative women, we used the current cost of EUR 21.2 per cytology evaluation at our laboratory.

Results: Out of 19,599 women, 5.8% (1137/19,599) had abnormal cytology. Among Aptima HPV negative women, 0.11‰ (2/18,132) had histologically confirmed HSIL. One of the women was infected with HPV18 and the other with HPV73 at the diagnosis of HSIL. The calculated cost to find one HSIL, via adding cytology to HPV negative cases, was approximately EUR 200,000.

Conclusions: The clinical benefit of a single co-test of cytology at 40-42 years added to an HPV-based screening program appears hesitant and economically unreasonable.

Oral presentation

Dr Gori Silvia
silvia.gori@iov.veneto.it - Italy

Gori Silvia
silvia.gori@iov.veneto.it - Italy

#3614

CONSERVATIVE MANAGEMENT OF WOMEN WITH CIN2 LESIONS ENROLLED IN A PROSPECTIVE MULTICENTRIC STUDY: AGE PREVALENCE AND APPLICABILITY.

23 - Diagnostic procedures / management

Gori S, Frayle H, Pagan A, Solda' M, Romagnolo C, Insacco E, Laurino L, Matteucci M, Sordi G, Busato E, Zorzi M, Maggino T, Del Mistro A, Cin2 Study Working Group .

Background/Objectives: The detection and treatment of high-grade cervical lesions prevent the development of invasive cervical cancer. Excisional procedures can pose a risk for subsequent pregnancies, thus conservative management of Cervical Intraepithelial Neoplasia grade 2 (CIN2) lesions should be adopted in young women. We are conducting a prospective multicentre study aimed to evaluate feasibility and efficacy of CIN2 conservative management, as well as the role of different biomarkers. Here we analyse women's eligibility and acceptance.

Methods: Data on all women diagnosed with CIN2 between April 2019 and October 2021 in four centres within organised screening in the Veneto region were collected. Women fitting the study's predefined criteria and willing to participate were recruited. Inclusion criteria were: age 25-45 yrs, transformation zone and lesion fully visible at colposcopy. Exclusion criteria were: pregnancy, previous treatment of a CIN2+ lesion, immunodeficiency, and presence of an endocervical lesion not completely visible at colposcopy. Women are grouped as: not eligible; eligible but refused participation; enrolled. Women are followed for 24 months, with colposcopy and testing for several biomarkers. Upon signing an informed consent, women are enrolled in the study and cervical cell samples collected. Treatment is delayed and subsequently performed in the case of lesion progression, or persistence for >12 months.

Results: Overall, 640 women were diagnosed with CIN2; 228 (35,6%) not eligible; 93 (22,6%) eligible but refused; 319 (77,4%) enrolled. The most frequent reason of exclusion was age >45 yrs. Among eligible not enrolled women, refusal/willingness to be treated accounted for 78,5%. CIN2 prevalence, no eligibility, eligibility, refusal and acceptance rates by age (expressed as number and %) are reported in Table 1.

Conclusions: The distribution of CIN2 lesions by age in our area indicates slightly higher figures before 40 years of age. According to the inclusion/exclusion criteria set for our study, we could enrol half of the cases diagnosed. Acceptance was (not significantly) higher among 25-34-yrs-old than among older women. At the completion of the study, the clinical outcome and the results of the biomarkers could envisage more effective criteria.

Table 1

Oral presentation

Dr Sørbye Sveinung
sveinung.sorbye@unn.no - Norway

Sorbye Sveinung
sveinung.sorbye@unn.no - Norway

#3615

7-TYPE HPV MRNA TEST IN TRIAGE OF HPV DNA POSITIVE WOMEN

12 - Triage of HPV positive women

Sorbye S, Falang B, Antonsen M

Background/Objectives: In 2019, Norway implemented HPV DNA testing as a more sensitive method in primary screening for women 34-69 years of age. Reflex cytology was applied as triage of the screen positives in line with common international practice. Based on the known limitations and the effect of subjectivity for cytology interpretation, cytology has shown limited specificity in triage. This challenges the advantages of HPV primary screening by lowering the sensitivity. The search for improved triage alternatives is among the key priorities for the HPV expert community to improve diagnostic accuracy, allocate resources and reduce potential harm from over-treatment. We studied the performance of a 7-type HPV mRNA test in triage of HPV DNA positive women compared to cytology as the established reference method.

Methods: Since 2019 to date, the Department of Clinical Pathology, University Hospital of North Norway, has been using the HPV DNA test Roche Cobas 4800 in primary screening of women 34-69 years. All HPV DNA positives were evaluated by cervical cytology and a 7-type HPV mRNA test (PreTect HPV-Proofer⁷, identifying mRNA E6/E7 from the genotypes 16, 18, 31, 33, 45, 52 and 58). According to national guidelines, follow-up was based on cytology. Histologically confirmed CIN2+ was used as the study endpoint.

Results: A total of 16,729 women were screened by HPV DNA testing. The positivity rate was 5.0% (836/16,729). During follow-up through end of November 2021, a biopsy has been taken from 31.1% (260/836) where the detection rate of CIN2+ among the HPV DNA positives was 8.7% (73/836). In triage of the HPV DNA positive women, 55.0% (460/836) had abnormal cytology (ASC-US+) and 36.5% (305/836) had positive HPV mRNA test. The sensitivity, specificity, PPV and NPV for the detection of CIN2+ were 76.7%, 47.2%*, 12.2%** and 95.5% by cytology and 80.8%, 67.8%*, 19.3%** and 97.4% for the HPV mRNA test respectively (*p<0.01, **p<0.01).

Conclusions: The 7-type HPV mRNA test had higher sensitivity, statistically significant higher specificity, PPV, and lower positivity rate than cytology in triage of HPV DNA positive women. A low positivity rate translates into a low referral rate for colposcopy, which is very appealing for triage situations to minimize the burden on the health care system. Women with a negative triage test can be followed up with repeat HPV DNA testing after 1-2 years. Molecular triage by mRNA is applicable for self-collected samples, an approach with clear benefits made even more important during COVID-19 and its impact on health care and cancer services to increase screening participation and prevention of cervical cancer.

Oral presentation

Ms Korsgaard Andreassen Emilie
emilie.korsgaard.andreasen@regionh.dk - Denmark

Korsgaard Andreassen Emilie
emilie.korsgaard.andreasen@regionh.dk - Denmark

#3616

ANALYTICAL STABILITY OF ROVERS EVALYN AND COPAN FLOQSWAB HPV SELF-SAMPLING DEVICES; IMPLICATIONS FOR USE

13 - Self-sampling

Background/Objectives: HPV self-sampling is emerging as an alternative sampling modality to facilitate increased participation in cervical cancer screening. Typically, self-sampling kits are mailed to women who subsequently return them for analysis. In this process, the self-sample will be transported under diverse temperature conditions and time-to-analysis depends on the postal infrastructure. Here, we assess the analytical stability of the Rovers Evalyn and COPAN FLOQSwab self-sampling brushes under low, normal, and extreme temperatures and during prolonged storage after sampling.

Methods: The analytical quality was assessed using the human beta-globin (HBB) and HPV16 Ct scores of the BD Onclarity HPV test. Two endpoints were used: rate of analytically invalid self-sampling brushes upon testing and changes in Ct scores. The assay cut-off for HBB is Ct 34.2, the cut-off for HPV16 is Ct 38.3 (manufacturers specifications). A pool of 120mL non-fixated, confirmed HPV16 positive cervical samples constituted a master sample. The FLOQSwab brushes were kept at three different temperatures (4°C, room temperature, RT, and 40°C) and four different timepoints (0 days, 4 weeks, 8 weeks, and 16 weeks). The Evalyn brushes were kept at RT for the first timepoint (0 days) and 40°C for the remaining timepoints (4 weeks, 8 weeks, 16 weeks). A total of 300 FLOQSwab and 120 Evalyn brushes were prepared, with 30 brushes per time and temperature point. A similar set of data on 50°C is in preparation and will be reported too. The study is conducted as a quality development study approved by AHH-Hvidovre Hospital.

Results: The proportion of analytical invalid samples increased with time and temperature. For FLOQSwab, the proportion of invalids measured by HBB at RT were 20% at 4 weeks and 70% at 16 weeks. At 40°C, the number of invalid samples was 73.3% at 4 weeks and 100% at both 8 and 16 weeks. For the Evalyn brush, the proportion of invalid samples was 3.3% at RT and 60% at 40°C for timepoints 4 and 16 weeks. Time and temperature affected the analytical quality of samples. At RT, FLOQSwab showed a T0 mean HPV16 Ct of 27.43 (N=30). At RT, T16 weeks, the overall mean HPV16 Ct was 32.12 (N=30) and at 40°C, T16 weeks, the mean Ct-value was 38.13 (N=12). For the Evalyn self-sampling device at RT, the reference value at T0 was a mean HPV16 Ct of 28.93. At T16, 40°C the mean HPV16 Ct was 27.90 (N=29).

Conclusions: Overall, the FLOQSwab was less analytical stable than the Evalyn brush. The implications of these study data are that Evalyn can be safely used with prolonged transport at RT, whereas FLOQSwab has a time limited stability at no longer than 4 weeks. Increased ambient temperature to 40°C affected both brush types negatively, with FLOQSwab being less analytically stable than Evalyn.

Oral presentation

Dr Ikenberg Hans
hans.ikenberg@cytomol.de - Germany

Dr Ikenberg Hans
hans.ikenberg@cytomol.de - Germany

#3617

COMPARISON OF THE GENIUSTM SYSTEM FOR DIGITAL CYTOLOGY WITH THE THINPREP® IMAGING SYSTEM

17 - Automation in cytology

Ikenberg H, Lieder S, Ahr A, Xhaja A

Background/Objectives: Up to now the gold standard in cervical cytology was liquid based cytology (LBC) with computer-assistance (CAS). A further increase in sensitivity and specificity would be welcome especially as experience with abnormalities will decrease with increasing rates of HPV vaccination. A new approach is digital cytology (DC) with artificial intelligence (AI) which also allows assessment by external experts worldwide. We compared the two systems in a retrospective study.

Methods: 1994 ThinPrep® LBC slides (Hologic, Wiesbaden, Germany) pre-analyzed with the Imaging system (CAS) in routine cotesting cytology and HPV in 2020 were reviewed in a blinded fashion with the Genius™ System for digital cytology (Hologic, Wiesbaden, Germany). In 555 cases, among them all >LSIL, a histology result was available. The slides were digitally scanned (volumetric scan) in 14 levels which were integrated to one. AI algorithms finally present a gallery of 6 "tiles" each in 5 categories. 6 additional "tile-rows" may be shown. Therefrom the diagnoses are made. If necessary, the digitalized full slide can be fully screened on the monitor. All cytotechnicians and cytologists had successfully attended a morphology training and had experience with over 10.000 DC cases when beginning the study. All cases with a mismatch between DC and Imaging system results were reviewed by an independent cytologist.

Results: Over 95% of the cases could be technically successful digitalized. The familiarization with the system was rapid, only the micrometer screw was sometimes missed. However, the acceptance of the presented "tiles" by the cytotechnicians was very high. In the retrospective analysis we found in 86,5% of the slides a complete match between both systems. Using the same cytology categories and considering the histology result the match was 90,37%. Applying in addition a cytology follow-up and/or a retrospective review it reached 97,34%. Only in 0,65% a major discrepancy (two grades of cytology or a LSIL / HSIL shift) was observed. All of them in favour of DC. The screening time was significantly shorter with DC.

Conclusions: With the Genius™ system for digital cytology sensitivity and specificity for LSIL and HSIL are at least as high as with LBC and CAS. Screening time is significantly lower. The acceptance by cytotechnicians and cytologists is high.

Oral presentation

Dr Sørbye Sveinung
sveinung.sorbye@unn.no - Norway

Aranda Flores Carlos Eduardo
aranda_floresc@hotmail.com - Mexico

#3618

SELF- VERSUS CLINICIAN-COLLECTED SAMPLES FOR THE DETECTION OF HPV BY 14-type DNA AND 7-TYPE mRNA TESTS

13 - Self-sampling

Aranda Flores C, Sorbye S

Background/Objectives: Cervical cancer remains a major public health problem in Mexico. Self-sampling for HPV testing increases access to screening and population coverage. E6/E7 mRNA detection has been identified as a potential biomarker for triage of DNA-positive women. The aim of this study was to compare self-sampling versus clinician-sampling for complete molecular HPV diagnostics, with respect to sample quality, positivity rates and relevance for referral to colposcopy

Methods: 505 women aged 30-65 years at Mexico General Hospital underwent self-sampling (Mia by XytoTest) and clinician-sampling (Cervex by Rovers). Samples were tested for HR-HPV DNA (RealTime hrHPV, Abbott (14 types)) and E6/E7 mRNA (PreTect HPV-Proofer⁷ (7 hrHPV-types in the 9-valent vaccine)).

Results: HPV-DNA prevalence was 22.8% in self-collected versus 19.2% in clinician-collected samples ($P=0.19$). Overexpression of mRNA E6/E7 from 7 HPV types was 7.1% and 6.3%, respectively ($P=0.71$). Overall agreement between the two collection methods was fair, with a concordance rate of 78.2% (390/505), $k=0.34$ (95% CI: 0.25-0.44), $P<0.001$, for the HPV-DNA test and 92.5% (467/505), $k=0.40$ (95% CI: 0.25-0.56), $P<0.001$, for the mRNA test, respectively. The self-sampled aliquot contained about 3 times more cells compared to clinician taken aliquot; 1.87 million cells/ml versus 0.63 million cells/ml, respectively.

Conclusions: Self-sampling (Mia by XytoTest) is as reliable as clinician-sampling for HPV testing and allows direct HPV mRNA genotyping. Self-sampling is an effective strategy to reach under screened women. The high prevalence of DNA positive results reflects the need for reflex triage, thereby reducing unnecessary colposcopies, women's uncertainties, and cost. A low positivity rate of concurrent DNA+/mRNA+ (6.3%) effectively discriminates women warranted for immediate colposcopy/biopsy from return to follow-up and suggests longer follow-up interval for single DNA+ women. Such a strategy will inevitably reduce over-referral for colposcopy but needs clinical and cost-benefit assessment in prospective studies.

Oral presentation

Ms Li Boning
lbnkyrie@163.com - China

Li Boning
lbnkyrie@163.com - China

#3619

25-hydroxycholesterol inhibits human papillomavirus infection in cervical epithelial cells via disturbance of cytoskeleton remodeling through lipid metabolism suppression

02 - Viral and molecular biology

Li B, Hua C, Lu L, Sui L

Background/Objectives: In spite of the fact that several prophylactic vaccines have been approved for prevention of high-risk human papillomavirus (HPV) infection and the occurrence of cervical cancer, not all high-risk HPV types are included in the vaccines; therefore broad-spectrum therapeutic strategies for HPV infection are crucial. 25-hydroxycholesterol (25HC) is the enzymatic product of cholesterol 25-hydroxylase (CH25H), which is encoded by an interferon-stimulated gene (ISG). It has recently been identified as a cholesterol biosynthesis regulator and virus repressor, targeting entry of enveloped virus, including SARS-CoV-2. However, the mechanisms of 25HC to limit non-enveloped viruses, such as HPV, remain to be explored. Here we investigated the anti-HPV potency of 25HC and illustrated probable mechanisms of this virus infectivity inhibitor.

Methods: We used HPV pseudoviruses (PsV) with reporter genes to identify the capability of 25HC to protect cervical epithelial cells from multi-types of HPV infection. IVIS in vivo imaging system was applied to evaluate the potency of 25HC in HPV infection prevention in a murine model. Immunofluorescence assays were conducted to determine the involvement of actin filaments in 25HC antiviral effect. RNA-seq, Western blot and Rho pull-down assays were used to study cytoskeleton reorganization mechanisms after 25HC treatment.

Results: 25HC exhibited broad-spectrum anti-HPV activity in cervical epithelial cells. Intravaginally applied and intraperitoneally applied 25HC inhibited HPV infection and reduced viral load in the cervical region of murine genital challenge models. 25HC did not block HPV PsV interaction with cervical epithelial cell receptors, but rather regulated cellular milieu. During HPV infection, 25HC disturbed cytoskeletal remodeling in cervical epithelial cells, which is reported to be vital in virion transportation. Further analysis revealed that, 25HC inhibited isoprenoid biosynthetic process and subsequently restricted prenylation of Rho GTPases. The decreased prenylation suppressed Rho GTPases activation, then restrained LIMK/cofilin activation, and ultimately interfered with polymerization status of filopodia and stress fibers.

Conclusions: Here we show a previously unrecognized role of 25HC in influencing cytoskeleton remodeling through its cholesterol regulator capacity. Dysfunction of actin filaments dynamics plays a critical role in the suppression mechanism of 25HC against HPV infection. Our study provided new clues and theoretical basis for therapeutic targets of HPV infection.

Oral presentation

Ms Payrich Eva-maria
eva-maria.payrich@charite.de - Germany

Payrich Eva-maria
eva-maria.payrich@charite.de - Germany

#3620

THE EMERGING ROLE AND OPINION ABOUT IMPLEMENTATION OF HPV SELF-SAMPLING IN GERMANY: A MIXED METHODS STUDY

13 - Self-sampling

Payrich E, Skof A, Struck M, Thies S, Schreckenberger C, Kaufmann A

Background/Objectives: The risk of developing cervical cancer is increased for non-participants in screening programs. In 2019, only 47.7% of all women entitled participated in the opportunistic screening program in Germany. Self-sampling increases participation rates (1) and is as sensitive as HPV tests performed on clinician samples (2). There are few qualitative data on opinions by women and physicians. Our aim was to involve potential users and learn about experiences, expectations, and possible applications of Self-sampling

Methods: This mixed-methods-study was a substudy of the FACTS project, a study to validate whether the use of the QuantiGene-Molecular-Profiling-Histology assay (QG-MPH) can triage women screened positive by cytology or HPV-tests. All included women (n=696) were asked to perform Self-sampling with the Evalyn® Brush in addition to a gynaecological specimen and to fill a questionnaire. Additionally, 25 one-to-one semi-structured interviews with FACTS participants (n=9), women recruited by a social media call (n=5), study gynaecologists (n=5) and gynaecologists not involved in the study (n=6) were performed.

Results: The Self-sampling participation rate was 86.6%. According to 496 questionnaires evaluated 413 (83.3%) women median age 40 (20-77) had participated in the screening several times. The application of Self-sampling and the Self-sampling compared to a gynaecological smear was described as "good or very good" by 89.3% and 82.9%, respectively. Nevertheless, all interviewed women (n=14) stated that they would not prefer Self-sampling to a visit to the gynaecologist, which was a concern by most of the gynaecologists interviewed. Self-sampling was seen more as a way of reassuring oneself, not having to extend screening intervals due to time constraints and an opportunity of female empowerment in gynaecological aspects. However, women and gynaecologists interviewed also indicated concerns about mishandling of the Self-sampling brush, lack of control opportunities and further physician involvement when results are abnormal.

Conclusions: Self-sampling as an additional option to office-based screening is well accepted among German women. However, the Self-sampling option is stated not to replace a visit to the gynaecologist, although this is concerned by many physicians. It would be conceivable to implement Self-sampling in addition to the office-based screening to facilitate access and to meet the desire for self-determination. It must be considered that our study mainly included women from urban areas with a high gynaecological office density and high screening participation.

References: 1. Bosgraaf RP, Verhoef VM, Massuger LF, Siebers AG, Bulten J, de Kuyper-de Ridder GM, et al. Comparative performance of novel self-sampling methods in detecting high-risk human papillomavirus in 30,130 women not attending cervical screening. *Int J Cancer*. 2015;136(3):646-55. Epub 2014/06/14. doi: 10.1002/ijc.29026. PubMed PMID: 24923998. 2. Arbyn M, Smith SB, Temin S, Sultana F, Castle P, Collaboration on S-S, et al. Detecting cervical precancer and reaching underscreened women by using HPV testing on self samples: updated meta-analyses. *BMJ*. 2018;363:k4823. Epub 2018/12/07. doi: 10.1136/bmj.k4823. PubMed PMID: 30518635; PubMed Central PMCID: PMC6278587

Oral presentation

Mrs Suominen Nelli
nelli.t.suominen@utu.fi - Finland

Suominen Nelli
nelli.t.suominen@utu.fi - Finland

#3621

PERINATAL HPV TRANSMISSION BETWEEN PARENTS AND THEIR OFFSPRING

03 - Epidemiology and natural history

Suominen N, Luukkaala T, Rintala M, Grénman S, Syrjänen S, Louvanto K

Background/Objectives: The knowledge on the routes of human papillomavirus (HPV) transmission in early childhood are scanty. Vertical transmission can occur from either mother or father although father-to-child HPV transmission is less studied. This study aimed to investigate the perinatal (i.e. prenatally or at birth) HPV transmission from mothers and/or fathers to their offspring.

Methods: Altogether, 321 mothers, 134 fathers and their 321 offspring from the Finnish Family HPV study cohort were included to the analyses. Parents' genital and oral brush samples in addition to semen samples were collected for HPV testing at baseline (36 week of pregnancy). Oral, genital and umbilical samples from the newborn in addition to placenta samples were collected immediately after delivery. HPV genotyping was performed with Luminex-based Multimetrix kit (Progen Biotechnik GmbH, Heidelberg, Germany) to identify 24 different low-risk (LR) and high-risk (HR) HPV genotypes. Type-specific HPV risk for the child was calculated for each HPV genotype separately from the mother or the father using univariable logistic regression analysis. Univariable- and multivariable-adjusted multinomial logistic regression analyses were used to determine associations between mother's, father's and child's site and LR- and HR-HPV specific status.

Results: Concordance between mother's and her newborn HPV genotype at any site was statistically significant for HPV genotypes 6, 16, 18, 31 and 56, OR ranged from 3.41 (95% CI 1.80-6.48) with HPV16 to 634 (95% CI 28.5-14087) with HPV31. HPV6 and HPV31 also showed statistically significant father-newborn HPV genotype concordance, OR 4.89 (95% CI 1.09-21.9) and OR 65.0 (95% CI 2.92-1448), respectively. Oral HR-HPV infections of the both parents increased the risk of newborn's oral HR-HPV presence, adjusted OR range from 4.87 to 7.72 (95% CI range from 2.10 to 28.3). Newborn's oral LR-HPV infection was predicted only by the mother's oral LR-HPV and the father's urethral LR-HPV infection (adjusted OR range from 7.12 to 7.93 (95% CI range from 1.12 to 56.1)). Mother's genital HR-HPV positivity increased the risk of newborn's genital HR-HPV by 3.3-fold (OR of 3.34, 95% CI 1.35-8.30).

Conclusions: The genotype specific HPV concordance between parents and their infant indicates that vertical HPV transmission occurs from both parents to their offspring already during the perinatal period of life.

Oral presentation

Dr Man Irene
mani@iarc.fr - France

Man Irene
mani@iarc.fr - France

#3623

Evidence-based impact projections of single-dose human papillomavirus (HPV) vaccination in India

40 - Public health

Man I, Georges D, De Carvalho T, Saraswathi L, Bhandari P, Kataria I, Siddiqui M, Muwonge R, Berkhof J, Sankaranarayanan R, Bogaards J, Basu P, Baussano I

Background/Objectives: Despite having the highest expected burden of cervical cancer worldwide, India still has limited access to HPV vaccination and cervical screening. Immunization by single-dose schedule could facilitate the scale-up of HPV vaccination in India and other low- and middle-income countries. Currently, evidence on the effectiveness of the single-dose schedule is expanding. In this work, we explore the impact of single-dose HPV vaccination in India and compare it to the impact of two-dose vaccination.

Methods: Based on Indian demographic and epidemiological data, and on data from the IARC India vaccine trial with 10-years follow-up, we constructed an HPV transmission model (EpiMetHeos) and adapted it to Indian states. Using the model, we projected the impact of HPV vaccination on the occurrence of HPV infection and cervical cancer incidence. Projections were obtained under different vaccination coverages, catch-up age ranges, as well as different scenarios of waning vaccine protection by single-dose vaccination.

Results: Under the base-case scenario, i.e., equal life-long vaccine protection by single-dose as compared to two-dose vaccination and 90% coverage, both single- and two-dose vaccination were projected to eliminate HPV 16 and 18 in the long-term and to provide some cross-protection to HPV 31, 33 and 45. Life-time risk (LTR) of cervical cancer was reduced by 70% in the first 5 routinely vaccinated cohorts and by 79% in the long term. The elimination threshold for age-standardized cervical cancer incidence rate (of 4 cases per 100,000 women) was reached throughout India, while an alternative threshold by standardized LTR (of 250 cases per 100,000 women born) was reached only in some Indian states with relatively low pre-vaccination LTR. Under scenarios with waning single-dose protection, vaccination could still reduce the LTR of cervical cancer by at least 50% if efficacy sustained above 75% for at least 20 years and vaccination coverage remained above 60%. Finally, under all considered scenarios with waning single-dose protection, vaccinating 10 routine and 10 catch-up cohorts with single-dose schedule had a higher impact on cancer cases than using the same number of doses to vaccinate only 10 routine cohorts with two doses.

Conclusions: Capitalizing on data on Indian cervical cancer epidemiology and from the IARC India vaccine trial, we demonstrated the potential of single-dose HPV vaccination and its comparative advantage to two-dose vaccination in India. Our context-specific projections could provide guidance to policymakers in deciding whether and how to implement cervical cancer preventive measures.

Oral presentation

Dr Siegler Efraim
siegler@netvision.net.il - Israel

Siegler Efraim
siegler@netvision.net.il - Israel

#3626

Clearance of High-Risk HPV after Large Loop Excision of Transformation Zone of Early Stage Cervical Cancer and Adenocarcinoma in Situ has a high correlation with absence of residual disease

26 - Cervical neoplasia

Siegler E, Goldberg Y, Siegler Y, Mazarebe S, Kugelman N, Sabu E, Lavie O, Shaked-mishan P, Mackuli L, Segev Y

Background/Objectives: Better predictors of recurrence are needed after fertility preserving surgery in women with early-stage cervical cancer and adenocarcinoma in situ (AIS). We assessed a possible correlation of High-Risk human papillomavirus (HR-HPV) clearance after large loop excision of the transformation zone (LLETZ) in women diagnosed with early-stage cervical cancer and AIS with no residual cancer.

Methods: Data were collected from 102 women diagnosed with early-stage invasive cervical cancer and AIS, who HR-HPV positive and had a repeat cervical HPV test short time after the operation, 3-12 weeks post LLETZ, and before the final surgical treatment. We compared characteristics of women with negative and positive HR-HPV post LLETZ.

Results: At the post-LLETZ follow up visit, 45 women were negative and 57 were HR-HPV positive for HR=HPV. The pathological results of women who were HR-HPV negative compared to HR-HPV positive after LLETZ included a significantly higher incidence of adenocarcinoma and AIS: 21 (46.7% vs 8(14%), ($p=0.0005$): higher proportion with early stage cervical cancer. In the negative Hr-HPV after LLETZ 40(88%) women had normal histology and only 2(4.4%) had cancer in the final histological specimen. Among women who underwent radical hysterectomy/trachelectomy after LLETZ, a normal final histology was observed in 75% (9/12) of HR-HPV negative and 8.7% (2/23) of HR-HPV positive ($p<0.0001$). The negative predicted value of negative Hr-HPV after LLETZ was 95.6%.

Conclusions: Clearance of HR-HPV from the cervix a short period after LLETZ has a high correlation with absence of residual cancer in the outcome, either in the final pathology or the follow up. Testing for HR-HPV a short period after LLETZ might serve as a new parameter for risk assessment: a negative HR-HPV result may suggest the need for a less radical operation in women with early-stage cervical cancer.

Oral presentation

Ms Doshi Mishelben
mishelben.doshi@affinity-cmc.com - United States

Morais Edith
edith.morais@msd.com - France

#3628

UNDERSTANDING HPV ORAL BURDEN AND 9-VALENT HPV VACCINE EFFICACY: DESIGN OF THE PROGRESS, BROADEN AND V503-049 STUDIES

29 - Oral HPV infection

Morais E, Tu Y

Background/Objectives: The burden of human papillomavirus (HPV)-related head and neck cancers (HNC) is high, and routine screening for early detection of these cancers is not available. As such, HPV vaccination represents an attractive approach to prevent HNC. Oropharyngeal cancer, the most studied HNC relative to HPV infection, disproportionately affects men. The proportion of other HNCs attributable to HPV remains unclear. Although countries have increasingly implemented gender-neutral HPV vaccination, globally male vaccination coverage remains low. To inform public health decisions and routine medical practice regarding vaccination of both genders, a better understanding of oral HPV natural history and the burden of HPV-related HNC is needed.

Methods: Three studies have recently been initiated to fill knowledge gaps related to the (1) overall burden of HPV-related HNC, (2) the natural history of oral HPV infection and associated risk factors, and (3) efficacy of HPV vaccination in preventing persistent oral HPV infection, a surrogate endpoint for HPV-related HNC. PROGRESS (The Prevalence of Oral HPV Infection, a Global Assessment) and BROADEN (Absolute Burden of HPV-Related Head and Neck Cancers) are two observational studies designed to assess oral HPV prevalence and natural history (i.e., incidence, persistence and clearance) and to estimate the fraction of oropharyngeal and non-oropharyngeal HNC cases attributable to HPV, respectively. These studies have been initiated in several countries using a common methodology. In addition, an international, double-blind, placebo-controlled trial (Protocol V503-049; NCT04199689) was initiated to evaluate efficacy of the 9-valent HPV vaccine against oral HPV persistent infection in adult men. While efficacy of HPV vaccines against anogenital persistent infection and disease endpoints has been well established, V503-049 is the first large randomized clinical trial to evaluate efficacy against persistent oral HPV infection.

Results: N/A

Conclusions: Data from BROADEN and PROGRESS will fill knowledge gaps concerning the natural history of oral HPV infection and burden of HPV-related HNC. Together with direct evidence of vaccine efficacy against persistent oral HPV infection from the randomized clinical trial, these data will aid physicians, patients and healthcare decision makers in understanding the potential impact of vaccination on reducing the overall burden of HNC.

Poster

Dr Steinkasserer Lena
steinkasserer.lena@mh-hannover.de - Germany

Dr Steinkasserer Lena
steinkasserer.lena@mh-hannover.de - Germany

#3629

CHARACTERIZATION OF PATIENTS WITH VULVAR LICHEN SCLEROSUS ET ATROPHICUS AND ASSOCIATION TO VULVAR CARCINOMA: A RETROSPECTIVE SINGLE CENTER ANALYSIS

27 - Vulvar diseases and neoplasia

Steinkasserer L, Hillemanns P, Jentschke M

Background/Objectives: Lichen sclerosis (LS) is a benign, cutaneous, chronic inflammatory (autoimmunological) disease. The HPV independent or differentiated vulvar intraepithelial neoplasia (dVIN) accounts as a precursor lesion of vulvar squamous cell carcinoma and is often associated with lichen sclerosis. Although association between lichen sclerosis and vulvar carcinoma has long been recognized, there is a lack of evidence in literature.

Methods: This retrospective study examined pseudonymized data of 499 women diagnosed with vulvar pathology between 2008 and 2020 at the Department of Obstetrics and Gynecology of Hannover Medical School (MHH). Data were further stratified with respect to time of onset, location of disease, accompanying disease, HPV status and progression of disease into vulvar squamous cell carcinoma (VSCC).

Results: In total, 56 patients were diagnosed with vulvar lichen sclerosis et atrophicus. The median of onset of disease was 60.3 years of age (78.6% postmenopausal). The incidence of vulvar cancer in women diagnosed with lichen sclerosis was 48.2%, namely 27 women with lichen sclerosis developed a VSCC during patient's treatment. 25 patients reported a diagnosis of VIN in their self-reported history (44.6%). In 17 patients who developed vulvar carcinoma in course, lichen sclerosis and VIN were confirmed histologically (62.9%). 33.4% of VSCC were HPV negative, in 63% HPV status was not done.

Conclusions: Lichen sclerosis is an underestimated disease. The association between lichen sclerosis and vulvar cancer is well known, but further studies are missing to close the lack of data. In our retrospective study we showed a strong correlation between vulvar lichen sclerosis et atrophicus and VSCC. VIN and postmenopausal status at time of lichen sclerosis diagnosis tent to be important predisposing factors for development of a VSCC. Our results highlight the importance to diagnose Lichen sclerosis et atrophicus early to ensure adequate follow-up and prevent progression to VSCC.

Poster

Dr Tachezy Ruth
tachezr@natur.cuni.cz - Czech Republic

Ruth Tachezy
tachezr@natur.cuni.cz - Czech Republic

#3630

LEVEL OF CYTOTOXIC T CELLS POPULATIONS IN HEAD AND NECK SQUAMOUS CELL CARCINOMA MICROENVIRONMENT AS A PREDICTOR OF PATIENT PROGNOSIS

30 - HPV and oropharynx / Head and neck cancer

Pokryvkova B, Grega M, Klozar J, Vencalek O, Nunvar J, Ruth T

Background/Objectives: Head and neck squamous cell carcinomas (HNSCC) belong to a group of diverse tumors, which can be induced by infection of human papillomavirus (HPV) or by tobacco and / or alcohol consumption. The viral etiology of HNSCC relates to the better clinical outcome reflecting the different immune system response. Recently, spatial analysis of the tumor microenvironment has enabled in situ analysis of infiltrating immune cells.

Methods: Here, we retrospectively analyzed 97 tissue samples of oral and oropharyngeal carcinomas with the known etiology using multispectral fluorescent immunohistochemistry based on OpalTM chemistry. To evaluate the immune cell infiltration in tumor and stroma compartments we designed 4 panels of 5 - 6 antibodies each. We mainly focused on quantification of CD4+, CD8+, and FOXP3+ T lymphocytes as well as on their subpopulations expressing PD1, CTLA4, or ICOS molecules. The cell counts were compared according to the tumor etiology and univariate and multivariate survival analyses were performed.

Results: More CD4 + and CD8 + T cells were detected in both compartments of HPV + tumors, which may be explained by the presence of viral antigens. Higher prevalence of PD1 + T cells both in the parenchyma and the stroma of HPV+ tumors points to T cell activation rather than exhaustion status. The higher prevalence of ICOS+ Tregs and CTLA4 +CD4 + T cells in the HPV - group agrees with poor prognosis of these patients and may also explain their poor immune response. Additionally, we observed significantly higher VEGF production in both compartments of HPV- tumors. To evaluate the influence of TIL infiltration on patients' prognosis we used Cox models. We fitted 38 models (19 for the parenchyma and 19 for the stroma) for OS and DSS analyses. We also tested models for the whole tumour and models where HPV status was not included. The detailed results will be presented.

Conclusions: We confirmed the HPV status as a main predictor of patients' prognosis but the number of PD1+CD8+ T cells, and the number of CD8+ T cells, all T cells and CD8+/FOXP3+ ratio were independent factors influencing the overall and / or disease specific survival.

References: This research was funded by the Czech Health Research Council (project 17-28055A), by the European Regional Development Fund and MEYS, project CZ.02.1.01/0.0/0.0/16_019/0000785.

Oral presentation

Dr Dostalek Lukas
lukas.dostalek@vfn.cz - Czech Republic

Dr Dostalek Lukas
lukas.dostalek@vfn.cz - Czech Republic

#3631

The impact of lymph node micrometastases on disease free survival in cervical cancer patients - a retrospective subgroup study of the SCCAN (Surveillance in Cervical CANcer) project

26 - Cervical neoplasia

Cibula D, Borcinova M, Benesova K, Klat J, Falconer H, Kim S, Van Lonkhuijzen L, Lopez A, Isla Ortiz D, Landoni F, Kostun J, Ricardo D, Odetto D, Zapardiel I, Jarkovsky J, Javurkova V, Salehi S, Abu-rustum N

Background/Objectives: The impact of lymph node (LN) micrometastases (MIC) in cervical cancer patients remains a controversial topic given their low incidence and good prognosis of patients managed by primary surgery. We aim to evaluate the prognostic significance of MIC and isolated tumour cells (ITC) in a large cohort of patients from the SCCAN retrospective study (Surveillance in Cervical CANcer). SCCAN study analysed data from more than 4300 patients with early stage cervical cancer treated by primary surgery at 20 large tertiary institutions from Europe, North America, South America and Australia.

Methods: In this SCCAN sub-study, we included patients with early stage cervical cancer (T1a1 LVSI+ - T2b) treated between 2007 and 2016 with at least 1-year follow-up data availability, who were treated by primary surgery including sentinel lymph node (SLN) biopsy and in whom SLNs were processed by pathological ultrastaging protocol.

Results: Out of 969 included patients with at least 1 SLN detected, 174 (18%) had positive LN (Table 1). Maximal tumour diameter >2cm, positive LVSI, grade ≥ 2 , uncommon histological type (neuroendocrine, sarcoma, etc.) and macrometastasis (MAC) or MIC in LN were factors associated with significantly decreased five-years disease free survival (DFS) (Table 2). MAC, MIC or ITC was the largest LN metastasis in 84 (9%), 59 (6%) and 31 (3%) cases respectively. Adjuvant (chemo)radiation was administered in 89%, 85% and 58% of patients with MAC, MIC and ITC. DFS reached 75%, 73% and 83% in patients with MAC, MIC and ITC compared with 90% in the N0 patients. Patients with MAC and MIC had significantly decreased DFS than those with N0 disease (HR=2.36 and 2.55).

Conclusions: Early-stage cervical cancer patients with MIC in pelvic LN have significantly decreased DFS. Their management should follow the same principles as in patients with MAC.

Lukas Dostalek

Poster

Dr Picconi Maria Alejandra
mapicconi@gmail.com - Argentina

Dr Correa Rita Mariel
correamariel01@gmail.com - Argentina

#3632

DISTRIBUTION OF HUMAN PAPILLOMA VIRUS (HPV) GENOTYPES IN CERVICAL SAMPLES OF LATIN AMERICAN WOMEN, IN THE FRAMEWORK OF THE ESTAMPA STUDY.

14 - Genotyping

Baena A, Valls J, Colucci M, Mendoza L, Rol M, Wiesner C, Ferrera A, Fellner M, Gonzalez J, Basiletti J, Mongelos P, Rodriguez De La Peña M, Saino A, Venegas G, Calderon A, Rodriguez G, Barrios H, Herrero R, Almonte M, Picconi M, On Behalf Of T

Background/Objectives: Persistent infection with high-risk human papillomavirus (HR-HPV) is the leading cause of cervical cancer (CC). While the proportion of CC associated with HPV16 and 18 is greater than 70% and similar throughout the world, the relative importance of the other types of HR-HPV differs according to geographic region. The aim was to analyze the genotype-specific distribution of HPV in cervical samples with different histological diagnoses, within the framework of ESTAMPA trial ("EStudio multicéntrico de TAMizaje y triage de cáncer cervicouterino con pruebas del PApillomavirus") ongoing in nine Latin American countries.

Methods: A cross sectional study was carried out in a sub-cohort of 1,114 women, being HPV positive by screening tests (HC2, Qiagen and Cobas 4800 assay, Roche), at recruitment. Four groups of clinical categories by histological diagnosis were analyzed: 1) cervical intraepithelial neoplasia grade 1 or minus (\leq CIN1) (n=728); 2) CIN2 (n=115); 3) CIN3 (n=188), and 4) CC (n=83). Samples were selected by convenience in order to achieve a similar number of samples in each of the diagnostic categories. Type-specific genotyping was carried out by two PCR strategies combined with reverse hybridization: a) BSGP5 +/GP6+ bio and b) PGMY09/11 (CHUV), which can identify 37 low risk (LR) and HR-genotypes. Each genotype was counted independently.

Results: The most common genotype in all clinical categories was HPV16, varying between \leq CIN1 (17%) and CIN2 (20.9%), but increasing in CIN3 (53.2%) and CC (65.1%) ($p < 0.001$). A great diversity of LR and HR-HPV was observed in \leq CIN1 and CIN2; while in CIN3 and CC, the viral spectrum was more restricted, with a clear predominance of HR-HPV (98.4% and 97.6%, respectively) ($p < 0.001$); among them stand out, in addition to HPV16, HPV31 (13.3%), HPV52 (11.7%) and HPV58 (9.0%) in CIN3 and HPV45 (8.4%), HPV18 (7.2%) and HPV52/58 (4.8%) in CC, although without statistical significance.

Conclusions: Our results confirm the relevance of HPV16 in the whole clinical spectrum of cervical lesions, with a strong rise of its proportion in CIN3+; particularly, in CC, the top ranked HPV types are the seven HR-HPVs which are components of the nonavalent vaccine (HPV16, 18, 31, 33, 45, 52, and 58), being dominant HPV16, HPV18, HPV45. These baseline data on the distribution of HPV genotypes in women from Latin America in the context of screening will be valuable to predict how vaccination and screening based on the HPV test will specifically influence the prevention of CC in our region (Grants IARC-PRI; PICT 0364-2016 and INC, Argentina)

Oral presentation

Ms Doshi Mishelben
mishelben.doshi@affinity-cmc.com - United States

Luxembourg Alain
alain_luxembourg@merck.com - United States

#3633

LONG-TERM IMMUNOGENICITY AND EFFECTIVENESS OF THE 9-VALENT HPV (9vHPV) VACCINE IN PREADOLESCENTS AND ADOLESCENTS AFTER 10 YEARS OF FOLLOW-UP

06 - HPV prophylactic vaccines

Olsson S, Luxembourg A

Background/Objectives: The 9-valent human papillomavirus (9vHPV) vaccine was developed to protect against infection and disease related to HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58. The pivotal 36-month Phase III immunogenicity study of 9vHPV vaccine in boys and girls aged 9-15 years was extended to assess long-term immunogenicity and effectiveness through 10 years after the last 9vHPV vaccine dose (NCT00943722).

Methods: Participants aged 9-15 years who received three 9vHPV vaccine doses in the base study (Day 1, Months 2 and 6) were enrolled in the study extension (N=1272 [males, n=301; females, n=971]). Serum was collected at Day 1 and Months 7, 12, 24, 36, 66, 90, and 126 to assess antibody responses (by competitive Luminex immunoassay [cLIA; primary immunoassay] and IgG Luminex immunoassay [IgG-LIA]). For effectiveness analysis, genital swabs were collected (to assess HPV DNA by polymerase chain reaction PCR) and external genital examinations (to detect external genital lesions) were conducted every 6 months starting when participants reached age 16 years. Cervical cytology tests were conducted annually for females ≥ 21 years of age; participants with cytological abnormalities were triaged to colposcopy based on a protocol-specified algorithm. External genital and cervical biopsies on abnormal lesions were performed. Tissue samples were adjudicated by a pathology panel. Primary analyses were conducted in per-protocol populations.

Results: Geometric mean antibody titers peaked around Month 7, decreased most sharply between Months 7 and 12, then decreased gradually thereafter through Month 126. Seropositivity rates remained $\geq 81\%$ by cLIA and $\geq 95\%$ by IgG-LIA at Month 126 for each of the 9vHPV vaccine types. After up to 11.0 years (median 10.0 years) follow-up post-Dose 3, there were no cases of HPV6/11/16/18/31/33/45/52/58-related intraepithelial neoplasia or genital warts among male participants. Among females, there were no cases of high-grade HPV6/11/16/18/31/33/45/52/58-related intraepithelial neoplasia or genital warts and one case of HPV-16-related cervical intraepithelial neoplasia grade 1. Incidence rates of HPV6/11/16/18/31/33/45/52/58-related 6-month persistent infection in males and females in the per-protocol population were low (54.6 and 52.4 per 10,000 person-years, respectively) and within ranges expected in vaccinated cohorts, based on results from previous efficacy trials of the quadrivalent and 9vHPV vaccines.

Conclusions: The 9vHPV vaccine demonstrated sustained immunogenicity and effectiveness through approximately 10 years post-9vHPV vaccination of boys and girls aged 9-15 years.

Poster

Dr Picconi Maria Alejandra
mapicconi@gmail.com - Argentina

Dr Fellner María Dolores
fellnermd@gmail.com - Argentina

#3635

EVALUATION OF THE CLINICAL DETECTION OF THE HUMAN PAPILLOMA VIRUS (HPV) USING A VALIDATED PLATFORM: FIRST INTERLABORATORY STUDY IN ARGENTINA.

09 - HPV testing

Basiletti J, Zubillaga E, Rodriguez M, Sciara M, Padin V, Yene C, Picconi M, On Behalf Of T

Background/Objectives: The detection of high risk HPV (HR-HPV) DNA as a primary screening test is being gradually implemented by the Ministry of Health from Argentina, as part of a comprehensive strategy for the prevention of cervical cancer. There are different diagnostic platforms for the detection of HR- HPV validated for clinical use. In order to evaluate the performance of the laboratories and their competence, it is necessary to participate in external quality assessment programs. Since the participation in international programs is expensive and the panels are complex to import, a local program is of great interest for Argentina, with the possibility to be extended to other countries of the region. Our aim was to organize a first interlaboratory study to analyze the competence of laboratories that use a clinical validated platform.

Methods: A panel consisting of 6 vials containing cervicovaginal cells in Preservcyt medium with different concentrations of HPV16, HPV18 and other HR-HPV, was developed; it included samples with single genotype or several (coinfections). The panel was tested 10 times to analyze its homogeneity; the acceptance criterion, considering 1 logs = 3.3 Cp (crossing point), was: trimmed mean of $Cp \pm 0.25$ logs in 90% of the repetitions. Panels were sent to 13 laboratories that routinely use the HPV COBAS 4800 test (ROCHE). The Kappa (K) index was estimated to analyze the concordance in genotypes identification. To analyze the precision at different levels of detection, the mean of the reported Cps for each detected genotype in each of the vials by all the participants, was calculated; the acceptance criterion was trimmed mean of $Cp \pm 0.5$ logs.

Results: The 6 batches of samples showed Cps within the acceptance criterion of reproducibility. There was agreement in 77/78 results of the reported HR-HPV genotypes ($K = 0.98$). The 96.6% of the reported Cp were in an acceptable range and random deviations were detected in 14/117 (12.0%), without a trend for any of the participants.

Conclusions: The call was successful; all participating laboratories submitted their results in a timely manner and proved to be competent for the detection of HR-HPV, showing similar levels of analytical sensitivity. This first study constitutes the starting point for the establishment of an external quality assurance program, which will allow the continuous monitoring of the HPV tests performance in Argentina.

Oral presentation

Dr Berza Natalija
natalija.berza@gmail.com - Latvia

Berza Natalija
natalija.berza@rsu.lv - Latvia

#3637

Acceptability of HPV self-sampling test in a population of Latvian women (preliminary data)

13 - Self-sampling

Berza N, Zodzika J, Kivite-urtane A, Pärna K, Uuusküla A, Pole I

Background/Objectives: Participation in cervical cancer screening is the cornerstone of cervical cancer detection. Latvia has a low attendance rate for cervical cancer screening and a relatively high incidence of cervical cancer. HPV self-sampling is an alternative strategy to improve cervical cancer screening program compliance. The study aims to evaluate the acceptability of self-sampling for human papillomavirus (HPV) testing in the population of Latvian women.

Methods: From February till October 2021 women who attended the colposcopy unit in Riga Eastern Clinical University Hospital for known cervical pathology were invited to participate in the study. They were asked to perform unsupervised HPV self-sampling using written instructions, and after the procedure was asked to fill in the questionnaire regarding the course and admissibility of the procedure. The study was approved by the Ethics Committee of Riga Stradiņš University.

Results: A total of 262 women were invited to participate in the study. 237 (90.5 %) participants agreed to provide a self-sample; of them, 93 (35.5 %) women found self-sampling acceptable and equivalent to the clinician-administered test. 134 (51.1 %) of patients reported that they would like to have a test carried out by a medical professional next time. Nineteen females (7.3 %) graded their feelings during the self-sampling test as "embarrassed during the procedure" or "strongly embarrassed during the procedure". 37 (14.1%) participants recorded mild or severe discomfort during the procedure. On the other hand, 36 (13.7 %) patients felt strong self-confidence during self-sampling, and 20 (7.6%) participants were interested or intensely interested in taking the test.

Conclusions: Self-sampling for HPV testing was acceptable in many patients, and women could carry out the test alone, using simple written instructions. The main arguments against performing self-sampling were shame regarding this procedure and discomfort during the procedure. This issue will need to be addressed if self-sampling is introduced.

Oral presentation

Dr Gottschlich Anna
anna.gottschlich@cw.bc.ca - Canada

Gottschlich Anna
anna.gottschlich@cw.bc.ca - Canada

#3640

PRELIMINARY RESULTS FROM A PRAGMATIC SEQUENTIAL RANDOMIZED TRIAL ON INTEGRATED CERVICAL CANCER SCREENING IN MAYUGE DISTRICT UGANDA (ASPIRE MAYUGE)

10 - HPV screening

Gottschlich A, Payne B, Trawin J, Albert A, Mitchell-foster S, Rawat A, Mithani N, Namogusa R, Pedersen H, Orem J, Smith L, Nakisige C, Ogilvie G.

Background/Objectives: Approximately 87% of cervical cancer diagnoses and 90% of cervical cancer deaths occur in developing regions, particularly in Sub-Saharan Africa, where there is often an absence or poor uptake of comprehensive screening programs. In Uganda, barriers to implementation of effective screening programs include poor health infrastructure and lack of human resources. Self-collection testing for the human papillomavirus (HPV), the primary cause of cervical cancer, may alleviate these barriers and has been shown to be feasible and acceptable among diverse populations. However, knowledge gaps persist in how to best implement self-collection HPV testing within community-based health services.

Methods: The aim of the Advances in Screening and Prevention in Reproductive Cancers (ASPIRE Mayuge) pragmatic sequential randomized trial was to compare the effectiveness of two cervical cancer screening models for self-collected HPV testing: Arm 1) community health worker recruitment (door-to-door) and Arm 2) invitation to community health meetings. Participants ages 25-49 from 31 villages in the Mayuge district of Eastern Uganda were enrolled from 2019-2021 and randomized to Arm 1 (Screening at home, N=1055) or Arm 2 (Screening at the health meeting, N=964) by clusters (villages). Participants completed surveys and were offered self-collection HPV testing. HPV results were returned to participants and those who were HPV positive were referred to a local health clinic for visual inspection with acetic acid (VIA) and thermal ablation. The primary outcome was attendance for follow-up VIA at a designated health clinic when indicated among all participants based on intention to treat. This outcome was compared between arms by estimating a mixed effect logistic regression model with cluster as a random intercept and adjusted for all known confounders at the individual and cluster level.

Results: In Arm 1 and Arm 2, 28.1% (N=296) and 24.9% (240) were HPV positive, and 22.0% (N=232) and 16.6% (N=160) of participants attended VIA, respectively. After adjustment for age, education, and number of living children, the model gave results suggesting that Arm 2 participants were less likely than Arm 1 to attend VIA (OR = 0.71, 95% CI: 0.56-0.90, p = 0.004).

Conclusions: In this population, participants who were recruited by community health workers in their homes were more likely to attend follow-up VIA. It is important to note that Arm 1 occurred before the COVID-19 pandemic, while Arm 2 occurred during the pandemic. Effects of the pandemic could confound the relationship between arm and attendance, but we unfortunately have no way of assessing the level of potential confounding. This study provides evidence that door-to-door recruitment by community health workers may be an effective method for implementing self-collection HPV testing in settings where cervical cancer screening is strongly needed.

Poster

Mrs Musialowicz Victoria
victoria.musialowicz@gmail.com - Poland

Dr Przybylski Marcin
nicramp@poczta.onet.pl - Poland

#3646

A diagnose of advanced cervical cancer during longterm infertility treatment - removal of cervical tumor above 4 cm

41 - Fertility and HPV

Przybylski M, Musialowicz V, Millert-kalinska S, Wszolek K, Jach R, Pruski D

Background/Objectives: In 1974 profesor Harald zur Hausen had proved, that persistent infection with high-risk human papillomavirus is the direct cause of cervical intraepithelial neoplasia and invasive cervical cancer. In 2006 the first prophylactic vaccine against human papillomavirus was registered. Although, despite the knowledge of this pathomechanism, invasive cervical cancer still claim the lives of many women worldwide. Every year we loose about 300,000 patients.

Methods: A 32-years-old woman presented for a routine visit to the gynecologist. The physical examination suggested cervical cancer. Pap smear come back with High-Grade Squamous Intraepithelial Lesions result. The patient was qualified for a colposcopic examination - transformation zone type 3, Reid Colposcopy Index with result 8 points. A punch biopsy and endocervical curettage was performed. The result of the histopathological examination confirmed the diagnosis of cervical cancer. By analyzing the results, the diagnosis of cervical cancer stage IB2 was made.

Results: The patient was qualified for a radically modified hysterectomy with bilateral salpingo-oophorectomy and bilateral pelvic lymphadenectomy. In the postoperative material, a cervival tumor of the size 45x39x20 mm was found. The margins were negative. In one of the removed lymph nodes a metastatic tumor was described, also invasion of lymph vessels was confirmed after the histopathologic examination. The preliminary diagnosis of stage IB2 cervical cancer was confirmed. The patient was qualified for adjuvant chemo-radiotherapy. She underwent five cycles of cisplatin in the dose of 40mg/m² each and radiation - teleradiotherapy (50,4 Gy), radiation boost (9 Gy) and hight dose rate brachytherapy (18 Gy).

Conclusions: The presented case report of a patient treated for many years due to primary infertility may be an example of oversight, which can occur during specialist treatment with the omission of obligatory, preventive examinations. Correctly used knowledge concluded in Jeorjos Papanicolau and prof. Teresa Koprowska publications in the 1920s and the achievements of the 1970s, make it possible to diagnose and to treat pre-cancerous lesions.

Poster

Mrs Musialowicz Victoria
victoria.musialowicz@gmail.com - Poland

Mrs Musialowicz Victoria
victoria.musialowicz@gmail.com - Poland

#3647

Effective treatment of cervical pathology in pregnant woman during COVID-19 pandemic

01 - HPV disease and COVID-19

Przybylski M, Musialowicz V, Millert-kalinska S, Jach R, Pruski D

Background/Objectives: In the beginning of 2020 the whole world was paralysed by SARS-CoV-2 pandemic. The functioning of public health care system has been subordinated to fight against a previously unknown virus. Despite that, other health problems did not disappear.

Methods: In 2020 a patient presented for a routine gynecological visit. After abnormal Pap-smear (ASC-H) and positive HPV 16 test a biopsy was performed. The histopathology examination did not confirm any pathology. The patient qualified for LEEP-conization. Again, the histopathology results were correct. Parallel to the diagnostic process, the patient was vaccinated against human papilloma virus. The subsequent pap smear came in with result LSIL. The molecular test confirmed persistent HPV infection. She referred to the gynecologist to perform another colposcopy, biopsy and endocervical curettage. Histopathology report described adenocarcinoma in situ. In the meantime the patient suffered from COVID-19. After quarantine, the patient was admitted to hospital to undergo surgical conization of cervix. The postoperative histopathological report confirmed, that the tumor was removed radically with clean margins. Oncological supervision was recommended. In December 2020 a schedule visit was planned. The patient referred with a positive pregnancy test. Analyzing the patient's medical history, she was in the risk group of developing cervical insufficiency. During this visit the cervical length was about 30 mm. As part of the prevention of miscarriage and premature birth resulting from earlier treatment of the cervix, progestagen supplementation was joined. Next follow up visits showed no pathology till the 18 week of pregnancy. The cervix shortened to 20 mm. Because of that, the patient was admitted to the hospital. Mycoplasma and Ureaplasma infection was ruled out. During the stay, the cervix did not shorten, so the physicians did not decide for cervical cerclage. The patient was discharged from hospital in good condition, with stable cervix length. During pregnancy, colposcopic examinations were performed in which no abnormalities were found.

Results: In 38 weeks of pregnancy, the patient was referred to hospital for elective caesarean section. The male newborn with a weight of 3200 g was given 10 points in Apgar score.

Conclusions: The presented situation shows, that even during pandemic preventive examinations should not be missed. Early treatment of pre-cancerous lesions is significant to prevent cervical cancer.

Poster

Dr Gopalani Sameer
ruj4@cdc.gov - United States

Saraiya Mona
yys2@cdc.gov - United States

#3649

Incidence trends for HPV-associated anal squamous cell carcinoma in the United States, 1999 to 2018

03 - Epidemiology and natural history

Gopalani S.

Background/Objectives: Human papillomavirus (HPV) vaccines can prevent HPV-associated anal cancer; new promising trial data also suggest a role for screening and treatment for anal cancer among HIV-infected populations. HPV-associated anal cancer cases have increased in the past two decades in the United States (US), with an average annual burden of 7,000 cases.

Methods: Using population-based cancer registries covering approximately 97% of the US population, we calculated age-adjusted incidence rates for HPV-associated anal squamous cell cancers (SCC) using SEER*Stat (version 8.3.9). All cancers were malignant, microscopically confirmed, and restricted to the following ICD-O-3 histology codes for SCC: 8050-8084, 8120-8131. We also examined incidence trends from 1999 to 2018 using average annual percentage change (AAPC). AAPC was calculated using a weighted average of the slope coefficients of the underlying joinpoint regression line with the weights equal to the length of each segment over the interval using the Joinpoint regression program (version 4.9.0.0).

Results: Overall, an average of 5,346 incident cases (rate of 1.66 per 100,000 persons) of HPV-associated anal SCCs were diagnosed over the 20-year period. Incidence rates were higher among females (2.03) than males (1.24), higher among non-Hispanic persons (1.71) than Hispanic persons (1.27), and higher among White persons (1.72) than Black persons (1.60). By age, rates were highest in the older age groups: 50-59 (3.98), 60-69 (5.12), 70-79 (5.22), and >80 (5.12) years. From 1999 to 2018, incidence increased by 2.5% annually, most notably among persons aged 60-69 (AAPC: 4.2) and 50-59 (AAPC: 3.9) years. Incidence also increased among both sexes (female AAPC: 2.9; male AAPC: 1.9), ethnicity groups (non-Hispanics AAPC: 2.7; Hispanic AAPC: 0.8), and among White (AAPC: 2.6), Black (AAPC: 2.3), and American Indian and Alaska Native (AAPC: 1.6) people. Conversely, incidence declined by 1.7% each year among persons aged 30-39 years and remained stable among persons aged 20-29 (AAPC: 1.5; 95% CI: -0.6, 3.7) and 40-49 (AAPC: 0.2; 95% CI: -0.4, 0.7) years.

Conclusions: HPV-associated anal SCC incidence continues to increase overall and among both sexes, ethnicity groups, and most races and age groups; however, incidence declined among persons aged 30-39 years. Ongoing surveillance using population-based registries is needed to monitor trends in the US. Improving HPV vaccine coverage and increased anal cancer screening and treatment among HIV-positive populations could prevent many HPV-associated anal cancers in the future.

Oral presentation

Dr Gottschlich Anna
anna.gottschlich@cw.bc.ca - Canada

Van Niekerk Dirk
dvanniek@bccancer.bc.ca - Canada

#3650

HPV-BASED CERVIX SCREENING AMONG WOMEN LESS THAN 30 YEARS OF AGE IN THE HPV FOR CERVICAL CANCER (FOCAL) TRIAL

10 - HPV screening

Gottschlich A, Smith L, Gondara L, Melnikow J, Cook D, Lee M, Stuart G, Martin R, Peacock S, Franco E, Krajden M, Van Niekerk D, Ogilvie G

Background/Objectives: Early studies on HPV-based cervix screening in women less than age 30 suggested a significant increase in colposcopy referrals that could overwhelm healthcare systems and unnecessary treatment for lesions that would otherwise regress. Thus, most guidelines delay HPV screening until age 30. However, recent evidence indicates that an increase in referrals may only occur in initial rounds of HPV screening. We used data from the HPV For Cervical Cancer (FOCAL) randomized controlled trial to investigate outcomes among participants aged 25-29 at baseline after two rounds of HPV screening.

Methods: Participants were randomized to two arms: HPV screening or cytology. Each arm received their respective test at Round 1 (HPV arm: 25-29 N=829, 30+ N=8723, Cytology arm: 25-29 N=834, 30+ N=8623) and co-testing (HPV/cytology) at 48-month Round 2 (HPV arm: 25-29 N=618, 30+ N=7726, Cytology arm: 25-29 N=611, 30+ N=7467). In this analysis, we only considered the result of each arm's respective primary exit test for colposcopy referral. Attendees were stratified by age group (inf30 or sup30) and we calculated percent referred to colposcopy at each round and with cervical intraepithelial neoplasia findings grade 2 (CIN2) throughout the trial and over 8 years of follow-up after trial exit.

Results: Across rounds, attendees in the younger age group had higher rates of colposcopy referral than the older age group in the same arm (Round 1: HPV - 25-29: 18.0%, 95%CI: 15.5, 20.7, 30+: 4.5%, 95%CI: 4.0, 4.9; Cytology - 25-29: 8.0%, 95%CI: 6.4, 10.1, 30+: 2.5%, 95%CI: 2.2, 2.8). In Round 1, referrals in the HPV arm were higher than in the Cytology arm for both age groups, however at Round 2 referrals were similar for corresponding age groups across arms (Round 2: HPV - 25-29: 5.3% 95%CI: 3.8, 7.4, 30+: 1.4%, 95%CI: 1.2, 1.7; Cytology - 25-29: 6.6%, 95%CI: 4.8, 8.8, 30+: 1.7%, 95%CI: 1.4, 2.0) and lower than Round 1. Among those inf30, throughout the trial, more CIN2 was detected in the HPV arm (N=30, 3.6%, 95%CI: 2.5, 5.1) than in the Cytology arm (N=16, 1.9%, 95%CI: 1.1, 3.1). After 8 years of follow-up the cumulative incidence of CIN2 was similar across arms (HPV: 5.1%, 95%CI: 2.3, 5.7; Cytology: 4.0%, 95%CI: 3.5, 6.7).

Conclusions: While colposcopy rates were higher in HPV-arm attendees aged 25-29 at Round 1, by Round 2 rates were similar between HPV and Cytology arms. More CIN2 lesions were detected among those inf30 in the HPV arm, resulting in more treatment of potentially regressive lesions. However, this difference decreased throughout long-term follow-up, suggesting that these precancers were detected earlier by HPV screening compared to cytology.

Oral presentation

Ms De Waard Jolien
j.de.waard01@umcg.nl - Netherlands

De Waard Jolien
j.de.waard01@umcg.nl - Netherlands

#3652

CIN3+-specific methylation marker analysis to improve the triage of hrHPV-positive self-samples in the population-based cervical cancer screening

18 - Methylation

De Waard J, De Boer M, Bhattacharya A, Vermeulen K, De Bock T, Schuurin E, Wisman B

Background/Objectives: The current Dutch population-based screening (PBS) programme for cervical cancer consists of primary high risk human papilloma virus (hrHPV) testing with cytology as triage test on cervical scrapings. Material is collected by a general practitioner (GP) or by the woman herself using a self-sampling device. One disadvantage of the self-sampling device is that hrHPV-positive women need to visit their GP to determine whether referral for a colposcopy is needed, as cytological examination is not possible on self-collected material. This will result in delay of the correct diagnosis and less compliance to the PBS. These drawbacks may be circumvented by direct molecular triage testing using hrHPV-positive self-samples. Previously, we have reported that methylation marker analysis is feasible for molecular triage testing. The aim of this study is to determine the most optimal methylation marker panel to detect CIN3+ in hrHPV-positive self-sampled material.

Methods: We selected 23 methylation markers based on literature with high sensitivity and specificity for CIN3+ status (>70%) using GP-collected scrapings. Quantitative methylation-specific PCR (QMSP) was set up and performed on bisulfite treated DNA of 208 hrHPV-positive self-samples of women with a CIN2 or less (controls) and 96 women with CIN3 or worse (cases). Receiver operating characteristic (ROC) was performed to analyse the diagnostic performance by calculating the area under the curve (AUC), sensitivity and specificity.

Results: QMSP analysis of these individual markers showed discriminative DNA methylation levels between controls and cases in all markers ($p < 0.05$). ROC analysis for CIN3+ showed that 10/23 markers had an AUC of > 0.7 ($p < 0.001$). For 7/23 markers, sensitivity was $> 60\%$ with a specificity of $> 70\%$ for CIN3+.

Conclusions: Our study established promising methylation markers to be used in hrHPV-positive self-sampled material as putative triage test. To improve sensitivity, various methylation markers are presently combined to identify a panel with the highest combined sensitivity and specificity using cluster analysis and decision tree modelling. An external validation of this optimized panel will be performed in a large cohort of 2415 hrHPV-positive self-samples.

Oral presentation

Dr Olkov Ilya
777y77@gmail.com - Russia

Olkov Ilya
777y77@gmail.com - Russia

#3653

REALIZATION OF WHO GLOBAL STRATEGY ON CERVICAL CANCER ELIMINATION IN RUSSIA IN COVID-19 ERA

01 - HPV disease and COVID-19

Olkov I, Kononova I

Background/Objectives: To analyze the implementation of WHO Global Strategy on Elimination of Cervical Cancer in Russia during COVID-19 era.

Methods: Analysis of the HPV screening implementation, vaccination against HPV and treatment of patients with SIL was performed. To determine the adherence of the population to screening and vaccination a sociological study of 2900 medical workers and 236 590 women from various regions of Russia was carried out. To determine efficacy of HPV screening implementation 5 pilot regional screening programs were initiated in Russia for a cohort of 60 000 women at 30-49 years.

Results: Various screening techniques have been introduced in Russian regions: in remote areas PAP test is performed as part of a clinical examination of the population once every 3 years with detection of up to 22% of CIN and cervical cancer. In large cities and megalopolises of a number of regions, active HPV screening by HC2 is being introduced with detection of up to 75% of CIN and cervical cancer. 60 000 women from 30 to 49 years were enrolled into the screening program and 98,68% were tested. HPV High Risk positive and PAP positive women were referred for colposcopy. In this group 32,7% were diagnosed with various CIN grades and duly treated. The widespread introduction of modern screening technologies is prevented by limited resources during COVID-19 pandemic. Regional vaccination against HPV is being increased too, although lack of HPV vaccination in National Immunization Schedule remains critical. To facilitate implementation of the Global Screening Strategy, a unique model for the formation of professional competencies in training programs was developed, including visualization, virtualization, CIN treatment simulation environments, information management and CASE decision-making system, which increased competences of medical personnel by 12.5 times. Educational programs for the population included the use of innovative technologies for interactive discussion of HPV-associated diseases, enabled to carry out planned vaccination of 90 - 93% of children.

Conclusions: Implementation of WHO Global Strategy on Elimination of Cervical Cancer in Russia is being accelerated due to enthusiasm and determination of a number of medical activists, however, some problems remain and need to be addressed even in spite of a COVID-19 pandemic. We also faced some difficulties in the reaching of women enrolled into HPV screening regional programs during COVID-19 era. To overcome this, a complex informational and educational campaign was carried out both for local gynaecologists and local women with the help of local authorities, volunteers, mass media and new digital technologies.

References: I.R. Minniakhmetov, M.V. Zabelin, I.G. Olkov, R.I. Khusainova "Pilot project for cervical cancer screening using HPV testing" Problems in Oncology, Vol. 66, Dec 2020: 618-624 Kononova I.N., Bashmakova N.V., Dankova I.V., Vinokurova E.A. A model for the formation of professional competencies in training programs for medical workers in the organization of cervical cancer screening. - Russian Bulletin of Obstetrician-Gynecologist. 2019; 19 (2): 21-26

Oral presentation

Prof. Carozzi Francesca
frakaro@gmail.com - Italy

Carozzi Francesa Maria
frakaro@gmail.com - Italy

#3654

HPV screening in Bolivia: results by a Pilot study

39 - Low resource settings

Carozzi F, Royder Yanez R, Cannistrà S, Matucci M, Paganini I, Sani C, Von Borries S, Traina S

Background/Objectives: Cervical cancer is a major public health problem in Bolivia. Little is known about the prevalence of HPV infections in the Bolivian population and therefore about the feasibility of a screening program based on HPV and self-sampling. In 2019, the Italian Agency for Development Cooperation (AICS), La Paz regional office, conducted a pilot study to estimate the prevalence of High-risk human papillomavirus (hr-HPV) and the feasibility of HPV screening in Bolivia, using self-sampling and portable & Transportable lab Units for HPV test in urban and rural districts

Methods: The project enrolled women 20-65 age in 3 different areas: La Paz City, Toro Toro and Acasio (rural centers). Self-sampling was carried out with Viba-Brush® system and preservation in Thinprep® containers. We used Gene Expert system for HPV HR that detects E6/E7 of 14 HPV types in Real Time PCR, a system of easy execution with minimal manual intervention of the operator. The system provides the results in about 1h. Women testing positive for any HPV 16 or HPV 18/45 returned for colposcopy while women positive for other HR types performed cytology triage and if Pap test was positive women are referred to Colposcopy. Enrollment began in December 2019 and lasted until June 2021. Enrollment was particularly challenging as the period coincided with the COVID-19 pandemic and follow-up test retrieval is still ongoing.

Results: 2201 subjects were enrolled of which 811 (36.84%) in Acasio, 622 (28.26%) in La Paz and 768 (34.89%) in Toro Toro. Samples with valid HPV test (neg/pos for HPV) were 2076 (94.31%). Total positivity to hr-HPV was 12.72% with a classical trend by age. The two rural areas show lower positivity than women enrolled in the city (15.31% in La Paz, 10.57% in Acasio and 12.89% in Toro Toro; RR urban vs rural: 1.4; 95%CI: 1.04-1.80). The overall data from the 3 municipalities show that the highest frequency group is HPV 31/33/35/52/58, followed by HPV 39/56/66/68, HPV 51/59, HPV 16 and HPV 18/45, finding in part already reported in 2 previous Bolivian studies, despite the very low number of enrolled women. The number of colposcopies performed on screening cases is low and we are trying to retrieve them. However, from currently available data, the high-grade lesions are all related to HPV 16 or HPV 18/45, single or in co-infections with other HR-HPV

Conclusions: Prevalence of HR HPV infection, measured with a validated test for screening, showed total and by age estimates comparable to the ones observed in Europe and USA. This finding confirmed 1) the feasibility of applying a screening protocol based on HPV test with self-sampling and 2) the high prevalence of cervical cancer in Bolivia is likely related to the lack of screening programs, not to a higher occurrence of Hr-HPV infections. We adopted a screening system with pre-prepared cartridges that requires minimal intervention by the operator. Performing HPV tests locally allows to avoid the problems of transport and storage of the material collected, the possibility of having the woman waiting in the clinic for the test response of the negative screening test in an hour. In positive cases, the staff scheduled an appointment for the pap test or colposcopy, depending on the type of HPV present. Sustainability of HPV testing is certainly still a major problem. The cost is about 15 euros, too high for low-income nations. A joint action between the national scientific community, international institutions and manufacturers is needed.

Oral presentation

Dr Olkov Ilya
777y77@gmail.com - Russia

Kononova Irina
irkonmed@mail.ru - Russia

#3656

LOCAL IMMUNE STATUS RELATED TO SIL: SPECIALITIES AND POSSIBILITIES FOR IMMUNOMODULATION WITH SOLUTIONS TREATED WITH ULTRASOUND CAVITATION.

08 - Immunotherapy - Immuno-oncology - New treatments

Olkov I, Kononova I

Background/Objectives: To evaluate efficiency of immunomodulation in patients with unfavorable clinical course of local immune status of SIL with solutions treated by ultrasound cavitation.

Methods: Molecular-biological examination of vaginal microbiota was performed using immunohistochemical analysis of biopsy with expression of CD3, CD56, CD20, CD138, M-CSF receptors followed by evaluation of treatment efficiency by with peptide solutions treated by ultrasound cavitation in patients with LSIL with possible progression to HSIL using Fotek ultrasound cavitation technique.

Results: HSIL progression is accompanied by the dominance of *Lactobacillus iners* in combination with *Gardnerella v.*, *E. coli*, *Enterococcus faecalis* ($\chi^2 = 3421$ at $p = 0.006$, $\chi^2 = 32.1$ at $p = 0.003$, respectively). HSIL is characterized by an increase in the proportion of cells expressing receptors CD20 (1.5%), CD3 (45%), CD138 (in stromal cells) (12%) and a decrease in M-CSF epithelial infiltration ($r = -0.55$ at $p = 0.000$ for CD20 and for $r = -0.50$ at $p = 0.00$ CD3), which led to creation of a mathematical model for predicting neoplasia was. The sensitivity of the method was 75%, specificity - 92.3%, efficiency - 83.7%. Physio medical immunomodulation with peptide solutions treated by ultrasound cavitation was administered to patients with unfavorable prognosis. A 4,5 times reduction in the risk of neoplasia progression was observed.

Conclusions: Physio medical immunomodulation appears to be efficient in prophylactics of LSIL progression in patients with unfavorable clinical course of the disease.

References: 1. Patent for invention "Method for predicting the risk of progression of recurrent cervical intraepithelial neoplasias of low oncogenic risk associated with human papillomavirus infection", registration number: 2020127844 from 08.20.2020. 2. Shmakova N.A. Features of the local immune microenvironment in cervical intraepithelial neoplasia associated with human papillomavirus. / ON. Shmakova, I.N. Kononova, G.N. Chistyakova, I.I. Remizova // Bulletin of SurGU, Medicine. 2020. No. 4 (46). S.68-73. DOI: 10.34822 / 2304-9448-2020-4-68-73 3. I.R. Minniakhmetov, M.V. Zabelin, I.G. Olkov, R.I. Khusainova "Pilot project for cervical cancer screening using HPV testing" Problems in Oncology, Vol. 66, Dec 2020: 618-624

Oral presentation

Dr Gillis Jennifer
jennifer.gillis@cw.bc.ca - Canada

Gillis Jennifer L.
jennifer.gillis@cw.bc.ca - Canada

#3657

SELF-REPORTED PARTICIPATION IN CERVICAL CANCER SCREENING AMONG A VOLUNTEER COHORT OF VACCINATED WOMEN IN BRITISH COLUMBIA, CANADA

40 - Public health

Gillis J, Donken R, Marty K, Racey C, Smith L, Mitchell-foster S, Dawar M, Krajden M, Naus M, Van Niekerk D, Dobson S, Dionne M, Sauvageau C, Gilca V, Mayrand M, Money D, Kellner J, Sadarangani M, Ogilvie G

Background/Objectives: The province of British Columbia (BC) in Canada (population ~5 million) implemented a voluntary, school-based HPV vaccine program in 2008 for girls in grade 6 (age: 11-12). As in other jurisdictions with early HPV vaccine adoption, these individuals are now becoming eligible for population-based cervical cancer screening. To monitor participation in screening among vaccinated individuals, our objective was to quantify cervical cancer screening uptake in a cohort of young women who received the HPV vaccine and became eligible for screening between 2013 and 2019 in BC, Canada.

Methods: This analysis was conducted in a cohort of young women from BC participating in the QUadivalent HPV vaccine Evaluation Study (QUEST), which follows participants who received 2 or 3 doses of the HPV vaccine for up to 10 years post-vaccination. They received frequent newsletters about HPV throughout the study. Participants completed annual questionnaires and were asked to self-report having ever had a Pap test. Eligibility for screening was assessed according to BC guidelines, where women 21 years or older or who had been sexually active for 3 or more years were eligible for screening from 2013 to 2015. Eligibility changed to 25 years or older in 2016 with no additional eligibility criteria based on sexual activity. This descriptive analysis quantifies uptake of screening among eligible participants in BC.

Results: As of January 2021, 3333 of 3543 young women from BC in QUEST had completed at least one questionnaire; median age at first questionnaire: 16 years (range: 13-22). As per BC guidelines, 237 (7%) became eligible for screening between 2013 and 2019; the majority (94%) were eligible based on being sexually active for 3 or more years. Of those eligible for screening, 100 (42%) self-reported having had a Pap test, of whom 74% reported having had their Pap within 1 year of becoming eligible, 18% within 2-3 years, and 8% over 3 years after becoming eligible. Of the remaining 3096 young women not indicated for screening per guidelines, 301 (10%) self-reported receiving a Pap test, of whom the majority (96%) were sexually active. Most will become eligible for screening in BC by 2023.

Conclusions: In this volunteer, engaged cohort of vaccinated individuals, self-reported cervical cancer screening among screen eligible participants was low with less than half of those eligible reporting having been screened. However, to validate these results and monitor participation in cervical cancer screening as all participants become eligible, we will be linking QUEST data to administrative data available from the BC provincial cervical cancer screening program.

Oral presentation

Mr Beyers Koen
koen.beyers@novosanis.com - Belgium

Beyers Koen
koen.beyers@novosanis.com - Belgium

#3660

EXPLORATIVE STUDY ON DEVICE ARCHITECTURES FOR DEFINED RANGES OF FIRST-VOID URINE COLLECTION

13 - Self-sampling

Beyers K, Ríos Cortés A, Jordaens S, Meers N, Hendrickx J, Pasmans D, Tijsen J, Vankerckhoven V

Background/Objectives: A properly collected first-void urine (FVU) sample is important for accurate detection of e.g. sexually transmitted infections, human papilloma virus or cancer biomarkers. FVU contains a higher number of specific analytes, improving diagnostic accuracy. For this reason, Novosanis developed Colli-Pee®, a user-friendly FVU self-collection device. Moreover, Novosanis expanded its Colli-Pee® portfolio with specific collection tubes in different sizes and volumetric capacities (e.g. 20mL and 10mL) compatible with diagnostic carriers. The aim of this study was to explore the capability of the different Colli-Pee® variants to collect small, regular and large volumes, while maintaining the quality of the FVU sample and respecting the urine-to-preserved ratio.

Methods: Scaling of device architectures in combination with known limitations of injection molding techniques, requires complex miniaturization efforts while tackling unscalable laws of hydrodynamics, allowing for ranges of collected FVU from e.g. 4mL up to 40mL. Development items we segregated in: (i) a device platform that allows push-fit or threaded tube connection in different sizes with Colli-Pee®, (ii) an inlet funnel capable of processing a urine flow of up to 25mL/s, (iii) selection of hydrophobic polymers for clean sample collection, without loss of materials sticking to inner sidewalls, (iv) an inflow duct to access the collection tube, (v) immediate mixing with preservative, (vi) a floater with an air pocket, with sufficient buoyancy for lifting the valve assembly, (vii) a valve on top of the floater that closes access to the tube when a specific range of FVU has filled the tube, (viii) and the same valve that opens pass through flow of midstream urine through the device's outlet into the toilet. Banking on 2 patents, a number of conceptual designs, simulation models, 3D CAD drawings and prints, the final Colli-Pee® designs were manufactured in injection molded parts.

Results: The resulting differences in physical design between the volumetric variants showed similar performance with regard to FVU sample quality and relative collected range, guaranteeing proper sample stabilization with prefilled chemistry volumes. All designs proved to be well-suited for upscaling and mass manufacturing, with good sample stability, allowing additional clinical evidence to be acquired.

Conclusions: Currently, Colli-Pee® variants collecting small volumes (approx. 10mL, CE-IVD or approx. 4mL, RUO), regular volumes (approx. 20mL, CE-IVD), and large volumes (approx. 40mL, RUO) are available and performing well. These variants give customers and users the possibilities to tailor fit their diagnostic solution with analyte concentrations, high-throughput carriers, and home-based postal solutions.

Poster

Dr Jermakova Irina
opals19@inbox.lv - Latvia

Dr Jermakova Irina
opals19@inbox.lv - Latvia

#3662

CLASSIFICATION OF HIGH-GRADE CIN BY KI-67 STATUS-SIGNIFICANT CONSTITUENT ELEMENT IN PROGNOSIS OF PERSONALLY TAILORED MANAGEMENT .

23 - Diagnostic procedures / management

Jermakova I, Dace R, Mara P, Inta L, Una K, Jana Z, Olga P

Background/Objectives: Histological assessment of cervical biopsies is a "gold standard" to confirm cervical intraepithelial neoplasia (CIN) and its grading, but misinterpretation may occur due to intra- and interobserver variability. High-grade cervical intraepithelial neoplasia (CIN2 and CIN3) is a heterogeneous condition and with different potential for cancer progression. These difficulties mean that patients as well may be over-treated for CIN lesions, which will naturally regress, as undertreated. Biomarker Ki-67 indicative for a cellular proliferation could improve diagnostic accuracy and quality and could personalize management in women with high-grade CIN.

Methods: A cross-sectional study that enrolled a total of 110 women aged 18-65 with abnormal cytology referred for colposcopy to Reference Colposcopy Centre in Riga East Clinical University Hospital in July 2016-July 2017. Histological evaluation of cervical biopsy samples taken under colposcopy control was performed to each patient and all specimens were examined for Ki67 expression through IHC technique. The cumulative score of immunohistochemical expression Ki-67 (score 0 - 3), referred to as the "immunoscore" (IS), in the corresponding cervical scrape were evaluated.

Results: In the total group of CIN2/3 lesions (79 cases), 5 lesions were classified as IS group 0 - 2 (6.0%), 24 lesions as IS group 3 - 4 (30.4%) and 50 lesions as IS group 5 - 6 (63.6%), confirming heterogeneity within high-grade CIN lesions. Increasing Ki-67 expression was associated with increasing CIN grade ($p < 0.05$).

Conclusions: We have found a significant heterogeneity in the expression of Ki-67 in high-grade CIN lesions. Additional use of Ki-67 biomarker with classical histology interpretation complete each other to reach the higher accuracy on diagnosis of high-grade CIN, might help detect the prognosis and more personalised management so preventing overtreatment, especially in young women.

References: 1. Three-tierd score for Ki-67 and p16 improves accuracy and reproducibility of grading CIN lesions. van Zummeren M, Leeman A, Kremer WW, Bleeker MCG, Jenkins D, van de Sandt M, Heideman DAM, Steenbergen R, Snijders PJF, Quint WGV, Berkhof J, Meijer CJLM. J Clin Pathol. 2018 Nov;71(11):981-988. doi: 10.1136/jclinpath-2018-205271. Epub 2018 Jul 16. PMID: 30012698

Oral presentation

Dr Tan Jerry
jerrytan@truscreen.com - Australia

Chen Fei
yanfei19520344131@163.com -

#3676

PRELIMINARY PROOF AND SUGGESTION FROM THE CLINICAL DATA OF MULTICENTER RESEARCH ON THE APPLICATION OF REAL-TIME OPTOELECTRONIC DEVICE FOR CERVICAL CANCER SCREENING: IN THE REPEATED SURGES OF COVID-19 EPIDEMICS

22 - Artificial Intelligence

Background/Objectives: The main purpose of this study is to verify the feasibility of real-time optoelectronic device (TruScreen; TS) for cervical cancer screening based on the data of five hospitals, especially during the repeated surges of COVID-19 epidemics.

Methods: The participating hospitals in this study were divided into training set (n = 5) and test set (n = 64). A total of 17447 participants were included in the study and tested by cytology, HPV, TS and gynecological examination from September, 2018 to June, 2021. When each test is positive result, the patient will go forward to colposcopy. The informed consent forms were signed by all participants. The inclusion criteria were female patients older than 20 years with sexual activity. Patients who had acute infections in the lower genital tract, pregnant or undergoing treatment of the cervix were excluded.

Results: Participants in both training set and testing set have completed case collection and all tests. 1,022 women data of the training set were analyzed in the study. 62.4% of the women were cytological ASC-US+, 91.3% were high-risk HPV positive. TS test was abnormal in 41.5% women. 261 women were diagnosed as CIN2+ lesions. The highest risk of CIN2+ was 0.61 for women with HPV 16/18+ and abnormal TS results. For detection of CIN2+, the sensitivity and specificity for TS was 90.4% and 75.3%, and sensitivity was statistically higher than cytology (82.4%). The specificity of cytology and hrHPV DNA testing were significantly lower than that of TS ($P < 0.001$). Participants in the test set have completed case collection and all tests.

Conclusions: TS has the advantages of good sensitivity, specificity, objectivity and real-time. It is a noninvasive, simple and rapid method for the diagnosis of cervical lesions. TS is particularly suitable as primary screening method for cervical cancer that women and medical staff facing SARS-CoV-2 exposure and infection risks in hospital during the repeated surges of COVID-19 epidemics.

Oral presentation

Dr Li Lei
lileigh@163.com - China

Chao Xiaopei
xy7cxp@163.com - China

#3680

Novel methylated genes as a clinical predictor to reduce the missed diagnosis of cervical lesions (cancer) in women over 50 years old: a preliminary analysis of a multicenter study

18 - Methylation

Chao X, Li L, You Y, Wu H

Background/Objectives: Women over 50 years old have a higher risk of cervical lesions (cancer) than young population, but account for the majority of cervical missed diagnosis by three-step examination. Only a few articles pointed out non-invasive methods to reduce the missed diagnosis in postmenopausal women. Objectives: The main purpose of the study is to analyze whether the CisCer methylation genes (PAX1 & JAM3) can be as the new non-invasive method to increase the accuracy of cervical cancer diagnosis.

Methods: From 2019 to 2022, a prospective study of outpatient opportunistic cervical cancer screening was conducted in three hospitals with multiple centers. More than 20,000 subjects will be collected and all subjects will be follow-up for one year. The research team of Peking Union Medical College Hospital is responsible for preliminary experiment, clinical study planning, and process quality control. Cervical exfoliated cell specimen was collected and analysis for the system setup including SOP of sample collection, quality of specimen, and experimental operation. The analysis of methylation level was determined by using CisCer methylation real-time system (CISPOLY Co., China).

Results: A system set-up study was conducted on 905 subjects including normal uterine cervix (n=501), CIN1(n=170), CIN2(n=69), CIN3/CIS(n=110), SCC (n=15), and adenocarcinoma (n=17) of the uterine cervix diagnosed according to histological results. The sensitivity and specificity of CisCer methylation were 82% and 88% in all subjects and 87% and 86% in subjects over 50 years old compared with HPV16 / 18 (74% and 72%) and LBC (ASCUS; 96% and 24%). The cancer missed diagnosis rates of cancer of CisCer, HPV16 / 18, LBC (³ASCUS), and LBC (³LSIL) in women over 50 years old were 0%, 18%, 6%, and 29%, respectively.

Conclusions: The preliminary results indicated that the CisCer real time PCR-based testing is promised for cervical cancer detection with high sensitivity and specificity without cancer missed diagnosis for women over 50 years old.

Oral presentation

Ms Senese Francesca
francesca.senese@merck.com - Italy

Mrs Senese Francesca
francesca.senese@merck.com - Italy

#3684

Eliminating cervical cancer in Italy: results from mathematical modeling of HPV vaccination impact and DNA screening

36 - Economics and modelling

Background/Objectives: Since the late 1990's, when it expanded organized cervical screening programs (OCSPs), Italy has been working on the reduction of cervical cancer (CC) incidence. In 2007, Italy was one of the first EU countries to introduce a national human papillomavirus (HPV) vaccination program with gender neutral offered since 2017. Several regions also have multicohort approach including 25 y.o.a. ladies catch-up and immunization offered to women treated for HPV lesions. For CC screening, HPV DNA screening (introduced in 2014) has become an increasingly important component of the national screening program becoming the standard from the age of 30. However, uptake and coverage of both interventions are heterogeneous across Italian regions. The WHO has defined three targets that are key for the elimination of CC as a public health problem (incidence below 4 per 100,000 women): 90% vaccination coverage, 70% coverage of screening, and 90% of women with cervical disease being treated. The aim of this study is to understand how these efforts are contributing to the elimination of CC in Italy on a national and subnational level, and to forecast what year elimination will be met.

Methods: A mathematical model of cervical HPV infection and disease was adapted to Italy on a national and sub-national level. The model accounts for historical changes in screening programs and HPV vaccination (including the transition to gender neutral vaccination, and a nonavalent vaccine) throughout Italy and was validated against documented drops in CC incidence following increasing coverage of OPSCs among target populations. At national and subnational levels, CC incidence was projected forward from 2021 and the year at which incidence reaches various thresholds of interest. Counterfactual scenarios were also run to ascertain the contribution of each intervention (HPV DNA screening and HPV vaccination) in achieving these thresholds.

Results: At the national level, screening alone, while responsible for significant near-term declines, is unlikely to lower CC incidence below 4 per 100,000 women. But when combined with HPV vaccination, this threshold is expected to be reached by approximately 2048. These results can vary among subnational regions.

Conclusions: Italy's efforts in CC screening and HPV vaccination have yielded significant benefits and will continue to drive CC incidence down over the next decades. Nevertheless, HPV immunization as a means of primary prevention is the key component of the strategy that will make the difference in whether elimination is reached, and higher vaccination coverage rates will contribute to an earlier year of elimination.

References: Li X, et al. Vaccine Impact Modelling Consortium. Estimating the health impact of vaccination against ten pathogens in 98 low-income and middle-income countries from 2000 to 2030: a modelling study. *Lancet*. 2021 Jan 30;397(10272):398-408. doi: 10.1016/S0140-6736(20)32657-X. Erratum in: *Lancet*. 2021 Feb 20;397(10275):670. Mennini FS, et al. Cost-effectiveness analysis of the nine-valent HPV vaccine in Italy. *Cost Eff Resour Alloc*. 2017 Jul 11;15:11. doi: 10.1186/s12962-017-0073-8.

Oral presentation

Dr Dong Zhe
dzpkuih@163.com - China

Dong Zhe
dongzhe@pkuih.edu.cn - China

#3685

New evidence of methylation biomarkers in cervical adenocarcinoma especially HPV independent type

15 - Molecular markers

Dong Z, Ren Z, Liu C, Xia W, Li T, Xiong G

Background/Objectives: Although cervical cancer screening and the use of HPV vaccine have greatly decreased the incidence of cervical cancer, the frequency of cervical adenocarcinoma (CAC) has increased these years. About 2/3 (60.3%) of CAC patients are younger than 50 years old. The three-step cervical cancer screening process is not suitable for early detection of CAC because the lesion is located in the cervical canal or HPV independent type. The main purpose of this study is to find clinical evidence of CisCer DNA-methylation test (methylated PAX1 and JAM3 gene) in CAC, especially HPV independent type.

Methods: 10 paraffin block carcinoma tissues and 17 liquid-based cytology (LBC) specimens of CAC were collected from Peking University International Hospital. The results of LBC, HPV genotyping test, immunohistochemical staining (p16 & ki67), and methylation test were compared and analyzed. Paraffin blocks and cervical exfoliated samples were subjected to DNA extraction, bisulfite conversion and DNA methylation detection in CisCer DNA methylation detection system according to the manufacturer's protocol (CISPOLY Co., Beijing, China).

Results: The 10 tissue samples of CAC included 4 common types, 3 neuroendocrine types and 3 gastric types. The positive rate of tests was high-risk HPV (hrHPV; 70.0%), HPV16/18 (60.0%), LBC (ASCUS; 70.0%), LBC (LSIL; 50.0%), p16 (partially +; 60.0%), ki67 (80.0%), and CisCer methylation (90.0%), respectively. The 17 cervical exfoliated cells of CAC were collected for cytology, HPV genotype, and methylation detection. The positive rates of tests were hrHPV (76.5%), HPV16/18 (58.8%), LBC (LSIL; 41.2%), LBC (ASCUS; 82.4%), and CisCer methylation (88.2%). In addition, the positive rate of CisCer methylation in rare cervical cancer was 100.0%, including gastric type (HPV independent) and neuroendocrine type, which were better than hrHPV and LBC testing.

Conclusions: In this study, new evidence from paraffin blocks and cervical exfoliated cell samples showed that the coincidence rate between CisCer methylation test and cervical adenocarcinoma was higher than that of other tests, especially HPV independent (gastric type). Methylation test can be used as a noninvasive detection of cervical adenocarcinoma including HPV independent and non-independent types in the future, which needs more evidence.

Oral presentation

Dr Qi Bingli
queenaqbl@163.com - China

Liu Shikai
commasll@163.com - China

#3686

Management of Minimally Abnormal Cervical Cancer Screening Test Results by DNA Methylation Detection

24 - Risk management

Qi B, Miao Z, Liu S

Background/Objectives: The ASCCP released consensus guidelines that formalized a strategy for management of cervical cancer based on estimates of underlying high-grade precancerous lesions or cancer (CIN3+). The remainder were minimal abnormalities including high-risk HPV (hrHPV; +)-cytology (-) and cytology (LSIL and ASCUS). Due to the great differences in cytological techniques, pathological CIN3 + missed diagnosis often occurs in minimally abnormal screening results, especially in areas with limited resources. The main purpose of this study is to find clinical evidence of methylated PAX1 and JAM3 genes in minimally abnormal cervical cancer screening test results, especially in ASCUS.

Methods: The subjects with minimally abnormal cervical cancer screening results were recruited in the Cangzhou central Hospital, China from 2019 to 2022. The inclusion criteria were female with age ≥ 20 and sexual experience. The exclusion criteria included: women had history of gynecological cancers, surgery of cervix and uterine body, or at pregnancy. The results of the methylated PAX1 and JAM3 genes (P&Jm) were determined by CisCer qPCR system (CISPOLY Co., Beijing, China). Positive rate, sensitivity, specificity, and accuracy for cytology, hrHPV and the methylation level of both genes were analyzed. Epidata3.1 was used for data entry and SPSS18.0 software was used for statistical analysis.

Results: Total 1200 subjects were recruited and analyzed in the study. The results showed that the CIN2+ of patients with minimal abnormalities was higher than 5 %, especially in ASCUS group. The final diagnosis was confirmed by colposcopy or histology of colposcopy directed biopsy. The results showed that the combination of P&Jm genes was significantly higher in patients with minimal abnormalities in CIN3+ than those in CIN2, CIN1, and normal cervix ($P < 0.001$). The sensitivity of P&Jm for CIN3+ were more than 75%, and the specificity and positive predictive rate were more than 90%, respectively. P&Jm still shows good results for clinical application on ASCUS and hrHPV triage in the study. P&Jm test reduces more than 20% colposcopy referral rate without cancer missed diagnosis than patients with minimally abnormal-HPV16/18 positive results. Compared the patients with minimal abnormalities with HPV16 / 18 (+) results, P&Jm detection can reduce the referral rate of colposcopy by more than 20%, and there is no missed diagnosis of cancer.

Conclusions: The current results show that the combined detection of methylated PAX1 and JAM3 genes is a promising triage biomarker for the patients with minimal abnormalities in the primary screening of cervical cancer in Chinese population. The single center clinical evidence can improve the clinical effectiveness and decrease the numbers of colposcopy and follow-up without cancer missed in the study.

Oral presentation

Prof. Zhang Mengzhen
13683806877@163.com - China

Zhang Mengzhen
13683806877@163.com - China

#3687

CLINICAL SIGNIFICANCE OF METHYLATED ZNF582 AS A TRIAGE MARKER FOR HR-HPV INFECTED WOMEN

18 - Methylation

Zhang M

Background/Objectives: The aim was to investigate the value of the methylated genes and cytology as shunt in hr-HPV infected women, especially in women with cervical transformation zone III (TZ III).

Methods: Total of 204 patients infected with hr-HPV were enrolled in this study from the First Affiliated Hospital of Zhengzhou University. Cervical exfoliated cells were collected for liquid-based cytology (LBC) and ZNF582 / PAX1 methylation (ZNF582m / PAX1m) tests before the colposcopy and biopsy. All enrolled patients were divided into normal/cervicitis (n=59), CIN1 (n=42), CIN2 (n=38), CIN3 (n=40), and cervical cancer groups (n=25) according to pathological results.

Results: In the hr-HPV infected women, the sensitivity / specificity of ZNF582m, PAX1m and LBC tests for detecting CIN3 + were 70.2% / 89.9% , 58.4% / 87.8% and 70.8% / 60.4%, respectively. The sensitivity of the ZNF582m and LBC tests for CIN3+ were significantly higher than PAX1m test ($p<0.05$) and the specificity of the ZNF582m and PAX1m tests for CIN3+ were significantly higher than LBC test ($p<0.01$). In the hr-HPV infected women with TZ III, the sensitivity of ZNF582m and PAX1m for detecting CIN3+ were 88.5% and 69.3%, which were similar to that (80.8%) of LBC ($p>0.05$), and the specificity of ZNF582m and PAX1m were 90.9% and 88.3%, which were significant higher than that (57.1%) of LBC ($p<0.01$).

Conclusions: The ZNF582m test was a promising triage marker for the hr-HPV infected women in China, which was superior than LBC and PAX1m tests.

References: none

Oral presentation

Ms Olthof Ellen
e.olthof@erasmusmc.nl - Netherlands

Ms Olthof Ellen
e.olthof@erasmusmc.nl - Netherlands

#3688

THE IMPACT OF COVID-19 ON THE DUTCH CERVICAL CANCER SCREENING PROGRAMME

01 - HPV disease and COVID-19

De Kok I, Siebers A, Van Kemenade F, Aitken C

Background/Objectives: The COVID-19 pandemic has had a major impact on cancer screening programmes worldwide. The Dutch government decided to temporarily stop the cervical cancer screening programme in March 2020 by not sending new invitations, reminders or self-sampling kits to women. Following the end of the first lockdown in June 2020, there was a progressive resumption of the programme. Additional catch-up measures were taken in October-November to speed up invitations and participation, such as sending out 120% of the usual monthly invitations and offering more prominent self-sampling in the invitation letter. We aimed to investigate the effect of the COVID-19 measures on short term outcomes of the Dutch cervical screening programme, by comparing these with data from 2018 and 2019.

Methods: Data from the national information system for cervical cancer screening (ScreenIT, Topicus, Deventer, the Netherlands) and the nationwide network of cyto- and histopathology in the Netherlands (PALGA) were used for this study. Women eligible for screening in 2018, 2019 and 2020 were included. These three cohorts were compared using the following indicators: participation rate stratified by age, region and screening history; time between invitation and participation; type of screening test used (self- or clinician sampled); time between an hrHPV positive self-sampling result and reflex cytology; hrHPV positivity; cytology result and direct referral rates.

Results: In total, 297,076 women participated in the Dutch cervical screening programme in 2020. The participation rate in 2020 (49.8%) was lower than that of 2018 (57.6%) and 2019 (56.1%). The decrease in participation was partly caused by a low participation rate of women invited between January and March 2020. This group also participated later compared to previous years. The use of self-sampling was higher in 2020 (16.1% of all participants) compared to 2018 (6.7%) and 2019 (8.6%). This was observed in all age groups and screening regions. There were no clinically relevant differences observed for the other indicators.

Conclusions: The interruption of the Dutch screening programme due to COVID-19 had impacted women who were invited in the first three months of 2020 the most; these women had lower participation rates compared to women invited in the same months in 2018 and 2019. The broader offer of self-sampling that was introduced in November 2021 had clearly an effect on the number of women who have used self-sampling. It is recommended to send re-invitations to women who were invited between January and March 2020 and did not participate to ensure these women have sufficient opportunity to participate.

Oral presentation

Dr Giachini Claudia
c.giachini@ispro.toscana.it - Italy

Giachini* Claudia
c.giachini@ispro.toscana.it - Italy

#3689

Peer Review Cytology Plus (PRCP): a new software for the Internal Quality Control in cytology

21 - New technologies

Giachini* C, Pompeo G, Meoni M, Cannistrà S, Paganini I, Di Stefano C, Sani C, Carozzi* F, Matucci* M

Background/Objectives: * These authors contributed equally. Peer Review is an internal quality control (IQC) system used in the laboratory of ISPRO (Institute for Cancer Research, Prevention, and Clinical Network) - where about 60,000 Pap tests from all over Tuscany are performed annually. It consists of the daily examination, by all cytologists of the lab, of cervical smears judged abnormal or of difficult interpretation, which emerge routinely. The examination is performed blindly, without opinion exchanges, and each cytologist records his/her own evaluation. Smears are then collegially discussed and reviewed using a multi-head microscope and a consensus on the final diagnosis (FD) is reached; FD generally corresponds to the majority diagnosis (MD). The aim of the study was the implementation of software, called Peer Review Cytology Plus (PRCP), designed to simplify ICQ data recording and statistical analysis aimed to improve the evaluation of the laboratory performance.

Methods: Software design has been driven by the non-computerized procedure with which peer review has always been performed in our laboratory since 1989 (1,2). The main goal was to achieve the following aims: - to simplify the recording of data of the peer review process by each cytologist, which was previously done without computerized tools - to generate a report for each smear with the following elements: the majority diagnosis (MD), automatically calculated; an histogram showing the distribution of the results recorded by each reader; the final and histological diagnosis (HD). - to generate a report for statistical analysis aimed to evaluate: i) the participation (%) of each reader to the peer review sessions; ii) the level of concordance of each cytologist with MD, FD and HD; iii) the level of agreement among readers.

Results: The Peer Review Cytology Plus (PRCP) has already been successfully introduced in our laboratory; some features are currently being implemented and validated such as the report showing the level of concordance of each cytologist with MD, FD and HD and the level of agreement among readers. Among the advantages of the introduction of this computerized tool are the complete traceability of all cases sent for peer review and the possibility of carrying out statistical analysis to evaluate the performance of each cytologist and of the whole laboratory. Moreover, this software allows comparing the diagnosis distribution of different smears from the same patient (for example baseline vs follow-up) as well as the revision of smears whose cytological and histological results were discordant.

Conclusions: The Peer review procedure represents an important method of IQC; the continuous on-going discussion of positive and difficult patterns among cytologists with different levels of experience should result in an improvement in the uniformity of reading and, consequently, in the achievement of optimal values of reproducibility, as well as improving levels of diagnostic accuracy. The Peer Review Cytology Plus (PRCP) software represents a very important evolution allowing the reduction of time-consuming for the daily IQC - through the facilitation of the peer review process - and the strictly monitoring of the laboratory performances in terms of inter-observer agreement, reproducibility and functioning of the whole laboratory.

References: 1. Palli D, Confortini M, Biggeri A, Russo A, Cariaggi P, Carozzi F, Minuti PA. A quality control system involving peer review of abnormal cervical smears. *Cytopathology*. 1993;4(1):17-25. 2. Confortini M, Di Stefano C, Biggeri A, Bulgaresi P, Di Claudio G, Grisotto L, Maddau C, Matucci M, Petreschi C, Troni GM, Turco P, Foxi P. Daily peer review of abnormal cervical smears in the assessment of individual practice as an additional method of internal quality control. *Cytopathology*. 2016 Feb;27(1):35-42.

Poster

Liu Xin
Xin.Liu@kreftregisteret.no - Norway

Liu Xin
Xin.Liu@kreftregisteret.no - Norway

#3690

Human Papillomavirus genotype specific concordance between physician-taken and self-collected cervical samples

13 - Self-sampling

Nygård S, Stratford E, Leinonen M, Lahlum E, Nygård M

Background/Objectives: Human papillomavirus (HPV) self-sampling does increase participation in cervical cancer screening among unscreened or under-screened women and HPV genotype specific guidelines improve clinical management of high-risk (hr) HPV- positive women. We have demonstrated earlier good concordance for hr HPV DNA detection between self-collected and cervical samples collected by physician, but comparative data on HPV-genotype specific concordance between different self-collection methods is limited. Our study aim to 1) Assess the positivity rate for selected HPV-genotypes separately and in combination (HPV-genotypes grouped as 2 v", "4 v" and "9 v" based on HPV types targeted by prophylactic vaccines and the combined carcinogenic HPV types which excluded carcinogenic HPVs targeted by 2 v or 4 v and 9 v vaccines).2) Evaluate concordance in detecting HPV genotypes and in combination between self-taken and physician-taken samples using two sampling devices.

Methods: Study participants were recruited from the treatment units for premalignant lesions and cancer care units. Each participant collected at home the day before their appointment at the hospital two cervicovaginal specimens using a brush (Evalyn®Brush) and a swab (FLOQSwabs™). Before the treatment, a physician took a cervical specimen using a brush, into PreservCyt® buffer for HPV testing. All specimens were tested using Anyplex™ II HPV28. We estimated HPV-genotype specific and combined positivity rate for each sampling method. Cohen's kappa (κ) was used to measure agreement of self- and physician-collected specimen regarding positivity to HPV-genotypes alone and combined. 95% confidence intervals (CI) were estimated using the bootstrap method provided in an ado file, kapci, written by Michael E. Reichenheim using Stata (version 15.0 StataCorp, College Station, TX, USA).

Results: 251 women were included in the final analysis. 43 women with normal or CIN1 diagnosis, 160 with CIN2+ diagnosis, 48 with cancer diagnosis. Of 48 histologically confirmed cervical cancers, 30 were squamous cell carcinomas, 16 adenocarcinomas and 2 of other type. The overall HR HPV positive rate was 89.2%, 89.6% and 81.6% separately for Physician sample, Evalyn Brush and FLOQSwab. The overall HR HPV, 2-valent vaccine types, 9-valent vaccine types, Non 2-valent HR HPV types and non 9-valent HR HPV types positive rate was 89.2%, 55.4%, 80.9%, 33.9%, 8.4% respectively. The kappa value for Evalyn brush on HPV16, HPV18, HPV31, HPV33, HPV52, any HR HPV type, 2-valent vaccine types, 9-valent vaccine types, non 2-valent vaccine HR types, non 9-valent vaccine HR types was 0.85, 0.81, 0.90, 0.94, 0.79, 0.81, 0.78, 0.80, 0.80 respectively. The kappa value for FLOQswab on HPV16, HPV18, HPV31, HPV33, HPV52, any HR HPV type, 2-valent vaccine types, 9-valent vaccine types, non 2-valent vaccine HR types, non 9-valent vaccine HR types was 0.74, 0.72, 0.90, 0.86, 0.62, 0.67, 0.61, 0.69, 0.60 respectively.

Conclusions: 1.Specimens collected by different sampling methods resulted in comparable positivity rates in detecting HPV genotypes alone and in combination; 2.Compared to swab-based device, brush-based device has higher concordance with physician taken specimens for individual HPV and for HPV genotypes combined compared to swab based device 3.Self-sampling can be used for detecting HPV genotypes, but device specific differences exist.

Oral presentation

Dr Schmitz Martina
martina.schmitz@oncgnostics.com - Germany

Hums Anna-bawany
anna-bawany.hums@oncgnostics.com - Germany

#3692

Clarification testing/triage of women tested HPV DNA-positive in cervical cancer screening using a DNA methylation marker-based test as well as an HPV mRNA test

15 - Molecular markers

Campana G, Knoll K, Schuster C, Zimmermann-marroquin J, Schmitz M, Hansel A, Hums A

Background/Objectives: HPV-testing is more and more implemented in cervical cancer screening, due to its higher sensitivity compared to cytology. As HPV screening is less specific, however, reliable triage methods are essential in such a setting in order to avoid overtreatment and higher screening costs. DNA methylation markers may provide a suitable tool especially with respect to keeping false-positive rates low. GynTect® is a DNA methylation assay using six human genomic marker regions, ASTN1, DLX1, ITGA4, RXFP3, SOX17, and ZNF671. Its utility in screening triage was assessed, especially in comparison to HPV mRNA testing.

Methods: Cervical smear samples from 231 women with cytology findings \geq Pap III, collected in a new liquid-based cytology medium (BestPrep®; CellSolutions), were selected for the study. For all samples results from HPV DNA testing (Roche Cobas HPV test) were available. For the study, HPV mRNA testing was performed using the Aptima HPV test (HOLOGIC). Testing for the six DNA methylation markers comprising GynTect was performed for all samples following the instructions of the supplier.

Results: The overall positivity for both tests within this cohort was higher for HPV mRNA than for GynTect (155 vs. 86 samples). Regarding cytology, 170 of the 231 cervical smear samples fell into the Pap III group, 53 into the Pap IV group, and 8 into the Pap V group. For both tests, positivity was significantly lower in the Pap III group than in the Pap IV group (HPV mRNA 64% vs 85%; GynTect 24% vs 81%). Of the 231 samples 33 were from women tested HPV DNA-negative, 94 were from women at least positive for one of the HPV types 16 or 18, and 104 samples were from women positive for any of the other 12 high-risk HPV types. Of the 33 HPV-negative samples, five were positive for HPV mRNA, and six were positive for GynTect. Two of the samples were positive for both tests. Among the 94 samples being HPV16 and/or HPV18-positive, 70 contained HPV mRNA, and 57 were GynTect-positive. Concordant positive results for both tests were obtained for 51 samples. Among the 104 samples with (at least) one of the other 12 high-risk HPV types, 80 were positive for HPV mRNA, whereas only 23 were positive for GynTect. Of these, 22 were also HPV mRNA-positive. The remarkable difference between HPV mRNA positivity and GynTect positivity in this group is mainly seen within those samples that showed a Pap III cytology. Of these 90 samples, 68 (75.6%) were HPV mRNA-positive, whereas only 12 (13.3%) were GynTect-positive. 11 of these 12 GynTect-positive samples were also HPV mRNA-positive.

Conclusions: Detection of mRNA of the HPV oncogenes E6 and E7 provides a sign for a persisting infection with HPV and thus a higher risk of the patient for developing a high-grade cervical lesion. In contrast, the DNA methylation markers comprising GynTect are a direct sign for carcinogenesis. Therefore, it is not too surprising that triage using GynTect may show much higher specificity than using HPV mRNA. Interestingly, the main difference in positivity of both tests is seen in the group of samples that are HPV DNA-positive and show a cytology Pap III. Thus, GynTect may help to clarify the malignancy status of HPV-positive women with cytological signs of max. mild dysplasia.

Poster

Prof. Inácio ângela
minacio@medicina.ulisboa.pt - Portugal

Bicho Maria Clara
mclara.bicho@gmail.com -

#3693

ASSOCIATION OF CBS, p22Phox, NOS3 AND GSTM1 GENES WITH CERVICAL CANCER OR PRECURSOR LESIONS

15 - Molecular markers

Background/Objectives: HPV is the leading necessary cause of cervical cancer, a malignant tumour that initiates in the cells of the cervix and later may spread to other parts of the body. CBS, p22Phox, NOS3 and GSTM1 are genes involved in several events crucial to tumour formation and biology, such as the body's response to oxidative stress. The purpose of this study was to understand the role of the variants 844Ins68 (CBS), C242T (p22Phox), 27-bp VNTR 4b/a (NOS3) and GSTM1-0/1 (GSTM1) in the development of HPV caused cervical cancer or precursor lesions.

Methods: The genotyping was made by PCR in the CBS and NOS3 genes, by PCR-RFLP in the p22Phox gene and by PCR-multiplex in the GSTM1 gene.

Results: Results show an association of the -/+ genotype of the CBS gene ($p = 0.036$; OR = 0.219 [0.047-1.018]), the CC genotype of the p22Phox gene ($p = 0.012$; OR = 2.057 [1.166-3.63]), the 4b4b genotype of the NOS3 gene ($p = 0.042$; OR = 2.032 [1.019-4.052]) and the GSTM1-1 ($p = 0.047$; OR = 0.594 [0.355-0.994]) of the GSTM1 gene with the disease.

Conclusions: Altogether, results demonstrate the relevance of mechanisms related with ROS formation and oxidative stress response in the onset of cervical cancer or precursor lesions.

Oral presentation

Saxena Kunal
kunal.saxena@merck.com - United States

Saxena Kunal
kunal.saxena@merck.com - United States

#3695

Burden of Cervical Conization in Commercially Insured Young and Mid-Adult Women in the United States

26 - Cervical neoplasia

Saxena K, Yande S, Sawhney B, Chatterjee S, Kathe N

Background/Objectives: The Advisory Committee on Immunization Practices (ACIP) recommends catch up Human Papilloma Virus (HPV) vaccinations for women ages 13-26 and shared decision making between health care providers and mid-adult (ages 27-45) patients who may benefit from HPV vaccine. However, providers may be hesitant to recommend vaccinations because of lack of data on HPV burden in the adult population. Our analysis estimates the incidence and health care costs for treating cervical lesions, i.e; cervical intraepithelial neoplasia grades 2, 3, and adenocarcinoma in situ (CIN2+) with loop electrosurgical excision procedure (LEEP) or cold knife conization (CKC) among commercially insured women, which if untreated may progress to cervical cancer.

Methods: This retrospective cohort study using IBM MarketScan commercial claims database included women aged 18-45, if they had at least 6 months of continuous enrollment, diagnosed with CIN2+, and treated with one of the conization procedures (CKC or LEEP). The annual incidence of conization was estimated from 2016-2019, stratified by age groups: 18-26 years, and 27-45. Adjusted two-year health care costs post conization were computed using multivariable general linear model (GLM) with a gamma distribution accounting for follow-up time, sociodemographic characteristics, and comorbidities, stratified by the same age groups.

Results: The inclusion criteria were met by 6735 women, mean age 33.9 years (SD=6.2). Annual conization incidence declined each year, starting at 156 procedures/100,000 women-years in 2016 to 143 procedures/100,000 women-years in 2019. Conization incidence was lowest for women ages 18-26 (41/100,000 to 62/100,000 women-years) and highest for women ages 31-35 years (243/100,000 to 269/100,000). The adjusted health care costs were \$7,279 and \$9,249 in the 18-26 and 27-45 age groups, respectively. The adjusted costs for disease-specific care were \$3,609 and \$4,557 for women ages 18-26 and 27-45 respectively.

Conclusions: Conization burden was found to be high across all age groups, especially in women aged 31-35 years old. The results from this assessment highlight the importance of catch-up vaccination for adult women to preserve their health and reduce future healthcare costs.

Oral presentation

Ms Pankakoski Maiju
maiju.pankakoski@cancer.fi - Finland

Ms Pankakoski Maiju
maiju.pankakoski@cancer.fi - Finland

#3696

IS ORGANIZED CERVICAL CANCER SCREENING MORE EFFECTIVE THAN OPPORTUNISTIC TESTING? A POPULATION-BASED CASE-CONTROL STUDY

40 - Public health

Pankakoski M, Ahti A, Tytti S, Sirpa H

Background/Objectives: The aim was to study the effectiveness of cervical screening, both organized and opportunistic, on the risk of cervical cancer.

Methods: We conducted a case-control study with 1600 women with invasive cervical cancer diagnosed in 2010—2019 in Finland. On each case, 10 controls were individually matched on age and hospital district. Data on organized and opportunistic screening for these women were derived from the mass screening registry, the pathology laboratories, and the health insurance reimbursement registers. Cervical cancers were linked from the cancer registry. Conditional logistic regression was used to calculate odds ratios for associations between cervical cancer and screening participation three and five years before cases' diagnosis by mode of testing (organized program, unorganized or both). Models were adjusted for socioeconomic status, education, and mother tongue.

Results: Preliminary results will be presented at the conference.

Conclusions: There seems to be little additional preventive effect of opportunistic screening if a well-organized nationwide screening program is in place.

Oral presentation

Dr Nobre J. Guilherme
joseguilhermenobre@campus.ul.pt - Portugal

Nobre J. Guilherme
joseguilhermenobre@campus.ul.pt - Portugal

#3697

THE INTERPLAY BETWEEN HPV, CVM DYSBIOSIS AND CERVICAL CANCER DEVELOPMENT (MICROCERVIXHPV STUDY)

19 - Microbiome

Nobre J, Matos A, Santos A, Veiga L, Brito M, Pires M, Cardoso C, Bicho M, Bicho M

Background/Objectives: The importance of Cervico Vaginal Microbiota (CVM) in the protection against infections (as Human Papilloma Virus (HPV)) is already well established, namely through Lactobacillus species. Furthermore, is it also well-known the influence of HPV in the development of Cervical Neoplasia (CN). However, it is not possible to classify HPV as a complete carcinogen. That is why CVM dysbiosis should be explored, intending to decipher this curious interplay with HPV. This work aims to compare CVM composition of healthy women without HPV, HPV infected women and women in several Cervical Intraepithelial Neoplasia (CIN) and CN stages. Furthermore, it is intended to evaluate mortality and morbidity outcomes amongst all the groups as well as characterize the vaginal cytokine microenvironment, regarding each group and each CVM composition.

Methods: This is a national, longitudinal, observational and retrospective study that has 4 experimental groups (3-6) and 2 control groups (1,2), with 25 participants in each group, with the following composition: 1) HPV negative and without cervical lesions; 2) HPV positive and without cervical lesions; 3) HPV positive and CIN1; 4) HPV positive and CIN2-3; 5) HPV positive and CN without metastasis (T1-4 N0-2 M0); 6) HPV positive and CN with metastasis (T>2 N+ M+). DNA was isolated from cervicovaginal exudate samples, and then CVM, through bacterial 16s RNA was analysed by NGS. CVM cytokines were obtained through the analysis of cervicovaginal exudate in a Multiplex platform. Statistical analysis was performed in SPSS v 24.0. An alpha of 0.05 was considered statistically significant, and 95% confidence intervals were reported when appropriate.

Results: In terms of cytokine analysis, we expect a rise of inflammatory cytokines in patients with HPV infection, and the maintenance of this inflammatory process, could lead to CIN1-3 and, subsequently, CN. Furthermore, the levels of cancer-related cytokines might be dysregulated in patients with CIN1-3 and CN. Additionally, INF-gamma will probably be increased in all patients with active HPV infection. These cytokine profiles are in line with the changes observed in patients' CVM.

Conclusions: The importance of using cytokine profile and CVM is highlighted in the possibility of early detection/prevention of HPV and/or CIN/CN development. Furthermore, it can also be used as prognosis and treatment efficacy biomarker. Altogether, these insights are one step closer to personalized medicine.

Oral presentation

Dr Wendland Eliana Marcia
elianawend@gmail.com - Brazil

Wendland Eliana Márcia
elianawend@gmail.com - Brazil

#3698

PREVALENCE OF ANAL HPV INFECTION AMONG UNVACCINATED MEN WHO HAVE SEX WITH MEN IN BRAZIL: PRELIMINARY DATA

28 - Anal neoplasia

Kops N, Bessel M, Miranda G, De Brito E, Dall'aqua C, Camargo T, Da Silva I, Comerlato J, Dall Soto M, Fernandes B, Pilz C, Fetzner T, De Souza F, Pereira G, Wendland E

Background/Objectives: Persistent anal human papillomavirus (HPV) infection is the major cause of anal cancer. Despite being rare in the general population, its incidence is elevated in men who have sex with men (MSM). Therefore, we aimed to evaluate the prevalence of anal HPV among unvaccinated MSM in Brazil.

Methods: Cross-sectional study using respondent-driven sampling in 10 capitals cities across all geographical regions. The targeted sample size was initiated with three seeds in each city. For the anal sample, the participants were instructed to insert a Dacron humid swab 3 cm into the anal canal and rotate it 360 degrees at least twice. All samples were processed in a certified central laboratory using Anyplex II HPV 28 detection (Seegene®).

Results: A total of 628 unvaccinated MSM were recruited. Most were single (65.55%), with a mean age of 29.73 ± 9.42 years, 44.83% white skin color and with university education (77.73%). The overall HPV prevalence was 74.68% ($n = 469$), being 16.24% for HPV type 16 and 6.53% for HPV type 18.

Conclusions: These preliminary data show a high prevalence of HPV in the unvaccinated MSM Brazilian population. Further studies are warranted on the impact of HPV vaccination on anal cancer in this specific population.

Oral presentation

Dr Rasmussen Christina Louise
clr@cancer.dk - Denmark

Rasmussen Christina Louise
clr@cancer.dk - Denmark

#3701

INFLUENCE OF HPV AND P16 EXPRESSION ON SURVIVAL AFTER VULVAR SQUAMOUS CELL CARCINOMA: A POPULATION-BASED DANISH STUDY OF ~1,300 CANCERS

27 - Vulvar diseases and neoplasia

Rasmussen C, Thomsen L, Baandrup L, Frederiksen K, Franzmann M, Larsen L, Mejlgaard E, Schledermann D, Villena N, Waldstrøm M, Winberg B, Oernskov D, Kjaer S.

Background/Objectives: Some studies have shown that human papillomavirus (HPV) status may be a prognostic marker for survival after vulvar cancer, alone or in combination with p16 expression. However, the previous studies were small and had heterogeneous findings. In this large, nationwide study, we estimated survival after vulvar squamous cell carcinoma (SCC) according to high-risk (hr) HPV status, alone and combined with p16 expression.

Methods: In a nationwide pathology register, we identified all cases of vulvar SCC diagnosed at 13 pathology departments in Denmark during 1990-2017 (n=1545). Archived formalin-fixed, paraffin-embedded tumor tissue blocks were retrieved from clinical biobanks, and diagnoses were reviewed by a gynecopathologist. At a central laboratory, all samples were tested for HPV with the INNO LiPA Genotyping Extra test, and CINtec histology staining for p16 was performed. All p16 slides were evaluated by two gynecopathologists independently, and tumors with >70% p16 staining cells were considered positive. From nationwide registries, we obtained information on emigration and deaths in the study population during up to 28 years of follow-up. We estimated the 5-year overall survival (OS) according to hrHPV and p16 status by Kaplan-Meier's method, and the hazard ratio (HR) of 5-year mortality was estimated by Cox regression, adjusting for age and year of diagnosis.

Results: Of the originally identified 1545 vulvar SCCs, we included 1278 cases (83%) for whom an archived tissue block could be located, the diagnosis was verified and we had valid HPV and p16 results. The median age at diagnosis was 72 years (range 27-102). The majority of cases were histologically classified as SCC, not otherwise specified (67.0%), followed by keratinizing SCC (23.3%). The prevalence of hrHPV was 52.0% (95% CI: 49.1-54.6), and 31.0% (95% CI: 28.4-33.5) of cases were positive for both hrHPV and p16. hrHPV positive women had better survival than hrHPV negative women (5-year OS: 61.0% versus 43.9%; HR 0.82, 95% CI, 0.66-0.98). Furthermore, women positive for both hrHPV and p16 expression had better survival than women who were negative for both markers (5-year OS: 69.7% versus 43.9%, HR 0.67, 95% CI, 0.45-0.89).

Conclusions: These preliminary analyses support that patients with hrHPV positive vulvar SCC have improved overall survival compared with hrHPV negative patients. Combined positivity for HPV and p16 expression appears to be a stronger prognostic marker than positivity for HPV alone. Further analyses will be presented at the conference, including estimates of disease-specific survival and analyses adjusted for other prognostic factors, including tumor stage.

Oral presentation

Ms Lagheden Camilla
camilla.lagheden@ki.se - Sweden

Lagheden Camilla
camilla.lagheden@ki.se - Sweden

#3703

METAGENOMES IN HPV NEGATIVE CERVICAL CANCERS

19 - Microbiome

Lagheden C, Ure A, Arroyo Mühr L

Background/Objectives: Background: Human papillomavirus (HPV) is a necessary cause of cervical cancer, although some invasive cervical cancers may test negative by HPV PCR. HPV-test negative cervical cancers may constitute a biologically distinct subgroup, associated with symptomatic detection, late-stage diagnosis and worse prognosis that may need different targeted therapeutic strategies. It is thus essential to find other biomarkers to early identify these HPV PCR negative cancers. Objective: We aimed to analyze and compare the metagenomes present in HPV PCR positive cervical cancers with the metagenomes present in HPV negative cancers.

Methods: Methods: We previously requested all invasive cervical cancers in Sweden during 10-year period and subjected them to HPV genotyping using Luminex. FFPE (formalin fixed paraffin embedded) specimens that were HPV PCR negative (392/29850), together with a set of 59 HPV PCR positive specimens (used as positive controls), were further subjected to an unbiased total DNA and cDNA sequencing using Novaseq 6000 in order to detect "truly" HPV negative specimens. A total of 223/392 cervical cancer cases were still negative for HPV after sequencing. For this study, we randomly selected 49/223 HPV negative specimens and compared the metagenomes present with the metagenomes detected in the 49 HPV PCR positive controls using the Kraken2. A total of 6 blank paraffin blocks were used as negative controls.

Results: Results: Overall, 53 bacterial genera were detected with the genera Klebsiella, Staphylococcus and Pasteurella often contributing to more than 75% of the total bacterial reads for each sample. No significant bacterial difference was found among HPV positive and HPV negative specimens. Analysis of viruses revealed a total of 50 viral genera showing more than 1% of the viral reads. Gorganvirus (known to infect vagina), Orthobunyavirus and Alphapapillomavirus represented most of the viral reads. HPV was only detected in the HPV PCR positive samples and no other significant difference was detected among the HPV PCR positive and HPV negative specimens.

Conclusions: Conclusions: Metagenomic analysis of bacteria and viruses present in HPV positive and HPV negative cervical cancers shows no significant difference in bacteria or viruses present, besides the HPV for HPV positive specimens.

Oral presentation

Ms Eklund Carina
Carina.Eklund@ki.se - Sweden

Eklund Carina
Carina.Eklund@ki.se - Sweden

#3704

INTERNATIONAL QUALITY ASSURANCE OF HPV DNA GENOTYPING SERVICES: THE 2021 GLOBAL HPV DNA PROFICIENCY STUDY

09 - HPV testing

Eklund C, Lagheden C, Dahlin Robertsson K, Forslund O, Arroyo Mühr L, Dillner J

Background/Objectives: Objective: The International Human Papillomavirus (HPV) Reference Center supports quality and order in HPV research and diagnostics. Notably, the center assigns HPV type numbers to novel HPV types, maintains a reference clone repository, and issues international proficiency panels for HPV genotyping. This international HPV DNA genotyping study issued in 2021 assesses the proficiency of the different HPV typing assays used routinely in laboratories worldwide as well as the performance of the laboratory.

Methods: Method: Participating laboratories were asked to perform HPV typing using one or more of their usual assays on 41 coded samples composed of purified whole genomic plasmids of sixteen HPV types (HPV 6, 11, 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68a and 68b) in a background of human cellular DNA. Proficient typing requires detection in both single and multiple infections of 50 International Units of HPV 16 and HPV 18 DNA/ 5ml and 500 genome equivalents in 5 ml for the other types, with no false positive results.

Results: Results: The 2021 proficiency study had a record high number of participants: 161 panels were distributed to 138 laboratories worldwide: Most participants are from laboratories in Asia and Europe. Participating laboratories were both public health laboratories, research laboratories and diagnostic test manufacturers.

Conclusions: Implications and Impact: A continuing and increasingly popular global proficiency program promotes comparability and reliability of HPV genotyping assay performance worldwide.

Oral presentation

Mr Floore Arno
an.floore@self-screen.nl - Netherlands

Mr Floore Arno
an.floore@self-screen.nl - Netherlands

#3705

STREAMLINED WORKFLOW FOR METHYLATION ANALYSIS ON CERVICAL SAMPLES

18 - Methylation

Floore A, Arno F

Background/Objectives: Triage testing of HPV-positive women by methylation analysis to select women with (pre)cancerous lesions that need colposcopy is emerging. Methylation detection assays usually require pretreatment of the sample DNA with sodium bisulfite conversion to separate methylated from unmethylated cytosines. Streamlined workflows for this pretreatment that comply with the needs of laboratories for high-throughput testing are needed.

Methods: This study evaluated a direct cell conversion protocol on cervical samples in PreservCyt as alternative to isolated genomic DNA as input. The direct conversion protocol is capable of handling 96 samples simultaneously (manually or automated). Clinician-collected cervical samples (n = 120) were subjected to a direct conversion protocol, or genomic DNA was isolated with a fixed amount used for subsequent bisulfite conversion. Converted samples were compared for ACTB control gene and methylation of FAM19A4 and miR124-2 genes using quantitative methylation-specific PCR (QIASure Methylation Test).

Results: Direct conversion resulted in a high success rate, i.e., 119/120 (99.2%) samples reported a valid test result. $\Delta\Delta Cq$ values of FAM19A4 and miR124-2 were significantly correlated between both protocols (Spearman Rho 0.708 and 0.763, respectively).

Conclusions: A direct cell conversion protocol shows good technical and analytical performance and offers a streamlined workflow for methylation analysis.

Oral presentation

Mrs Da Silva Isabel Cristina Bandeira
isabel.silva@hmv.org.br - Brazil

Wendland Eliana
eliana.wendland@hmv.org.br - Brazil

#3706

The Quality of Anal Sampling for HPV Genotyping: Experience of an Epidemiological Study

14 - Genotyping

Bandeira I, Kops N, Comerlato J, Bessel M, Moreno Alves De Souza F, Mendes Pereira G, Wendland E.

Background/Objectives: High-risk HPV anal infection is associated with anal cancer. Screening for anal HPV among men who have sex with men (MSM) are crucial due to the increased incidence of anal cancer in this population. Therefore, we aimed to evaluate the quality of the anal self-sampling and associated factors in an epidemiological study.

Methods: We analyzed data from a cross-sectional, multicentric study about the prevalence of HPV among MSM in Brazil (SMESH study). Participants were instructed to self-collect HPV samples with a single Dacron swab soaked with saline solution. They swab must be introduced around 3 cm into the anal canal and rotate it 360 degrees at least twice. Sample collection was done under health professionals' supervision. The samples were submitted to DNA extraction (MagNA Pure LC 2.0 System, Roche) and subsequently genotyping by real-time PCR (AnyplexTM II HPV28 Detection, Seegene).

Results: Of the 1,177 samples genotyped, 420 (36.33%) showed some type of inadequacy in the collection process and only 22 (1.89%) of all samples were considered invalid to HPV genotyping. The most common inadequacy considered were tube with changed color (300 samples; 71.4%), presence of feces (146 samples; 34.76%), followed by tube without liquid (12 samples; 2.86%). However, none of the inadequacies were significantly associated with the invalid genotyping condition.

Conclusions: Few samples were invalid to HPV genotyping and the presence of inadequacies in the samples does not impact on HPV genotyping. The results reinforce that HPV detection through anal self-sampling is a valuable strategy to screen patients at high risk for anal cancer.

Oral presentation

Ms Kaljouw Sylvia
s.kaljouw@erasmusmc.nl - Netherlands

Kaljouw Sylvia
s.kaljouw@erasmusmc.nl - Netherlands

#3707

REALISTIC POPULATION-BASED COST-EFFECTIVENESS ESTIMATES FOR WIDER DEPLOYMENT OF HRHPV SELF-SAMPLING: A MODELLING STUDY

13 - Self-sampling

Kaljouw S, Jansen E, De Kok I

Background/Objectives: Background/Objectives: The hrHPV self-test has been suggested as an alternative to the hrHPV-test administered by the general practitioner to reach women who might otherwise not participate in cervical cancer screening. Recently, population-based data on loss-to-follow-up (LTFU) for reflex cytology (after self-sampling) (-7.8% LTFU) and relative sensitivity of the self-test have been reported in the Netherlands (0.94 relative sensitivity for CIN3+). We aimed to present population-based realistic cost-effectiveness estimates of wider deployment of self-sampling, taking into account this recent data.

Methods: Methods: We used simulation model MISCAN-Cervix to calculate the cost-effectiveness of several self-test deployment strategies based on feasible options to adjust the guideline (such as actively sending the self-test to all women, to 30-year old women or with the reminder letter) (Table 1). Scenarios varied in the percentage of originally participating women using self-sampling and the percentage of original non-attenders using self-sampling. Main outcome measures were life years gained, QALYs gained and costs compared to a scenario without screening. Table 1: Overview of strategies included in the analysis. The strategies vary in the percentage of self-sampling of all used tests without extra attendance and the percentage of extra attendance using self-sampling. Extra attendance (using self-sampling) Self-test deployment strategy % self-sampling of all used primary tests (without extra attendance) No extra attendance 10% of non-attenders 30% of non-attenders Women need to actively request the self-test 7% self-sampling all ages Current Send self-test with the reminder letter 7% self-sampling all ages (A) A10 A30 Send self-test to 30-year old women 50% self-sampling at 30 years old (B) B0 B10 B30 Send self-test to all women 30% self-sampling all ages (C) C0 C10 C30 Send self-test to all women 40% self-sampling all ages (D) D0 D10 D30 Send self-test to all women 50% self-sampling all ages (E) E0 E10 E30 * If two self-sampling percentages are given for a certain age, the highest percentage will be used when strategies are combined.

Results: Preliminary results: There was very little relative variation in the life years gained and QALYs gained for any of the scenarios (0.00-0.01%), however in absolute values 134 life years per 100,000 women can be saved if the attendance increases by 30% of non-attenders. Replacing 40% of the hrHPV tests at the GP by self-sampling without additional attendance leads to a decrease in costs (-3%), whereas increasing the attendance with the same strategy leads to an increase of 8% in costs. The highest costs are obtained when 7% of women use self-sampling with 30% additional attendance (+13%).

Conclusions: Conclusions: Taking into account the slightly lower relative sensitivity of self-sampling and loss-to-follow-up after self-sampling, we found that wider deployment of self-sampling is desirable. High attendance in combination with a high percentage of self-sampling is most cost-effective, due to the relatively low cost and slightly lower relative sensitivity of self-sampling compared to GP-based hrHPV-testing. More analyses are currently being done.

Oral presentation

Ms Middeldorp Marit
marit.middeldorp@rivm.nl - Netherlands

Mrs Middeldorp Marit
marit.middeldorp@rivm.nl - Netherlands

#3708

Vaccine effectiveness against persistent genital HPV infections up to ten years after three doses of the bivalent HPV vaccine

40 - Public health

Joske H, Middeldorp M, Audrey J K, Prof Johannes B, Hester E D

Background/Objectives: In the Netherlands, the bivalent HPV vaccine (2vHPV) has been offered to preadolescent girls via the National Immunization Program in a three-dose (3D) schedule between 2010 and 2014, and was replaced by a two-dose schedule from 2014 onwards. We aimed to estimate vaccine effectiveness (VE) against HPV persistent infections among girls eligible for a catch-up campaign 3D immunization up to ten years post-vaccination.

Methods: Data were used from a longitudinal cohort study (HAVANA) in which participants were annually asked to fill out a questionnaire and provide a vaginal self-swab for HPV determination using the SPF10-LiPA25 assay. Vaccination status of participants was acquired through the national vaccination registration system. Adjusted VEs against persistent HPV16/18 and cross-protective type (HPV31/33/45) infections following 3D immunization were estimated using the Prentice Williams-Peterson total-time approach. Persistent infections were defined as two HPV positive samples for the same HPV type in two consecutive years, preceded by a negative sample. Additionally, adjusted VE estimates up to 5 years post-vaccination and 5-10 years post-vaccination were estimated for HPV16/18 and HPV31/33/45. VEs were adjusted for age, urbanization degree, contraceptive use and sexual activity.

Results: A total of 1635 girls were included in the VE analyses, of whom 875 (53.5%) were vaccinated according to a 3D schedule. Preliminary data showed an adjusted pooled VE against persistent HPV16/18 infections of 95.8% (95% CI, 86.6-98.7%), and 64.6% (95% CI, 37.6-79.9%) against HPV31/33/45. Moreover, VE against persistent HPV16/18 infections amounted to 100% (95% CI could not be estimated) and 93.2% (95% CI, 78.1-97.9%), respectively for less than 5 years and 5-10 years post-vaccination. These estimates amounted to 43.4% (95% CI, -48.3-78.4%) and 71.9% (95% CI, 42.6-86.2%) for HPV31/33/45.

Conclusions: We found high VE estimates following 3D immunization against persistent genital infections in women in a population-based observational study after a 10-year follow-up. Furthermore, we did not find any indication of waning against persistent HPV16/18 and HPV31/33/45 infections over time. Continued surveillance of the HPV vaccination program is important to monitor future changes in protection or shifts in HPV trends over time.

Oral presentation

Ms Lujan Elena
lujan.e@procarehealth.com - Spain

Mr Gaslain Yann
lujan.e@procarehealth.com - Spain

#3710

EFFICACY OF A MULTI-INGREDIENT CORIOLUS VERSICOLOR-BASED VAGINAL GEL IN HIGH-RISK HPV WOMEN OVER 40: SUB-ANALYSIS OF THE PALOMA CLINICAL TRIAL & PAPILOBS REAL-LIFE STUDY

26 - Cervical neoplasia

Gaslain Y, Cortés J, De Santiago J, Gaslain Y

Background/Objectives: HPV clearance and resolution of cervical HPV-dependent lesions become difficult in peri and postmenopausal women. The objective of this analysis was to evaluate the effect of the Papilocare®, a multi-ingredient Coriolus versicolor-based vaginal gel in repairing the high-risk (HR) HPV-dependent low-grade cervical lesions in women over 40 years.

Methods: Paloma study (ClinicalTrials.gov NCT04002154) was a multicenter, randomized, open-label, parallel-group, watchful waiting approach-controlled clinical trial. Unvaccinated HPV positive women aged between 30-65 with cytology of ASCUS or LSIL and concordant colposcopy image were randomized into 3 groups: A) Papilocare® 1 cannula/day for 1 month + 1 cannula/alternate days for 5 months; B) Papilocare® 1 cannula/day for 3 months + 1 cannula/alternate days for 3 months; C) Control group: watchful waiting approach. Papilobs study (ClinicalTrial.gov: NCT04199260) was an observational, multicenter, prospective, one-cohort study. Vaccinated or not HPV-positive women aged > 25y with cytology of ASCUS or LSIL and concordant colposcopy were included. Patients were treated with Papilocare® 1 cannula/day for 21 days during first month + 1 cannula/alternate days for 5 months. Percentages of patients with normal cytology and concordant colposcopy after 6 months of treatment in the HR-HPV population are presented.

Results: A total of 30 and 68 HR-HPV patients above 40yo were evaluated in Paloma and Papilobs studies, respectively. In the Paloma trial, normal cytology and concordant colposcopy was observed in 90% vs 33% patients in A+B Papilocare® and control groups, respectively, ($p=0.003$, Fisher test). In the Papilobs study, normal cytology and concordant colposcopy was achieved in 73,5% of patients.

Conclusions: After a 6-month treatment period, Papilocare® showed a clinically robust and statistically significant efficacy in repairing cervical HR-HPV lesions in women over 40 years vs watchful waiting approach. This efficacy was corroborated in the real-life study in more than 2/3 of the HR-HPV patients above 40.

ASCUS/LSIL cervical lesions normalization

Poster

Ms Presthus Gro Kummeneje
gro.kummeneje.presthus@ahus.no - Norway

Mr Stosic Milan
milansto@oslomet.no - Norway

#3711

OPTIMIZATION OF AN NGS-BASED HPV-GENOTYPING PROTOCOL

14 - Genotyping

Presthus G, Lagström S, Andresen A, Costanzi J, Rounge T, Ambur O, Christiansen I

Background/Objectives: High analytical sensitivity and specificity of the HPV assay is important for epidemiological research and vaccine surveillance. HPV genotyping is also increasingly used for clinical purposes and a better understanding in terms of risk management is warranted. HPV detection and genotyping can be done precisely, efficiently and with high resolution by next generation sequencing (NGS). The aim of this study was to optimize an NGS-based genotyping protocol previously developed in our laboratory.

Methods: Material used in optimization studies included HPV-positive cell lines, WHO international standards, in-house plasmids for 37 genotypes harbouring HPV L1 sequences, as well as material from the Equalis proficiency panel. The in-house WHO validated PCR-based Luminex hybridization assay, Anyplex II HPV28 (Seegene) and in-house E6/E7 type-specific PCR were included as comparable methods. The first step of the NGS workflow, equal to the Luminex assay, is the amplification of an L1 gene fragment with modified general primers (MGP), followed by an index PCR. The performance of different DNA polymerases was assessed. Samples were pooled and quantified prior to sequencing, which was conducted on an Illumina MiSeq platform generating 2x151bp paired-end reads. The obtained data was analysed with an in-house genotyping pipeline combining Cutadapt to remove adapters, primer sequences, and low-quality bases, and Bowtie2 for aligning reads to HPV reference genomes retrieved from the PaVE database. featureCounts function was used for counting the mapped reads.

Results: The presented NGS method is time and cost-efficient and shows high-quality results for the identification of single HPV types, with identical sensitivity as Luminex in dilution series. For samples containing multiple HPV types however, some inconsistency in type identifications was observed, likely associated with stochasticity in PCR target hybridization. To address positive/negative HPV type classifications, several cutoff parameters were applied. Due to variation in amplification efficiency for the different HPV types, implementation of individual HPV type-specific cutoffs is highly considered.

Conclusions: Method optimization has initially been performed on plasmids and cell lines and showed consistent results for samples containing one HPV type. Further improvement is needed for multiple genotype identifications and optimization with emulsion PCR is planned. For confirmation of performance and for further validation, we are currently in an application process for the use of clinical diagnostic biobank material. Results and findings including clinical material will be presented.

Poster

Ms Lujan Elena
lujan.e@procarehealth.com - Spain

Dr Sanmartín Patricia
sanmartin.p@procarehealth.com - Spain

#3712

VULVAR CONDYLOMATA ACUMINATA: NEW LOCAL SYNERGISTIC TREATMENTS

32 - Genital warts

Sanmartín P, Estrada Z

Background/Objectives: Condylomata acuminata or genital warts are the clinical expression of low oncogenic risk (No. 6 and 11) human papillomavirus (HPV) infection. Genital infection by HPV is one of the most common sexually transmitted infections. However, condylomata acuminata are not included in surveillance systems in most countries, so worldwide epidemiological data is limited. The lack of a single effective treatment and the high post-treatment relapse rate, as new lesions appear in treated or untreated areas makes the management difficult. We present the clinical case of a young woman, presenting multiple vulvar condylomatous lesions previously evaluated and treated.

Methods: A 26-year-old patient visiting our office in CIMEG MADRID in August 2020 due to presenting multiple vulvar condylomatous lesions that had already been evaluated and treated in a dermatology center using local cryotherapy and sinecatechins (Veregen) ointment twice a day at home; she decided to seek out another medical opinion. In our unit, after adequate physical and colposcopy examination we decided to start local therapy with ErYag Laser (Fotona laser) with a wavelength of 2940nm together with a Coriolus versicolor-based external genital gel as an adjunct to local laser therapy. The gel was prescribed as co-adjuvant treatment and it was administered for 6 weeks at home to help re- hydrate, epithelialize and repair the area. At the end of the 6th week, a second local vulvar laser session was performed, and the Coriolus versicolor-based external genital gel was prescribed as co-adjuvant treatment again for a total of 6 months.

Results: After the 2nd laser session and 42 days of treatment, the patient was re-evaluated and fewer lesions were observed. The last evaluation was performed after 6 months of treatment and no the vulvoscopy was normal and no esions were observed macroscopically. Currently the patient is asymptomatic, with completed healed lesions.

Conclusions: The combination of Laser Therapy with the Coriolus versicolor-based external genital gel was suitable for this patient with a satisfactory evolution, showing total elimination of the lesions and absence of recurrences after 6 months of treatment.

Oral presentation

Ms Castañeda Kelly
k.m.castaneda.vanegas@umcg.nl - Netherlands

Castañeda Kelly
k.m.castaneda.vanegas@umcg.nl - Netherlands

#3713

HAS PARTICIPATION IN DUTCH CERVICAL CANCER SCREENING DECREASED IN THE LAST YEARS?

11 - Screening for women difficult to reach

Castañeda K, De Waard J, Vermeulen K, Greuter M, Van Der Vegt B, Siebers A, Sidorenkov G, Wisman B, Schuurin E, De Bock G

Background/Objectives: The benefits of regular cervical cancer screening (CCS) attendance are recognized worldwide (1). In The Netherlands, women aged 30-60 are invited to participate in the national CCS program every five years (2). In 2017, the screening changed from a cytology-based to an hrHPV-based program (3). Since then, the participation has decreased from 64% to below 57% (4). The participation rate is yearly provided by the CCS monitoring and is estimated within a 15-months window (4). As participation after that time window is not included in the estimates, this project aimed to evaluate the participation rate after the 15-months window and identify factors associated with delayed participation.

Methods: Data from the Lifelines, a population-based cohort in the north of The Netherlands, was linked to CCS-related data in PALGA (the nationwide registry of histo-cytopathology in the Netherlands) between 2001-2020. Women were included from the age of 30 till hysterectomy, death, or the age of 60. Women were categorized as timely participants and delayed participants when there was a primary screening result within 15 months or between 15-60 months after the start of the invitation year, respectively. The timely and the delayed participation rate was presented per year. The data were divided into 5-years blocks to identify the factors associated with delayed participation. For this, a multivariable logistic regression analysis was performed in each block, using age, education level, and pregnancy during the invitation year as covariates.

Results: Per year, on average, 13684 (SD:544) women in this cohort were eligible for screening. The timely participation rate was about 66% between 2001-2006, increased to 75% between 2007-2013, and started to decrease in 2014 to 69% in 2019. The delayed participation rate was approximately 5% between 2001-2013 and in 2014 began to increase to 8% in 2017. In the block 2001-2005, delayed participants were more likely to be 30-35 years old (OR,95%CI: 7.1, 5.8-8.6), be highly educated (OR,95%CI: 1.8, 1.6-1.9), and be pregnant during the invitation year (OR,95%CI: 9.5, 8.5-10.6) compared to timely participants. Similar results for age and education were found for 2006-2010, 2011-2015, and 2016-2020.

Conclusions: In Lifelines, the delayed participation in the CCS program has increased in the last five years. Considering both, timely and delayed, the participation was over 70% from 2001 to 2019. Delayed participation is associated with an age between 30-35, higher education, and pregnancy. These results might not be generalizable to the national data, and further analysis must be done to understand the low participation in recent years.

References: 1. Cervical cancer screening programme [Internet]. Vol. 2020. RIVM Committed to health and sustainability; 2020. Available from: <https://www.rivm.nl/en/cervical-cancer-screening-programme> 2. Framework for the Execution of Cervical Cancer Population Screening | RIVM [Internet]. [cited 2021 Jul 16]. Available from: <https://www.rivm.nl/documenten/framework-for-execution-of-cervical-cancer-population-screening> 3. Aitken, et al. Introduction of primary screening using high-risk HPV DNA detection in the Dutch cervical cancer screening programme: a population-based cohort study. BMC Med. 2019;17(1):228. 4. Monitor bevolkingsonderzoek baarmoederhalskanker 2019. Integraal Kankercentrum Nederland; 2019.

Poster

Mr Zupancic Mark
mark.zupancic@ki.se - Sweden

Mr Zupancic Mark
mark.zupancic@ki.se - Sweden

#3715

Long-Term Survival and Recurrence in Oropharyngeal Squamous Cell Carcinoma in Relation to Subsites, HPV, and p16-Status

30 - HPV and oropharynx / Head and neck cancer

Wendt M, Hammarstedt-nordenvall L, Friesland S, Landin D, Munck-wikland E, Dalianis T, Näsman A, Marklund L

Background/Objectives: Long-term survival data in relation to subsites, human papillomavirus (HPV), and p16INK4a (p16) for patients with oropharyngeal squamous cell carcinoma (OPSCC) is still sparsely studied. P16 alone may be suboptimal surrogate marker for assessing an active HPV infection in tumour tissue. It has been suggested that p16- in combination with HPV DNA-status could predict clinical outcome better than p16 alone in OPSCC. Furthermore, reports have indicated atypical and late recurrences for patients with HPV and p16 positive OPSCC. Therefore, we assessed long-term survival and recurrence in relation to oropharyngeal subsite and HPV/p16-status.

Methods: A total of 529 patients with OPSCC, diagnosed in the period 2000-2010 in Stockholm and Gotland Region, treated with curative intent, with known HPV DNA- and p16-status, were included. HPV/p16-status and subsites were correlated to disease-free and overall survival (DFS and OS respectively).

Results: The overexpression of p16 (p16+) is associated with significantly better long-term DFS and OS in tonsillar and base of tongue squamous cell carcinomas (TSCC/BOTSCC), but not in patients with other OPSCC. Patients with HPV DNA+/p16+TSCC/BOTSCC presented better DFS and OS compared to those with HPV DNA-/p16- tumours, while those with HPV DNA-/p16+ cancer had an intermediate survival. Late recurrences were rare, and significantly more frequent in patients with p16- tumours and other OPSCC, while the prognosis after relapse was poor independent of HPV DNA +/- p16 +/- status and OPSCC subsite.

Conclusions: In conclusion, patients with p16+ OPSCC do not have more late recurrences than p16-, and a clear prognostic value of p16+ was only observed in TSCC/BOTSCC. Finally, the combination of HPV DNA and p16 provided superior prognostic information compared to p16 alone in TSCC/BOTSCC.

Poster

Ms Hendrickx Jhana
jhana.hendrickx@novosanis.com - Belgium

Pasmans Danielle
danielle.pasmans@novosanis.com - Belgium

#3718

FIRST-VOID URINE AND VAGINAL SELF-SAMPLING IN CERVICAL CANCER SCREENING: FEEDBACK FROM A COLPOSCOPY REFERRAL POPULATION

13 - Self-sampling

Hendrickx J, Jordaens S, Meers N, Pasmans D, Beyers K, Vankerckhoven V

Background/Objectives: Cervical cancer (CC) is the seventh most common cancer in Europe. Almost all cases (99%) of CC are caused by oncogenic infections with high-risk human papillomavirus (hrHPV) strains. Many efforts have been made in several countries on hrHPV CC-related screening programs, although there is a great disparity between high- and low-income countries. First-void urine (FVU) is being more often positioned as the golden sample for HPV detection because it is non-invasive, has the ability to reach non-attendees of screening programs, and can overcome cultural barriers. The aim of the study was to evaluate sample preference as part of several studies in a colposcopy referral population.

Methods: Questionnaire data from the Predictors 5.1 study (United Kingdom)¹, the VALHUDES study (Belgium, NCT03064087)² and the Cocoss trial (Germany)³ were analyzed. The main goal of all studies was to investigate the clinical accuracy of hrHPV testing on FVU (Colli-Pee®) and vaginal self-sample (VSS) in a colposcopy referral populations. All participants were asked to fill out a questionnaire on their general experience and convenience on all different sampling methods. For comparison matters, only dry vaginal SS were compared between studies. Data are presented as mean \pm SEM.

Results: In total, 1034 women aged 21 - 76 years consented their participation and provided all sample types. Self-collection was well accepted, with $48 \pm 4\%$ of women indicating a preference for FVU over $39 \pm 4\%$ who preferred a vaginal SS. On average, $77 \pm 9\%$ of women indicated that FVU was "very easy to use" compared to $73 \pm 9\%$ for VSS, and $76 \pm 8\%$ and $60 \pm 1\%$ of women indicated that they were confident that samples were taken correctly for FVU and VSS, respectively.

Conclusions: Here, we show that the majority of women enrolled in several European triage population preferred FVU over a vaginal SS, found that FVU was more easy-to-use and that they were confident that samples were taken correctly. This outcome highlights the potential of FVU to become the golden sample for detection of HPV infection and offers future perspective to screen non-attendees in low-income countries and regions where VSS is less well accepted.

References: 1Cadman, L., et. al., Cancer Epidemiology and Prevention Biomarkers (2021) 2De Pauw, H., et. al., Archives of Public Health (2021) 3Ertik, F. C., et. al., International journal of environmental research and public health (2021)

Oral presentation

Dr Schmitz Martina
martina.schmitz@oncgnostics.com - Germany

Schmitz Martina
martina.schmitz@oncgnostics.com - Germany

#3720

GynTect DNA methylation marker - longitudinal observational study in patients with CIN2/3. Results from the GynTect-PRO trial

18 - Methylation

Schmitz M, Wunsch K, Hansel A, Hippe J, Heller T, Hoyer H, Dürst M

Background/Objectives: Precancerous lesions of the cervix, depending on their grade of severity, are known to have a high potential to regress to normal. Especially among young women, only a few patients are at high risk to develop invasive cancer. Up to now, current methods are not able to distinguish between high risk patients and those patients who are likely to show regression of their precancerous lesions. DNA methylation markers such as the CpG islands in the promoter regions of the genes ASTN1, DLX1, ITGA4, RXFP3, SOX17, ZNF671 comprising the diagnostic test GynTect® are a new class of biomarkers which are discussed to have prognostic potential. GynTect-PRO is a prospective, longitudinal and multi-centered trial in Germany aiming to address the probability of regression in GynTect-negatively tested women having a CIN2 or CIN3 lesion (Negative Predictive Value, NPV). The study tests the null hypothesis $NPV \leq 70\%$ against the alternative hypothesis $NPV \geq 90\%$.

Methods: 77 women <25 years were included between December 2017 and February 2021, recruited from nine clinical centers in Germany. All patients had a histopathology confirmed CIN2 or CIN3 lesion at first visit. Follow-ups were planned for all patients every 6 months. CIN2 patients were observed for up to 24 months (a max. of 5 visits) and CIN3 patients for up to 12 months (a max. of 3 visits). At all visits, colposcopy was done and, if indicated, a biopsy was taken. HPV testing, cytology and GynTect testing was also done at every timepoint. The outcome was defined as progression, persistence or regression with respect to the last biopsy, conization or colposcopy findings if no histopathology was done at the last visit. In the end, 60 women fulfilled all inclusion criteria.

Results: In total, 18 of 24 CIN2 and 27 of 36 CIN3 patients had a negative GynTect result at the study entry point. Over time, 12 of 18 GynTect negative CIN2 patients showed regression ($NPV=67\%$, $90\%CI$ 44%-85%), whereas 5 persisted during the follow-up period (2 of them with follow-up <1.5 years). One patient showed progression. Of the 27 GynTect negative CIN3 lesions 15 regressed ($NPV=56\%$, $90\%CI$ 38%-72%) and 12 showed persistence (5 of them with follow-up <0.8 years). 4 CIN2 and 6 CIN3 were tested GynTect positive and 2 CIN2 and 3 CIN3 showed an invalid result at entry point.

Conclusions: The objective of the study to show a negative predictive value of $\geq 90\%$ for a GynTect negative result was missed. Since follow-up of some patients was shortened the probability of a final regression could be underestimated and that of a final persistence overestimated. On the other hand, regression of a lesion may be induced by repeated biopsies taken for safety reasons during follow-up. For valid NPV estimates we need to study the natural development of CIN2/3 lesions over a relevant period, a task which is difficult to implement in the target clinical setting.

Poster

Ms Hendrickx Jhana
jhana.hendrickx@novosanis.com - Belgium

Vankerckhoven Vanessa
vanessa.vankerckhoven@novosanis.com - Belgium

#3722

HOME-BASED SELF-SAMPLING OF FIRST-VOID URINE FOR HRHPV TESTING IN CERVICAL CANCER SCREENING: USABILITY FEEDBACK FROM A BELGIAN COLPOSCOPY REFERRAL POPULATION

13 - Self-sampling

Hendrickx J, Jordaens S, Meers N, Rios Cortes A, Beyers K, Vankerckhoven V

Background/Objectives: Cervical cancer (CC) is the seventh most common cancer in Europe with 99% of cases caused by oncogenic infections with high-risk human papillomavirus (hrHPV) strains. Women with socio-economical disadvantages and lower education are often under-screened and hard-to-reach leading to an increased risk of developing CC. Successful CC screening programs strongly depend on the participation of the target population. Various barriers are known to contribute to lower participation such as physical discomfort, poor access to health services, time constraints and lack of knowledge. Here we report interim usability feedback results from the CASUS study (NCT04530201, Belgium) using first-void urine (FVU) as a liquid biopsy.

Methods: Women were asked to self-collect two FVU samples at home with the Colli-Pee® Small Volumes (10mL) variant (Novosanis, Belgium), prefilled with a non-toxic nucleic acid preservative the day prior to their colposcopy appointment. Afterwards, women completed a questionnaire indicating their usability feedback of the Colli-Pee® Small Volumes urine self-sampler. A specific Systems Usability Scale (SUS) score was incorporated in the questionnaires. A SUS score greater than 68 is considered good. Data are presented in relative percentages and as mean \pm SEM.

Results: Up until the beginning of September 2021, a total of 280 women (25-64y) were enrolled and consented their participation in the CASUS study. The interim analysis includes data of 204 questionnaires with 197 valid SUS scores. Overall, 57% of women indicated to prefer FVU self-sampling over a physician taken PAP smear (37%) for their next CC screening. Additionally, 61% women indicated to prefer the use of Colli-Pee® Small Volumes over a urine cup whereby 95% of women experienced Colli-Pee® Small Volumes as easy to use and 97% who would use the device again. An average SUS score of 85.49 ± 15.60 was calculated.

Conclusions: The results of this study show that the majority of women would prefer a urine self-sample at home over a physician-taken PAP smear for their next CC screening. Moreover, Colli-Pee® Small Volumes was considered an easy-to-use and well-accepted self-sampling device for CC screening in a Belgian colposcopy referral population. From a future perspective, these results highlight the possibility of home-based urine self-sampling as a liquid biopsy in CC screening where hard-to-reach populations could be approached more easily.

Oral presentation

Wiik Johanna
johanna.wiik@gmail.com - Norway

Ms Wiik Johanna
johanna.wiik@gmail.com - Sweden

#3723

MATERNAL HUMAN PAPILLOMAVIRUS INFECTION DURING PREGNANCY AND PRETERM DELIVERY, A MOTHER-CHILD COHORT STUDY IN NORWAY AND SWEDEN

03 - Epidemiology and natural history

Wiik J, Værnesbranden M, Jonassen C, Staff A, Lødrup Carlsen K, Hilde K, Sundet B, Skjerven H, Rehbinder E, Nordlund B, Vettukttil R, Granum B, Haguen G, Hedlin G, Söderhäll C, Jacobsson B, Nilsson S, Sengpiel V, Sjöborg K

Background/Objectives: Human papillomavirus (HPV) is a common infection in reproductive age. Infection, inflammation and placental dysfunction are leading causes for preterm delivery. This study aimed to assess the association between HPV infection in pregnancy and preterm delivery (PTD), prelabour rupture of membranes (PROM), preterm prelabour rupture of membranes (pPROM), chorioamnionitis and neonatal sepsis.

Methods: Pregnant women participating in a prospective multicenter cohort study of a general population in Norway and Sweden (PreventADALL, ClinicalTrials.gov, NCT02449850) provided samples for HPV-DNA detection in first void urine at mid-gestation and at delivery, and from placenta after delivery. The association between HPV-DNA presence; any HPV (28 genotypes), any high risk HPV (12 HR-HPV) or HPV 16 and PTD, pPROM, PROM, chorioamnionitis and neonatal sepsis was analyzed using logistic regression adjusted for relevant confounders. In women positive for HR-HPV at mid-gestation, we examined if persistence of HPV, viral load or presence of multiple HPV infection were associated with obstetric outcomes.

Results: The study included 950 women (mean age 32 years) with singleton pregnancies and valid urine HPV test at mid-gestation and obstetric outcomes, whereof 753 also had a valid urine HPV test at delivery. At mid-gestation, 24% were positive for any HR-HPV and 40% for any HPV. The frequency of PTD was higher in HR-HPV positive (3.5%) than in HR-HPV negative women (2.1%) at mid-gestation but the association was non-significant (adjusted odds ratio 1.71 (95% CI 0.70-4.19), $p=0.24$). We found no significant association between presence or persistence of HR-HPV or HPV at mid-gestation or delivery and PTD, pPROM, PROM, chorioamnionitis or neonatal sepsis. HPV-DNA was not detected in any of the placentas in women with PTD, pPROM or chorioamnionitis.

Conclusions: HPV infection during pregnancy was not associated with PTD, pPROM, PROM, chorioamnionitis or neonatal sepsis in this general population with a low incidence of adverse obstetric outcomes.

Oral presentation

Dr Martinelli Marianna
marianna.martinelli@unimib.it - Italy

Dr Martinelli Marianna
marianna.martinelli@unimib.it - Italy

#3724

HPV TESTING OF SELF-COLLECTED FIRST-VOID URINE SAMPLES: COMPARISON OF PERFORMANCE USING DEVICES COLLECTING DIFFERENT URINE VOLUMES

13 - Self-sampling

Giubbi C, Perdoni F, Musumeci R, Monticciolo I, Di Meo M, Fruscio R, Landoni F, Cocuzza C

Background/Objectives: HPV testing of self-collected samples has been demonstrated to be a valid alternative to reach women reluctant to undergo clinician-based cervical cancer screening. First-void urine (FVU) has been recently proposed to be a less invasive sample with a similar clinical accuracy to cervical sample in detecting high-grade cervical intraepithelial neoplasia. Colli-Pee® device allows standardized collection of different volumes of FVU captured in a collector tube prefilled with a preservation solution. The aim of the study was to evaluate the accuracy of HPV testing on urine self-samples using two different devices able to collect different volumes of FVU (20ml and 10ml) as compared to cervical samples (gold standard) in women referred to colposcopy.

Methods: Clinician-collected cervical (L-shaped Endo/Esocervical FloqSwab®, Copan) and FVU samples (Colli-Pee®, Novosanis), were obtained from 90 women attending the Colposcopy Clinic, San Gerardo Hospital (Monza, Italy). Two different devices using volume variants have been tested: 45 women collected FVU using the 10mL and 45 women using the 20mL Colli-Pee® device. All samples were extracted using MicroLab Nimbus starting from 200 µl of sample and HPV detection carried out using AnyplexII HPV28 (Seegene). Concordance in HPV detection between sample types was determined using Cohen's Kappa value (k).

Results: 44% (20/45) and 51% (23/45) of women that used respectively the 10ml and the 20ml Colli-Pee® devices resulted positive in clinician-collected cervical sample for one or more hrHPV infections. A good concordance for hrHPV was observed between clinician-collected cervical and urine samples collected using both devices (82.2% total agreements for 10mL and 20mL, respectively). In particular HPV16 and HPV18, responsible for over 70% of cervical cancer cases, showed high agreement rates between cervical and urine samples collected using both devices (10 mL Colli-Pee®: HPV16: k= 0.897 HPV18 k=1.000; 20 mL Colli-Pee®: HPV16: k=1.000 HPV18: k=1.000).

Conclusions: These preliminary data demonstrate promising results for the use of both urine collecting devices in cervical cancer screening. The high concordance rate for hrHPV detection, including the lower volume collection device, if confirmed by larger number of tested samples, would suggest the possibility and feasibility of performing FVU home-collection in screening programs, allowing increased and safer participation of women to cervical cancer screening even during COVID-19 pandemic.

Oral presentation

Ms Hendrickx Jhana
jhana.hendrickx@novosanis.com - Belgium

Hendrickx Jhana O.
jhana.hendrickx@novosanis.com - Belgium

#3725

FIRST-VOID URINE: A RELIABLE APPROACH FOR CERVICAL CANCER SCREENING

10 - HPV screening

Hendrickx J, Jordaens S, Meers N, Rios Cortes A, Pasmans D, Beyers K, Vankerckhoven V

Background/Objectives: A global analysis of cervical cancer (CC) incidence and mortality in 2018 revealed 570 000 cases of cervical cancer and 311 000 deaths from the disease, making it the fourth most common cancer in the world.^{1,2} CC is primarily caused by infections with high-risk strains of the human papillomavirus (hrHPV). The golden standard to test for hrHPV infections is a physician taken smear (PTS). However, this screening method has a suboptimal attendance rate due to reluctance, discomfort and cultural barriers. The Colli-Pee® device by Novosanis is user-friendly allowing for the collection and stabilization of first-void (FV) urine to enable self-sampling for the purpose of HPV screening. The aim of this abstract is to provide consolidated data on the clinical performance of the Colli-Pee® device compared to PTS from several studies that explored the detection of HPV in self-samples.

Methods: Novosanis developed the Colli-Pee® portfolio for standardized and volumetric (10mL, 20mL) self-sampling (SS) of FV urine. Additionally, Colli-Pee® can be prefilled with Novosanis' UCM, to allow preservation and stabilization of HPV DNA in urine. Colli-Pee® containing UCM is CE-IVD marked, and has been registered in several countries worldwide. In total, eight clinical trials have been reviewed to assess the potential of Colli-Pee® -collected FV urine compared to PTS and different vaginal SS devices for the detection of HPV.

Results: A total of 2443 women were enrolled and consented their participation in the trials. Seventeen different institutions are involved covering multiple European regions and the state of New Jersey, USA. Overall, 9 different HPV assays were used to investigate the clinical performance of Colli-Pee® against a PTS and nine different vaginal SS devices. Throughout, high overall HPV positivity was detected with Colli-Pee® collected FV urine on high-throughput HPV assays and the concordance for hrHPV detection between FV urine and cervical samples ranged from good to very good.

Conclusions: The clinical performance of FV urine for hrHPV detection is non inferior to that PTS as tested via multiple HPV high-throughput assays. Thus, FV urine can be considered an appropriate and user-friendly sample type for CC screening that can improve attendance rates because it is painless, can be used in a home setting, and does not create cultural barriers making it easier to target hard-to-reach and under-screened populations. Furthermore, new generation of Colli-Pee® devices are being evaluated in the framework of the CASUS study for the collection and stabilization of smaller volumes and the compatibility of collection tubes with high-throughput machines in favor of CC screening.

References: 1 Arbyn, M., Weiderpass, E., Bruni, L., de Sanjosé, S., Saraiya, M., Ferlay, J., & Bray, F. (2020). Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *The Lancet Global Health*, 8(2), e191-e203. 2 World Health Organization (WHO), Cancer Today, available on: <https://gco.iarc.fr/today>, accessed on: December 13, 2021

Oral presentation

Mrs Tian Pu
19111250016@fudan.edu.cn - China

Tian Pu
19111250016@fudan.edu.cn - China

#3726

Mechanism of HPV16 E2 promoting its persistent infection by inhibiting cGAS-STING signaling pathway

05 - Immunology

Background/Objectives: HPV employs multiples strategies to escape immune surveillance. And one important approach is to inhibit DNA sensing pathway in the host cells. Previous studies have confirmed that E6/E7 can inhibit the activation of cGAS-STING signal pathway. However, how HPV can escape immune recognition during its persistent infection without gene integration is unclear. We hypothesized that HPV E2 may impair innate immune recognition of HPV and promot persistent infection.

Methods: To confirm if cGAS-STING signal Pathway can be activate by HPV, HPV16PsV was applied to infect HaCaT and the activation of STING was measured by Western-blot and Immunofluorescence. And we overexpressed E2 in HaCaT by trasfecting HPV16 E2 plasmids to observe whether E2 has an effect on the activation of cGAS-STING signal pathway. Quantitative PCR and western blotting were performed to determine the RNA and protein expressions of STING and its downsream cytokines, respectively.

Results: STING was activated and its expression increased in HaCaT after being infected by HPV16PsV. Compared with the control group, the expression of STING, IRF3 and ISG15 significantly decreased in HaCaT overexpressed HPV16 E2.

Conclusions: HPV16 impairs innate immune detection via reducing STING expression in host cells during its persistent infection without gene integration.

References: [1] Luo X, Donnelly CR, Gong W, et al. HPV16 drives cancer immune escape via NLRX1-mediated degradation of STING. J Clin Invest, 2020, 130(4):1635-1652. [2] Lau L, Gray EE, Brunette RL, Stetson DB. DNA tumor virus oncogenes antagonize the cGAS-STING DNA-sensing pathway. Science. 2015;350(6260):568-571.

Poster

Ms Lujan Elena
lujan.e@procarehealth.com - Spain

Mr Gaslain Yann
gaslain.y@procarehealth.com - Spain

#3727

EFFICACY OF A MULTI-INGREDIENT CORIOLUS VERSICOLOR-BASED VAGINAL GEL IN HIGH-RISK HPV INFECTED PATIENTS: RESULTS OF 6 DIFFERENT STUDIES

26 - Cervical neoplasia

Cortés J, Dexeus D, Santiago P, Serrano L, Gajino C, Marín E, Criscuolo A, Riera M

Background/Objectives: To evaluate the consistency of the efficacy of a multi-ingredient Coriolus versicolor-based vaginal gel, Papilocare®, on high-risk human papillomavirus (HR-HPV) clearance in 6 different studies.

Methods: Results from 4 independent observational studies (6 month-treatment period with Papilocare®) were compared to results from a randomized, open, parallel, controlled trial (Paloma: NCT04002154) and an observational, multicenter, prospective, one-cohort study (PapilOBS: NCT04199260). Vigo study: Prospective one-cohort. Secondary endpoint (SE), HPV clearance in 25 patients infected by HPV 16 and/or 18. Coruña study: Retrospective one-cohort. Primary endpoint (PE), HPV clearance assessed in 57 medical patients' records. Hospitalet study: Retrospective one-cohort. PE, Composite efficacy variable (patients with normal cytology and/or HPV clearance) in 91 HR-HPV patients. Roma study: Retrospective controlled. PE, HR-HPV clearance in 183 patients. Paloma trial: SE, HR-HPV clearance in 65 patients. PapilOBS study: SE, HR-HPV clearance in 176 patients.

Results: 48% of patients cleared 16-/18 -HPV in Vigo study. 58% of reduction was observed in the number of HR-HPV patients (Coruña) and 72.5% normalized cytology and/or cleared HR-HPV (Hospitalet). 67% HR-HPV clearance was observed (treated group) vs 37.2% (control group), in the Roma study. In the Paloma trial, HR-HPV clearance reached 63% (treated group) vs 40% (control group). 57.4% HR-HPV clearance was observed in the PapilOBS study.

Conclusions: Papilocare® has shown significant consistent rates of efficacy with a 64% of HR-HPV clearance in average in 6 different studies involving 597 patients which reinforce its beneficial effect for HR-HPV patients.

Results in different studies

Poster

Mrs Giubbi Chiara
chiara.giubbi@unimib.it - Italy

Mrs Giubbi Chiara
chiara.giubbi@unimib.it - Italy

#3731

IMPACT OF VAGINAL SELF-SAMPLES RESUSPENSION VOLUME ON HUMAN PAPILLOMAVIRUS (HPV) TESTING

13 - Self-sampling

Giubbi C, Monticciolo I, Di Meo M, Perdoni F, Musumeci R, Castriciano S, Fruscio R, Landoni F, Cocuzza C

Background/Objectives: Validation of Human Papillomavirus (HPV) testing in combination with self-collection devices offers the possibility to improve cervical cancer (CC) screening coverage rates reaching women who do not attend screening programs. Moreover, the possibility of performing HPV testing on vaginal self-samples eluted in alternative media and resuspension volumes as compared to the methanol-based media, presently used routinely to perform cervical cytology (Pap Test) for the triage of women HPV-positive, may further reduce screening costs. This ongoing study aims to evaluate the performance of HPV detection on self-collected vaginal samples resuspended in two different non-alcohol-based media, eNat® and MSwab® (Copan), using two different resuspension volumes (2 ml and 5 ml).

Methods: Two self-collected vaginal samples (FLOQSwab®, Copan) and a clinician-taken cervical specimen (L-Shaped FLOQSwab®, Copan) were collected from 17 women referred to colposcopy at San Gerardo Hospital, ASST Monza. Cervical swabs were resuspended in 20 ml of ThinPrep®PreservCyt® (Hologic®), while vaginal samples were transported dry to the laboratory. In the laboratory, vaginal samples from 10 women were resuspended in 2 ml and 5 ml of eNat® respectively; vaginal samples from the remaining 7 women in 2 ml and 5 ml of MSwab®. Nucleic acids were extracted from 200 µL using Microlab Nimbus platform (Seegene) and HPV detection carried out with AnyplexTMII HR-HPV real-time assay (Seegene). HPV test results obtained from the 2 vaginal specimens/woman were then compared to those obtained from the cervical sample.

Results: Preliminary results showed HR-HPV positivity rates of 70.6% (12/17), 76.5% (13/17) and 64.7% (11/17) of women respectively in cervical swabs, vaginal self-samples resuspended in 2 ml and in 5 ml. Independently from the suspension medium used, the overall agreement rate between cervical and vaginal specimens resuspended in a volume of 2 ml was 94.12% (16/17), while the percentage of agreement between cervical and vaginal swabs resuspended in 5 ml was 82.35% (14/17).

Conclusions: These preliminary data demonstrated a good agreement in HPV detection between cervical and vaginal self-collected samples independently from the resuspension medium and/or volume used, supporting the possibility of introducing alternative safer and more cost-effective media for the implementation of vaginal self-sampling in CC screening programs. Studies on a larger set of samples could confirm these findings.

Oral presentation

Dr Armaroli Paola
paola.armaroli@cpo.it - Italy

Dr Armaroli Paola
paola.armaroli@cpo.it - Italy

#3733

Results after two screening rounds for primary HPV screening for cervical cancer, in a population based organized screening program in Piedmont, Italy

10 - HPV screening

Rizzolo R, Ronco G, Gillio-tos A, De Marco L, Cristina S, Valentini S, Larato C, Massari L, Leo L, Giordano L

Background/Objectives: In Piedmont Region, Italy, cervical cancer screening has been conducted with a primary high-risk human papillomavirus (hrHPV) test and cytology triage since February 2014. We here compare HPV positivity, colposcopy referral, and detection of cervical intraepithelial neoplasia (CIN) 2+ at first and at second screening round.

Methods: Within the organised population based screening programme, women of 30-64 years of age were randomised for invitation to a first screening round with hrHPV test in 2014-2015. They were invited for a second screening round in 2019-2020. The primary hrHPV test used was Hybrid Capture 2. hrHPV-negative women were scheduled for re-screening after 5 years. Women with a positive hrHPV (HPV+) test and cytology triage \geq than ASC-US or unsatisfactory were directly referred to colposcopy, whereas HPV+ women with a negative cytology triage were re-invited for a hrHPV test after 1 year. If retesting was still positive they were referred to colposcopy, otherwise were scheduled for re-screening after 5 years

Results: In 2019-2020, 44162 women aged 30-64 were invited for the second screening round with hrHPV test, and 28552 participated (participation rate 65% vs 55% at first round, Relative Risk -RR- 1.17, 95%IC 1.15-1.18). At the second round, the observed positivity rate was 5.7%, compared with 7.6% at the first one (RR 0.74, 95%IC 0.70-0.78), and 344 women were immediately referred to colposcopy (1.2% compared with 1.5% at first round, RR 0.82, 95%IC 0.72-0.92). Compliance to colposcopy was 92%, not different from the one observed at the first round (93%). Immediate detection of CIN2+ lesions was 1.7‰, compared with 4.1‰ at the first round (RR 0.41, 95%IC 0.30-0.56). At hrHPV test repetition after 1 year, at second round 37% had a HPV+ result and were referred to colposcopy, compared to 53% at the first one (RR 0.70, 95%IC 0.60-0.81), but only the 40% (501) of the women negative at cytological triage could perform a hrHPV test after 1 year, due to the reduction of the screening activity during the COVID-19 emergency.

Conclusions: Among women at second screening round we observed a significantly higher participation, and significant reduction in the positivity rate (26%), in the immediate colposcopy referral rate (21%), and in the immediate CIN2+ detection (59%). Due to the COVID-19 emergency, data on HPV retesting at 1 year are not conclusive, but preliminary results would suggest an overall reduction in colposcopy referral and detection of CIN2+ lesions

Oral presentation

Dr Biederman Erika
emcconne@iu.edu - United States

Biederman Erika
biederman.51@osu.edu -

#3742

FACTORS ASSOCIATED WITH MAILED RETURN OF SELF-COLLECTION FOR HUMAN PAPILLOMAVIRUS (HPV) TESTING

13 - Self-sampling

Zimet G, Champion V

Background/Objectives: Low-income women in the United States (U.S.) have greater odds of dying from cervical cancer and lower cervical cancer screening rates compared to higher income women. Self-collection for human papillomavirus (HPV) testing has demonstrated acceptability in previous studies, but up to 50% of low-income women do not return self-collection kits, and characteristics of those who complete and return a self-collection kit have not been investigated. The purpose of the current study was to examine how sociodemographic factors (age, education, income, insurance and marital status, and race), health-related variables (seeing a provider in the past year, body mass index, smoking status, history of abnormal Papanicolaou test), and relative advantages were associated with mailed return of self-collection kits (returners vs. non-returners).

Methods: Low-income women (n=168) aged 30-65 with no history of hysterectomy who were not up to date with U.S. Preventive Services Task Force cervical cancer screening guidelines (i.e., no Papanicolaou [Pap] test in the past 3 years, HPV test in the past 5 years, or Pap and HPV test in the past 5 years) were recruited from Midwestern food pantries and online (Craigslist and Facebook). Women completed a survey and were mailed self-collection kits that could be mailed back. Bivariate analyses were initially conducted, with significant bivariate predictors subsequently entered in a multivariable logistic regression analysis to evaluate predictors of mailed return of self-collection kits.

Results: In bivariate analyses, age, education, body mass index, and relative advantages were significantly different between returners and non-returners and thus entered the multivariable logistic regression. Women in the 40-49 age-group vs. the 30-39 age-group (Odds ratio [OR]=3.32, 95% Confidence Interval [95%CI]=1.44,7.69), college education or more vs. high school education or less (OR=6.47, 95%CI=1.61,25.96), obese vs. non-obese (OR=2.24, 95%CI=1.06,4.72), and with a higher relative advantages score (OR=1.09, 95%CI=1.01,1.20), had greater odds of returning a kit.

Conclusions: Interventionists could use results from the current study to target women who are not in the 40-49 age group, college educated, obese, or who have a higher relative advantages score with an intervention to increase self-collection. Future studies should examine these results in a larger, more diverse sample.

Oral presentation

Tota Joseph
joseph.tota@merck.com - United States

Dr Palmer Cody
cody.palmer@merck.com - United States

#3743

Forecast Reduction in RRP in the US Attributable to Quadrivalent and Nonavalent HPV Vaccination: A Dynamic Transmission Modeling Study

36 - Economics and modelling

Palmer C, Tota J

Background/Objectives: Recurrent Respiratory Papillomatosis (RRP) is a rare life-threatening disease resulting from HPV infection in both children (Juvenile Onset RRP - JORRP) and adults (Adult Onset RRP - AORRP) with no cure. HPV infections that result in AORRP are generally contracted through sexual activity, while HPV infections in children are likely a result of vertical transmission through an infected mother. 95% of cases of RRP are thought to be caused by HPV types 6 and 11, which are both targeted by available quadrivalent (4vHPV) and nonavalent (9vHPV) vaccines. Using dynamic transmission disease modeling, we investigate how HPV vaccination has driven the observed decline of JORRP in the US, and the impact that it will have on future RRP incidence.

Methods: A dynamic transmission model for HPV infection and RRP disease was adapted to the US, utilizing pre-vaccination data on HPV prevalence and seroprevalence in US adults, birth cohort incidence of JORRP, age of diagnosis of JORRP, and incidence of AORRP. Rollout of 4vHPV and 9vHPV vaccination, accounting for known vaccine coverage rates since 2007, was simulated and projected forward. We also ran some counter-factual historical vaccination scenarios to investigate how vaccination in older adolescent age groups impacted JORRP decline. Finally, health economic results were collected, including costs-averted (2020 USD; discounted 3%) and QALYs gained compared to no vaccination.

Results: The model accurately reproduces the documented decline in JORRP among birth cohorts since 2007. Forecasting RRP incidence suggests that after 2030, we can expect to see 10 cases of JORRP per year or less among children under the age of 13. For AORRP, after 2040 virtually all new cases of AORRP are due to types not covered by the available vaccines. Averted treatment costs exceed 200 million USD over the time horizon, and QALYs gained were significant. We also show that 75% of the observed decline in incidence of JORRP since 2007 is due to vaccination of individuals aged 16 years and older.

Conclusions: Our modeling suggests that 4vHPV and 9vHPV vaccination has contributed to the documented decline of JORRP incidence by averting infection among adults and is providing protection for infants and children under 13 years of age. We have shown that vaccination of adults and older adolescents would be a key driver of rapid drops in RRP incidence in the early years of a national HPV vaccination program. Lastly, modeling suggests that, if current vaccination practice continues, after 2040 HPV6/11-related RRP will be eliminated, both among adults and children.

Oral presentation

Mrs Sonne Si Brask
si.brask.sonne@regionh.dk - Denmark

Sonne Si Brask
si.brask.sonne@regionh.dk - Denmark

#3745

A RANDOMIZED, LARGE SCALE COMPARISON OF DIRECT MAIL VERSUS OPT-IN HPV SELF-SAMPLING INVITATION STRATEGY: PARTICIPATION RATES, PARTICIPANTS, FOLLOW-UP AND COST OF LAST ATTENDER

13 - Self-sampling

Sonne S, Pedersen B, Andreasen E, Ejegod D, Pedersen H, Bonde J

Background/Objectives: Background HPV self-sampling is increasingly being offered as an alternative to ordinary screening. However, which invitation strategy yields the best cost-benefit in terms of participation versus number of self-sampling kits distributed. Is it directly mailing kits to all eligible women or an Opt-in strategy where kits are mailed only to those who actively accept the offer? In a large scale, randomized, study we evaluated participation and number of HPV self-sampling kits distributed by a Direct mail and Opt-in approach.

Methods: Methods Women with no registered cervical screening sample in at least 4 years were randomly assigned to either the Direct mail group (N=9,634, period June-Sept 2020) or the Opt-in group (N=39,754, period Sept 2020-January 2021). In the Direct mail group, all women received the HPV self-sample kit whereas the Opt-in group received the HPV self-sampling invitation by national e-mail with instructions for registration. Reminder letters to return the HPV self-sampling kit for analysis were sent 60 days after shipping the kit. For the Opt-in arm, only women who registered for a test kit received a reminder. Follow-up information was registered and retrieved from the Danish National Pathology Database (Patobank), latest data retrieval November 2021. The test kit contained an Evalyn self-sampling brush. Upon arrival in the lab, the sample was analyzed on a VIPER LT, using the Onclarity HPV assay from BD. Women found HPV positive by self-sampling were referred for follow up with a clinician collected liquid based cytology sample within 3 months.

Results: Results In the Direct-Mail arm, 25% (N=2,406) returned the self-sampling kit and in the Opt-in arm, 20% (7,994) returned the kit. In the Direct mail arm, 7,850 women received a reminder with 9% (N=685) responding. In the Opt-in arm, 7,462 received a reminder resulting in 30% (N=2,222) samples returned. The overall HR-HPV prevalence was 14.6%, and 89% of the HPV positive women adhered to the follow up recommendation, with no difference between the two arms, respectively. For each woman included in the Direct mail group, 3 tests were never returned, whereas 0.7 tests were lost for each participant from the Opt-in group. More women participated by direct mail, but at a cost of >13 HPV self-sample kits per additional woman compared to the Opt-in approach.

Conclusions: Conclusions Both invitation approaches returned a high number of participating women. A reminder strategy yielded a significant number of additional participants in both groups, and women in general adhered to the follow up recommendation. Finally, the cost of conducting a Direct mail approach seems rather high when balanced against the yield in participants.

Oral presentation

Dr Bottari Fabio
fabio.bottari@ieo.it - Italy

Bottari Fabio
fabio.bottari@ieo.it - Italy

#3748

THE EXTENDED VALIDATION OF HUMAN PAPILLOMAVIRUS ASSAYS AND COLLECTION DEVICES FOR HPV TESTING (EXTENDED VALHUDES): THE IEO EXPERIENCE.

13 - Self-sampling

Bottari F, Iacobone A, Passerini R, Pietri R, Piana A, Guerrieri M, Martella S, Preti E, Cocuzza C

Background/Objectives: Previous studies demonstrated that PCR-based HPV tests are similarly accurate on self-produced samples as on clinician-collected cervical samples [1] but validation of HPV assays in combination with the self-sample collection devices for are required [2]. This study, named Extended VALHUDES (Extended VALidation of HUMAN papillomavirus assays and collection DEvices for HPV testing on Self-samples), aims to extend a previous VALHUDES validation of BD Onclarity HPV Assay in combination with a new vaginal sample collection device, COPAN FLOQ Swab, on BD Viper LT automated instrument.

Methods: A total of 300 women referred to the European Institute of Oncology (IEO) in Milan and a Coordinamento Consultori Familiari, ASSL Sassari, ATS Sardegna because of a previous cervical abnormality were enrolled after obtaining informed consent. A dry vaginal self-sample was collected by each enrolled woman and transferred to BD HPV LBC Self-Collection diluent (3 mL) tubes on arrival at IEO laboratory and tested with BD Onclarity on the BD Viper LT at IEO. An aliquot of the clinician-collected cervical sample PreservCyt media was transferred to a BD HPV LBC diluent tube. Vaginal swab and the LBC cervical specimen were tested with BD Onclarity on the BD Viper LT at IEO to compare performance.

Results: 296 of the 300 women had evaluable results from both the vaginal self-sample and the clinician collected cervical specimen. 172 of the 296 women had cervical biopsies obtained during the colposcopy, with histopathology reports. There were 79 women with \geq CIN2 histology and 48 women \geq CIN3. The overall agreement between vaginal self-sample and cervical sample for BD Onclarity HPV Assay on BD Viper LT in IEO laboratory was 264/296 (89.2%; CI 79.1-99.3%), Kappa=0.75 (CI 0.630-0.867). The absolute sensitivity for the vaginal self-sample and cervical sample in CIN2+ detection was 89.9% and 88.6% respectively and the absolute specificity was 38.7% and 40.9% respectively.

Conclusions: This data demonstrates a good concordance and sensitivity between HPV detection on vaginal samples vs BD Onclarity HPV Assay (Viper LT) on clinician-collected samples.

References: [1] Arbyn M, Smith SB, Temin S, Sultana F, Castle P; Collaboration on Self-Sampling and HPV Testing. Detecting cervical precancer and reaching underscreened women by using HPV testing on self samples: updated meta-analyses. BMJ. 2018 Dec 5;363:k4823. [2] Arbyn M, Peeters E, Benoy I, Vanden Broeck D, Bogers J, De Sutter P, Donders G, Tjalma W, Weyers S, Cuschieri K, Poljak M, Bonde J, Cocuzza C, Zhao FH, Van Keer S, Vorsters A. VALHUDES: a protocol for VALidation of HUMAN papillomavirus assays and collection DEvices for HPV testing on Self-samples and urine samples. J Clin Virol. 2018; 117: 52-56

Poster

Dr Dickey Brittney
brittney.dickey@moffitt.org - United States

Mrs Giuliano Anna
Anna.Giuliano@moffitt.org - United States

#3753

A BIOMARKER PANEL FROM GENOME-WIDE METHYLATION TO PREDICT EARLY OROPHARYNGEAL CANCER

18 - Methylation

Dickey B, Putney R, Schell M, Chung C

Background/Objectives: As oropharyngeal cancer (OPC) associated with human papillomavirus (HPV) increases in men, the need for a screening test to diagnose OPC early is crucial. Prognosis is favorable though recurrence is likely and metastatic OPC is often incurable regardless of HPV status. We previously reported that methylation of the host tumor suppressor gene, EPB41L3, and the HPV16 genome assessed from oral gargles in pretreatment, male OPC cases (n=228) was significantly ($p<0.0001$) higher among cases compared to controls and predicted cases with an AUC of 0.82. However, prediction in early OPC cases alone (T1-T2, N0-N1; small tumors with a single ipsilateral node $<3\text{cm}$) was lower (AUC=0.78), indicating the need for additional biomarkers to improve screening performance.

Methods: Therefore, this study agnostically identified oral cell CpG sites differentially methylated among early OPC cases. DNA was extracted from oral gargles of 228 OPC cases (92 early cases) and 142 frequency matched healthy controls and processed for genome-wide methylation using the MethylationEPIC BeadChip (Illumina) array. Ten fold cross validation was used to independently compare detection at individual CpG sites and select sites for modeling. Selected sites were combined with our prior methylation data, which included three CpG sites (438, 427 and 425) in the EPB41L3 gene. Due to strong association with cancer, oral HPV16 status was considered as a binary variable (positive/negative). Lasso regression identified CpG sites strongly associated with early OPC. Receiver operator characteristic (ROC) curves with area under the curve (AUC) and optimized sensitivity and specificity cut points were generated. As a final step, the panel was validated utilizing bootstrap re-sampling.

Results: Analyses revealed 14 markers significantly associated with early OPC, including one EPB41L3 CpG site (438) and oral HPV16 status. Together this panel trained the model and was able to predict early OPC compared to controls with an AUC of 0.970. In the bootstrap validation set, the AUC was 0.930, indicating adequate internal validity. OPC, a still rare event, requires high specificity from a biomarker panel to minimize identification of false positives. A specificity of 99.1% was achieved with a sensitivity of 70.1%. Allowing specificity to drop to 98.1% resulted in a substantial gain in sensitivity to 78.0%.

Conclusions: Our data suggest this panel can detect OPC early. Internal validation showed high predictability, however external validation of this panel is needed. A panel of biomarkers to diagnose OPC earlier is needed to prevent complex treatment of OPC and associated co-morbidities, while reducing risk of recurrence.

Oral presentation

Dr Moreno Acosta Pablo
dajup63@yahoo.com - Colombia

Dr Moreno-acosta Pablo
dajup63@yahoo.com - Colombia

#3757

Predictive factors of response to treatment in invasive squamous cell cancer of the cervix

15 - Molecular markers

Moreno-acosta P, Gonzalez-prieto C, Gamboa O, Malaver G, Magné N

Background/Objectives: The most important risk factor for cervical cancer is HPV. Patients with cervical cancer in stages IIB and IIIB treated at the National Cancer Institute (Colombia) present a poor response to treatments such as radiotherapy (RT), at least 40% of these do not respond adequately. Within tumoral microenvironment, hypoxia in solid tumors, such as cervical cancer, induces genomic and proteomic changes, such as, for example, in IGF1R, GLUT1, HIF1a and hTERT, E6 HPV, changes that modulate response to treatment. We evaluate the possible relationship of the expression of the proteins IGF1R β , hTERT, HIF1a, GLUT1, the presence of certain HPV16 species and some clinical parameters with the response to radiochemotherapy and radiotherapy.

Methods: Type-specific HPV infection was detected using GP5 + / GP6 + PCR-RLB, and expression of the IGF1R, hTERT, HIF1a, GLUT1 proteins by immunohistochemistry in 21 patients with invasive squamous cell carcinoma of the cervix.

Results: HPV16 was the most frequent type followed by HPV58 and HPV56. HPV16 alpha 9 species was the most frequent, followed by alpha 6 and alpha 7. IGF1R was expressed in 85.7%, HIF1a in 61.9%, GLUT1 in 52.3% and hTERT was detected in both the cytoplasm and the nucleus. with the same frequency (90.4%). Hgb levels ≤ 11 d / dL (anemic hypoxia) were found in 28.6% and Hgb > 11 g / dL in 71.4%. 57.1% of the patients received radio chemotherapy and 42.9% exclusive radiotherapy. From the descriptive analysis of multiple correspondence in which two factorial axes were used accumulating 45.97% of the total variance, the MCA factor map showed a relationship between: nuclear and cytoplasmic expression of hTERT, and alpha 7 and alpha 9 species HPV16, observing similarity between HPV16 alpha 6 species and no expression of hTERT; GLUT1 and HIF1a; Hgb levels ≤ 11 d / dL and no response to radiotherapy, such as between Hgb levels ≤ 11 d / dL, expression of GLUT1 and HIF1a; HIF1a expression and hTERT expression; complete response and having received radio chemotherapy. In addition, a close relationship was observed between the nuclear and cytoplasmic expression of hTERT and the expression of IGF1R β , a relationship that was confirmed by applying Fisher's exact test ($p = 0.041$).

Conclusions: A better response to radiochemotherapy than to radiotherapy was confirmed, as well as non-response in the presence of anemic hypoxia and HIF1a expression. One of the findings was the subcellular location of hTERT, the possible relationship between alpha 7 and alpha 9 species of HPV16 and the expression of the hTERT protein, in addition, the relationship between the expression of hTERT and IGF1R, interaction that could contribute to resistance to radiation therapy.

References: Harima Y, Sawada S, Nagata K, Sougawa M, Ohnishi T. Human papilloma virus (HPV) DNA associated with prognosis of cervical cancer after radiotherapy. *Int J Radiat Oncol Biol Phys* 2002;52:1345-51. Badaracco G, Savarese A, Micheli A, Rizzo C, Paolini F, Carosi M, et al. Persistence of HPV after radiochemotherapy in locally advanced cervical cancer. *Oncol Rep* 2010;23:1093-9. Song YJ, Kim JY, Lee SK, Lim HS, Lim MC, Seo SS, et al. Persistent human papillomavirus DNA is associated with local recurrence after radiotherapy of uterine cervical cancer. *Int J Cancer* 2011;129:896-902. Ferdousi J, Nagai Y, Asato T, Hirakawa M, Inamine M, Kudaka W, et al. Impact of Human papillomavirus genotype on response to treatment and survival in patients receiving radiotherapy for squamous cell carcinoma of the cervix. *Exp Ther Med* 2010;1:525-30. Moreno-Acosta P, Gamboa O, Sanchez de Gomez M, Diaz GD, Romero A, BalartSerra J, et al. IGF1R gene expression as a predictive marker of response to ionizing radiation for patients with locally advanced HPV-16 positive cervical cancer. *Anticancer Res* 2012;32:4319-26. P. Moreno-Acosta, A. Vallard, S. Carrillo et al., "Biomarkers of resistance to radiation therapy: a prospective study in cervicalcarcinoma,"*Radiation Oncology*, vol. 12, no. 1, p. 120, 2017. P. Moreno-Acosta, S. Carrillo, O. Gamboa et al., "Potential biomarkers for personalized radiation therapy for patients with uterine cervical cancer,"in *Uterine Cervical Cancer*,S.Farghaly, Ed., Springer, Cham, 2019. Moreno-Acosta, P., Romero-Rojas, A., Vial, N. Magné, N. Persistent High-Risk HPV Infection and Molecular Changes Related to the Development of Cervical Cancer. *Case Reports in Obstetrics and Gynecology*, 2020; 17:1-6.

Oral presentation

Ms Guimarães Yasmin
yasminmg96@gmail.com - Brazil

Yasmin Guimarães
yasminmguimaraes@outlook.com - Brazil

#3758

Characteristics of ancestry patterns of multiracial women population with cervical lesions induced by HPV: Preliminary results from H2020 Elevate Study.

02 - Viral and molecular biology

Yasmin G, Luani G, Tauana D, Nathalia C, Heleen V, Olivier D, Ricardo D, Adhemar L.

Background/Objectives: Cervical cancer remains a public health problem worldwide, especially in middle and low-income countries as Brazil. The responsible for this tumor is the human papillomavirus (HPV), most notably types 16, 18 and 12 other high-risk types, capable of infecting epithelial tissues of the human body, especially the cervix. This infection is considered the most common sexually transmitted disease in the world and, in most cases, is eliminated by the immune system without major complications. However, persistent infection can induce carcinogenesis. Currently, the meaning of the infected woman's ancestry in the persistence of the HPV infection and the risk of developing cancer has been documented. However, in Brazil that has a highly admixed population, the knowledge about the biological influence of ancestry in cervical cancer is still incipient; for this reason, it is important to investigate whether differences in persistence caused by the women's ancestry can influence the HPV infection profile in distinct populations. Thus, we aimed to characterize the genetic ancestry of 500 Brazilian (250) and Belgium (250) women and to associate it with age, cytology status, and HPV genotyping data.

Methods: Liquid-based cervical samples are being used for ancestry and HPV genotyping. Ancestry analysis is performed using a set of 46 ancestry-informative markers (AIMs) among the most informative insertion and deletion polymorphisms (INDELs) for each population group. A 46plex PCR is performed and the amplified products are subsequently submitted to capillary electrophoresis in the ABI 3500 xL sequencer (Applied Biosystems). The electropherograms are analyzed and the genotypes are automatically assigned with GeneMapper v4.1 (Applied Biosystems). The proportions of ancestry are assessed using the Structure v2.3.3 software, considering the four main parental population groups: Native Americans, Europeans, Africans, and Asians as possible contributors to the current genetic background of the general population. For HPV genotyping, Cobas® 4800 HPV Test (Roche) is used. This PCR automatized platform can identify the infections by HPV16, HPV18 and/or a pool of other 12 high-risk HPV types (31, 33, 35, 39, 45, 51, 52, 56, 59, 66, 68). Statistical analysis will be performed with the Software SPSS 25.0 (IBM Corp) and only results with a p-value <0.05 will be considered significant.

Results: So far, 239 cases were fully analyzed for genetic ancestry. The estimated proportion of ancestry for Brazilian women is European=67%, African=15%, Native American=12%, and Asian=6%. For Belgium women, the estimation is European=89%, African=5%, Native American=3% and Asian=3%. Regarding HPV genotyping, 374 samples were already tested. For the 194 Brazilian samples, 57.3% were negative; among the positive ones, 28.6% were positive for the other 12 high-risk HPV types, 6.8% were positive for HPV16, 1.6% HPV18, 3.6% coinfection of others and 16, 1% coinfection of others and 18, 0.5% coinfection of 16 and 18, and 0.5% coinfection of other, 16 and 18. For the 180 Belgium samples, 68.8% were negative; among the positive ones, 18.8% were positive for the other 12 high-risk HPV types, 3.1% positive for HPV16, 1% for HPV18, and 1.6% coinfection of others and 16.

Conclusions: In the near future, we expect to verify if the HPV infection profile of Brazilian and Belgium women is associated with significant lesions' behavior differences when clustered by genetic ancestry, age, and cytology status.

Oral presentation

Prof. Dr Sláma Jiří
jiri.slama@vfn.cz - Czech Republic

Sláma Jiří
jiri.slama@vfn.cz - Czech Republic

#3761

FINAL DATA FROM LIBUSE TRIAL - ALGORITHM FOR CERVICAL CANCER SCREENING IN THE CZECH REPUBLIC WITH USAGE OF HPV DNA TESTING WITH HPV 16/18 GENOTYPING AND P16/KI-67 DUAL-STAINED CYTOLOGY

10 - HPV screening

Sláma J, Dvořák V, Trnková M

Background/Objectives: The incidence and mortality rates of cervical cancer in the Czech Republic have remained unchanged for more than 30 years regardless of the existing national screening based on annual Pap smear. The aim of our prospective study was to evaluate the role of HPV DNA testing with 16/18 genotyping and p16/Ki-67 immunocytochemical triage.

Methods: Women aged 30 to 60 years who underwent routine annual Pap smears at 12 collaborating centers were tested for HPV DNA with the addition of selective 16/18 genotyping (Cobas 4800, Roche) at 3-year intervals. All HPV 16/18-positive cases and/or cases with severe abnormal cytology were sent directly to colposcopy; HPV non-16/18-positive cases and LSILs were triaged using p16/Ki-67 dual-stain cytology (CINtec Plus, Roche) and positive cases were referred to colposcopy.

Results: Altogether 2407 patients were eligible for analysis. The mean age of the subjects was 43 years. Twelve cases with severe Pap smear abnormalities were identified throughout the study period. A total of 7.4% (180/2418) patients were found to be HPV positive and 5.4% were HPV positive at the end of the study, 2% were HPV 16/18 positive at baseline and 1.1% were HPV positive at the end of the study. Triage using p16/Ki-67 was positive in 22.5% of cases (29/129). After 3 years of follow-up, biopsy confirmed 51 HSILs and 2 glandular lesions, all of which were HPV positive.

Conclusions: Screening based on HPV testing with selective 16/18 genotyping and p16/Ki-67 triage demonstrated four times more high-grade lesions, including glandular lesions, than standard Pap smear-based screening over 3 years. After three years, there was a significant reduction in HPV positivity, HPV 16/18 positivity, high-grade lesions, and referral for colposcopy.

Oral presentation

Dr Iacobone Anna Daniela
annadaniela.iacobone@ieo.it - Italy

Iacobone Anna Daniela
annadaniela.iacobone@ieo.it - Italy

#3762

VALIDATION OF THE HPV ONCOPREDICT® ASSAY ON SELF-COLLECTED VAGINAL AND URINE SAMPLES: PRELIMINARY ITALIAN DATA OF THE EUROPEAN-VALHUDES STUDY.

13 - Self-sampling

Iacobone A, Bottari F, Preti E, Guerrieri M, Martella S, Giubbi C, Cocuzza C, Latsuzbaia A

Background/Objectives: HPV testing of minimally invasive self-collected samples represents a promising strategy to improve cervical cancer screening. The aim of the present study is to evaluate the performance of the HPV OncoPredict® assay, a double step reflex test consisting of an initial screening for HR-HPV E6/E7 sequences and a second quantitative assessment of normalized type-specific viral loads, on self-collected vaginal and urine samples as compared to clinician-taken cervical samples, according to VALHUDES (VALidation of HUMAN papillomavirus assays and collection DEVICES for HPV testing on Self-samples and urine samples) protocol [1].

Methods: Women aged 25-64 years referred to colposcopy at the European Institute of Oncology, Milan, were enrolled after giving informed consent. Prior to colposcopy, first-void urine and vaginal self-samples were taken with the ColliPee (Novosanis) and the FLOQSwab (Copan) collection devices, respectively. Physician collected cervical samples with the Cervex-Brush (Rovers Medical Devices). Colposcopy was performed with targeted biopsy if required. All samples were tested with the HPV OncoPredict® QT assay (Hiantis), a recently developed PCR-based test evaluated as part of WHO HPV Labnet international proficiency study [2].

Results: Overall, 288 samples were collected from 96 women. Forty-two out of 96 patients received colposcopy-targeted biopsy and histology report was CIN2+ in 60% (25/42) of cases. Sample host cell cellularity was similar in physician- and self-collected samples. Mean Log10 (copies CCR5/rx) was 3.840, 3.938 and 3.820 in cervical, vaginal and urine samples, respectively. The hrHPV test result concordance with cervical sample was 83.3% (80/96) for first-void urine and 82.3% (79/96) for vaginal samples. The most prevalent genotypes were HPV 16 and 31. Viral load was higher in cervical samples than self-collected samples; but it varied according to detected genotype. Interestingly, in nine HPV16-related CIN2+ mean viral load was 3.183, 3.407 and 2.748 Log10 (copies HPV16/rx) in cervical, vaginal and urine samples, respectively; whereas it was 3.870, 3.851 and 3.409 Log10 (copies HPV31/rx) in cervical, vaginal and urine samples of five women affected by HPV31-related CIN2+.

Conclusions: Preliminary data indicate a good overall agreement between HPV OncoPredict® assay results from self-collected and cervical samples. Furthermore, HPV genotyping and viral load may be useful as potential biomarkers in the triage of women with a hrHPV-positive self-sample. Acknowledgments: HPV OncoPredict® assays were developed as part of a European Commission funded SME Instrument Phase 2 Project (Grant agreement ID: 806551).

References: Arbyn M, Peeters E, Benoy I, et al. VALHUDES: A protocol for validation of human papillomavirus assays and collection devices for HPV testing on self-samples and urine samples. J Clin Virol. 2018;107:52-56. Eklund C, Mühr LSA, Lagheden C, et al. The 2019 HPV Labnet international proficiency study: Need of global Human Papillomavirus Proficiency Testing. J Clin Virol. 2021;141:104902.

Oral presentation

Dr Leite Sara
saradiasleite@hotmail.com - Portugal

Dr Dias Leite Sara
saradiasleite@hotmail.com - Portugal

#3763

THE WERTHEIM-MEIGS HYSTERECTOMY IN SURGICAL TREATMENT OF CERVICAL CANCER: SIX YEARS' EXPERIENCE

35 - Conventional therapies

Dias Leite S, Veiga Guimaraes M, Vide M, Petiz A

Background/Objectives: Cervical cancer is the third most common cancer among women. Surgical management is the primary therapy for most patients with early-stage cervical cancer. The Wertheim- Meigs surgery is an effective treatment for low risk patients with stage IA2, IB1, IB2 and IIA cervical cancer. Assess the treatment and follow-up of the patients when Wertheim-Meigs surgery has been performed for cervical cancer.

Methods: Retrospective observational study with descriptive data analysis of the patients who underwent Wertheim-Meigs surgery due to cervical cancer at the Portuguese Institute of Oncology - Oporto (IPO-Porto) between 2013 and 2018.

Results: During a six year period between 2013 and 2018, 70 patients underwent the Wertheim-Meigs surgery due to cervical cancer, with a median age of 49 years. The clinical FIGO stage was IA2 in 11%, IB1 in 83%, IB2 in 1.6% and IIA1 in 4.7% of the patients. A laparotomy surgery was performed in 89% of the cases. The vast majority of the patients did not present linfovacular invasion (73%) nor parametrial (90%). Six women (8,7%) had lymph node invasion, five with only one positive node and one patient with two. The majority did not have indication for adjuvant treatment, 29% underwent brachytherapy, 26% external radiotherapy and 1.4% chemotherapy. Squamous cell carcinoma was identified in 38 patients (54%), adenocarcinoma in 28 (40%), adenoid cystic carcinoma in 3 (4%) and adenosquamous carcinoma in 1 (1%). Two patients had disease recurrence (2,9%), one with adenoid cystic carcinoma and the other with adenocarcinoma. The minimum follow-up time in IPO-Porto was 5 years in the majority of the cases. Four deaths were reported (5,7%), with only one directly related to cervical cancer. This was a patient with adenoid cystic carcinoma with local recurrence and distant metastases identified 8 months post surgery, dying after a period of 37 months of follow-up.

Conclusions: The Wertheim Meigs surgery seems to be an effective treatment in the early stages of cervical cancer, with a low recurrence and mortality rate in the studied population.

Oral presentation

Mr De Bondt Daniël
d.debondt@erasmusmc.nl - Netherlands

Mr De Bondt Daniël
d.debondt@erasmusmc.nl - Netherlands

#3766

THE IMPACT OF HPV VACCINATION ON HPV PREVALENCE, PRE-CANCERS, CANCERS: A SYSTEMATIC REVIEW ON OBSERVATIONAL AND MODEL OUTCOMES.

40 - Public health

Naslazi E, De Bondt D, De Kok I, Jansen E, Hontelez J

Background/Objectives: HPV vaccination has been implemented in most high-income countries. With the new (partly) vaccinated cohorts approaching screening ages, policymakers need to re-evaluate how to shape screening for these cohorts. It will be vital to understand the impact of vaccination on the prevalence of high-risk HPV, pre-cancerous lesions, and/or cancers in vaccinated and unvaccinated women. Comparing observations to model predictions can give important insights into the validity of models and their predictions, and can help set benchmarks for improvement of modelling precision in the future.

Methods: We searched Cochrane Library, EMBASE, MEDLINE, Web of Science for studies published between 2005 to present. We performed three independent literature searches for studies that reported on effects of HPV vaccination in a high-income country: 1) Systematic reviews, 2) Observational studies performed after the last systematic review was published, and 3) Studies that used HPV transmission models. Studies were eligible if they reported both pre- and post-vaccination outcome figures on a population level, included vaccination coverage, and if they examined equal population characteristics before and after vaccination. Relative risk (RR) for HPV-related health outcomes was calculated comparing the pre-vaccination and post-vaccination periods. These risks were then compared to prevalence reductions found in the modelling studies.

Results: We identified a total of 7963 potential articles, and included five systematic reviews, 11 observational studies and 20 modelling studies in 18 different high-income countries. HPV prevalence, CIN and cervical cancer reduction were used as an outcome in different studies. All studies reported significant vaccination effectiveness in all three outcomes (mean RR: 0.30 for HPV, 0.51 for pre-cancers and 0.71 for cervical cancers). The effect was more heterogeneous for later disease outcomes. Most studies showed a relationship between vaccination coverage and effectiveness, with greater decreases in HPV-related outcomes with higher coverage. This relationship was also found among the modelling studies.

Conclusions: A combined assessment of both observed evidence and modelling predictions allows to more confidently evaluate the effects of HPV vaccination in high-income countries. This will assist policymakers in reshaping their screening programs.

Oral presentation

Prof. Sparen Par
par.sparen@ki.se - Sweden

Sparén Par
par.sparen@ki.se - Sweden

#3767

Quadrivalent HPV vaccination and long-term risk of adverse pregnancy outcomes

06 - HPV prophylactic vaccines

Lei J, Sparén P

Background/Objectives: Current available HPV vaccines have proven to be highly efficacious and effective in preventing HPV infection, genital warts, precancerous lesions, and cervical cancer. Recent studies also indicate that birth outcomes might be positively affected by HPV vaccination. However, long-term safety data regarding the impact of HPV vaccination on adverse pregnancy outcomes is absent. We aimed to evaluate the long-term association between the quadrivalent HPV vaccine and adverse pregnancy outcomes, leveraging nationwide high-quality Swedish register data

Methods: We included 470,000 singleton pregnancies (gestational week ≥ 22 weeks) among nulliparous women aged 16 to 35 in Sweden between Jan 1, 2006 and 31 Dec, 2018. We obtained the exposure to quadrivalent HPV vaccination, pregnancy outcomes and confounding variables from nationwide Swedish registers, linked at individual level. Confounding variables were: county of residence, calendar year of delivery (as a continuous variable), mother's country of birth, highest educational level, highest annual household income level, maternal age (as a spline term with 5 degrees of freedom), smoking, body mass index, assisted reproduction and maternal diagnoses. We estimated the risk of adverse pregnancy outcomes including stillbirth, preterm delivery, premature rupture of membranes (PROM) at 37+ weeks, preterm premature rupture of membranes (pPROM), low birth weight, small-for-gestational age, neonatal mortality and medically induced delivery in relation to HPV vaccination prior to child birth. We also performed a sensitivity analysis where women receiving quadrivalent HPV vaccination after the onset of pregnancy (n~100) were excluded.

Results: After controlling for potential confounders, we found a 13% lower risk (OR 0.87; 95% CI, 0.80 to 0.95) for small-for-gestational age among vaccinated women in comparison to not vaccinated women. In the main analysis we observed point estimates below unity for stillbirth, moderately preterm, PROM in deliveries at 37+ weeks and low birth weight, but none of these were statistically significant. No statistically significant differences were found for other pregnancy outcomes including preterm delivery, pPROM, neonatal mortality and medically induced delivery. In the sensitivity analysis the lower risk for SGA after quadrivalent HPV vaccination was confirmed, and there was also a statistically significant lower risk for stillbirth (OR 0.78; 95% CI, 0.61 to 0.99).

Conclusions: We found no evidence that quadrivalent HPV vaccination should be associated with elevated long-term risk of adverse pregnancy outcomes. We also found indications of lower risks for SGA and stillbirth, although possible mechanisms behind this could not be tested in the current study.

Oral presentation

Marini Bruna
b.marini@ulissebiomed.com - Italy

Marini Bruna
b.marini@ulissebiomed.com - Italy

#3769

A new PCR-based detection methodology for Human Papillomavirus genotyping on cervical brushes and self-collected vaginal samples

14 - Genotyping

Avian A, Clemente N, Mauro E, Isidoro E, Di Napoli M, Dudine S, Del Fabro A, Morini S, Perin T, Giudici F, Cammisuli T, Foschi N, Mocenigo M, Montrone M, Modena C, Polenghi M, Puzzi L, Riva E, Angeletti S, Ciccozzi M, Castriciano S, Pachetti M, Zella D, Centonze S, Gerin D, Banks L, Marini B, Canzonieri V, Sopracordevole F, Zanconati F, Ippodrino R

Background/Objectives: The study aims at evaluating the diagnostic performance of HPV Selfy, a new PCR-based detection and genotyping methodology for high risk Human Papillomavirus (HPV) on clinician-collected cervical brushes and self-collected vaginal samples in a screening population. HPV Selfy is able to perform full high risk HPV genotyping in the same PCR reaction.

Methods: From September 2018 to February 2019 all the women consecutively referring to Azienda Sanitaria Integrata Giuliano Isontina (ASUGI - Trieste, Italy) and to CRO - National Cancer Institute (Aviano, Italy) for the Italian cervical cancer screening programme were invited to participate in the study (n=1143). Eligible women were asked to perform both a vaginal self-sampling analyzed with our new PCR-based detection methodology for HPV and a clinician-collected cervical brush for Pap smear, analysed with the Hybrid Capture® 2 DNA test analysis. All the women enrolled were asked to fill a proper questionnaire about self-sampling acceptability. Moreover, a second group of women (n=500), whose traditional screening test was delayed because of the COVID-19 pandemic, was enrolled in the period February 2021-April 2021. Women were mailed a kit at home for self-collection and returned the sample to the hospital for the analysis with the HPV Selfy test.

Results: HPV Selfy was clinically validated on cervical brushes and vaginal swabs, reaching non-inferior performance to the reference test. HPV Selfy shows high concordance between self-collected and expert-collected specimens for HPV detection. According to the questionnaire response, 98.4% of women agreed that self-sampling test format was easy and effective. Genotyping analysis showed that prevalence of oncogenic HPV types was 9%; the five most frequent high risk HPV types detected in the cytology-negative screening population were HPV16 and HPV31, followed by HPV39, HPV51 and HPV66. On the other hand, in the CIN2+ population, HPV16 was the predominant strain followed by HPV31 and HPV58. Through a second prototype assay, able to genotype also 10 low-risk and probably high risk HPVs, we found a low and probable high risk HPV prevalence of 10% in the screening population.

Conclusions: The study demonstrated that HPV Selfy is potentially suitable for primary screening purposes performed on clinician-collected samples, providing also full genotyping information in the same reaction. According to several studies, genotyping information will become valuable for future triage strategies development. Despite hrHPV genotyping has not achieved widespread acceptance among clinicians, significant differences have been observed in the oncogenicity of hrHPV genotypes. HPV Selfy can also be used in combination with self-sampling tools to rescue women difficult to reach with traditional screening methods. HPV self-sampling could be an attractive solution to increase women's participation, regardless of age, education level, and other possible social parameters. It has been also demonstrated that self-sampling will be particularly relevant nowadays, when COVID-19 pandemic deeply impacted the effectiveness of hospital-based organized screening programmes.

Poster

Dr Smet Carolina
carol_smet@hotmail.com - Portugal

Dr Smet Carolina
carol_smet@hotmail.com - Portugal

#3772

CERVICAL CANCER DIAGNOSED AFTER THE DETECTION OF PROTEIN P16 IN AN ILIAC ADENOPATHY

26 - Cervical neoplasia

Beja M, Martins A, Cruz E, Pinto D, Costa C

Background/Objectives: The p16 protein is overexpressed in HPV-associated tumours and there appears to be a higher expression in more severe lesions. We report a case of cervical cancer suspected after the biopsy of an iliac adenopathy which was positive for p16.

Methods: Case report based on the information collected from the clinical records of the patient.

Results: A 44-year-old woman went to the emergency room with asthenia, anorexia, weight loss and fever. The patient mentioned these symptoms had begun a month prior. She denied gynecological symptoms. On physical examination she had multiple inguinal adenopathies and one supraclavicular. A computerized tomography was performed and detected multiple peri-aortic and iliac adenopathies. A biopsy of an iliac adenopathy revealed a squamous cell carcinoma which was strongly positive for p16, suggesting the primary origin of the metastasis to be a cervical cancer. The gynecological observation showed an apparently normal cervix and the gynecologic ultrasound was normal. The cytology revealed a HSIL. Colposcopy showed in the posterior lip a red lesion with irregular margins, areas of ulceration and exophytic lesions with atypical vascularization, that stained with acetic acid. Biopsies of the lesion confirmed the suspicion of a cervical cancer. Although the supraclavicular adenopathy was palpable, it did not show on the PET scan. The patient is currently under chemoradiotherapy.

Conclusions: p16 immunostaining can be an important aid in determining the probable primary tumour site in the context of metastatic squamous cell carcinoma. When positive, and in the appropriate clinical and imagiological context, it should raise the suspicion of a cervical primary.

Poster

Dr Cochicho Daniela
dcochicho@ipolisboa.min-saude.pt - Portugal

Martins Luís Carlos
lmartins@ipolisboa.min-saude.pt - Portugal

#3776

CERVICAL CANCER SCREENING: PRELIMINARY RESULTS OF 22 ACES IN THE REGION OF LISBON AND TEJO VALLEY

10 - HPV screening

Martins L, Cochicho D, Roque R, Cunha M, André S

Background/Objectives: The implementation of organized screening programs would allow early diagnosis. Abnormalities found on screening need follow-up, diagnosis and treatment, to prevent the development of cancer or to treat cancer at an early stage. Despite the high coverage of HPV vaccines in Portugal, introduced in 2008 in the National Immunization Plan for girls and more recently (2020) for boys, an organized screening based on the HPV test was implemented for cervical cancer at the national level. AIM We intend to verify the prevalence of HPV HR types in the first organized screening program in the region of Lisbon and Tejo Valley.

Methods: A total of 7793 women (age range 29-66 years) were screened with HPV HR test as the main method for cervical cancer screening, between December 2018 and August 2021. Samples were collected from 22 ACES (Public Health Centers). Clinical algorithms refer all HPV16 or HPV18 positive women for colposcopy and all samples with other HR HPV types were forwarded to cytological evaluation, owing to the type of cell abnormalities women could be referred for colposcopy.

Results: The percentage of women negative for HPV HR types is 88,49% (n=6897) while women positive for at least one HR HPV type is 11,02% (n=859). 20,37%. The number of women that test positive for HPV16 and/or HPV18 is 175 (20,37%) and positive for others HR HPV types (HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV56, HPV58, HPV59, HPV66, HPV68) is 684 (79,63%). As expected HPV infection tends to decrease with age. Women with HR HPV types who underwent reflex cytology, 55,26% (n=205) were negative, 21,58 (n=80) had ASCUS, 2,96% (n=11) ASC-H, 0,81% (n=3) AGC, 15,90% (n=59) LSIL, 3,23% (n=12) HSIL and 0,27% (n=1) CPC. In terms of HPV prevalence, HPV68 (17,69%) is the most prevalent, followed by HPV16 (14,67%), HPV31 (12,92%), HPV 52 (11,64%) and HPV51 (10,36%), all other HPV detected as a prevalence less than 10% (HPV18 has a prevalence of 6,29%).

Conclusions: HPV 16 is usually the most prevalent HPV detected in screening programs, the fact that in our study HPV68 is the most prevalent may be related to the introduction of the vaccine in the National Vaccination Plan since women that were vaccinated are already being screened. The collection of more data is being carried out to give more strength to the data already obtained, namely: HPV vaccination, age of menarche, type of contraception.

Oral presentation

Mrs Kepka Deanna
deanna.kepka@hci.utah.edu - United States

Dr Kepka Deanna
deanna.kepka@hci.utah.edu - United States

#3777

Intention to receive COVID-19 vaccination relates to HPV vaccination receipt among young adults ages 18-26 years in the western region of the United States

40 - Public health

Christini K, Mann S, Chang E, Lipscomb J, Kepka D

Background/Objectives: To assess similarities and differences between receipt of the HPV vaccine and intention to receive the COVID-19 vaccine among young adults by demographic factors in the western region of the United States.

Methods: An online, self-administered, cross-sectional survey was conducted from October 2020 to April 2021 that included young adults ages 18-26 years old who live in rural and urban communities in the western region of the United States. Participants were recruited through university, public health, and community list serves (N=2889). The questionnaire asked participants to provide information on their demographics; general levels of trust in vaccinations; HPV and HPV vaccination knowledge, trust in HPV vaccine effectiveness, and receipt of the HPV vaccine. The questionnaire also included COVID-19 questions related to COVID-19 testing, COVID-19 concerns, and COVID-19 vaccination intentions. We hypothesized that a difference in COVID-19 vaccination intention between those who had received an HPV vaccination or not would be found, expecting the decision-making process for getting the HPV vaccine and the COVID-19 vaccine to be similar. HPV vaccination was self-reported. Chi-square and penalized binary multivariate logistic regression were utilized.

Results: After controlling for the effect of gender, age, race and ethnicity, health insurance, frequency of religious services, and rurality, the odds of young adults who had received any HPV vaccination stating that they intended to get COVID-19 vaccinated was 1.58 (95%CI: 1.33-1.88) times higher than those who had not had any doses of the HPV vaccine. After model adjustments, being male (1.41, [1.16 - 1.72]), identifying as non-Hispanic racial minority (1.33, [1.03 - 1.72]), having health insurance (0.49, [0.39 - 0.61]), and living in rural areas (0.50, [0.41 - 0.59]) all had significantly different odds of COVID-19 vaccination intent. Rural young adults stated, at significantly higher frequencies, that they would be unlikely to receive the COVID-19 vaccine, even if recommended by their health care provider, than did urban young adults (24% vs 7%, $P<0.001$).

Conclusions: Our research indicates that the decision-making process of intent to receive COVID-19 vaccination among young adults and that of receiving HPV vaccination are similar, and that the presence of any HPV vaccination may be a strong predictor of intent to vaccinate against COVID-19 infection among young adults. Targeted interventions are needed to address both HPV vaccine hesitancy and COVID-19 vaccine hesitancy to improve vaccination rates. Rural communities in the western United States would benefit greatly from interventions that address HPV and COVID-19 vaccine hesitancy among young adults.

References: DAG modeled using Dagitty.net. Judea Pearl, Madelyn Glymour, and Nicolas P Jewell. Causality Inference in Statistics: A Primer. Wiley, New York, NY, USA. 1st edition, 2016.

Oral presentation

Ms Parker Susan
susan.parker2@bcm.edu - United States

Montealegre Jane
jrmontea@bcm.edu - United States

#3779

Influence of COVID-19 pandemic on participation in HPV self-testing trial

01 - HPV disease and COVID-19

Parker S, Daheri M, Benavides E, Kasbaum M, Escobar B, Chiao E, Schmeler K, Anderson M, Deshmukh A, Hilsenbeck S, Jibaja-weiss M, Bulsara S, Zare M, Scheurer M, Montealegre J

Background/Objectives: The COVID-19 pandemic led to health system restrictions and closures, leading to gaps in preventive care. To our knowledge, the relationship between the COVID-19 pandemic and use of HPV self-test kits has not been assessed among patients in a large urban safety net health system that provides services to predominantly racial/ethnic minority, medically underserved populations. We aimed to describe the influence of the COVID-19 pandemic on the use of mailed HPV self-test kits within a safety net health system.

Methods: We assessed the influence of the COVID-19 pandemic on the use of mailed HPV self-test kits as part of a randomized controlled trial to evaluate the effectiveness the kits. Participants are women, ages 30-65 years in an urban safety net healthcare system who are underscreened for cervical cancer. We conducted a telephone survey in English and Spanish among a subgroup of trial participants who were randomized to receive the at-home HPV self-test kits.

Results: Among the 124 women who have completed the survey to date, the majority are Hispanic (70.2%) and Black or African American (20.2%). Among the women surveyed, 78.1% reported using the kit to take a sample. Of those who used the kit, over half (58.5%) said that the COVID-19 pandemic influenced their decision to use the kit. When asked in what way the pandemic influenced their decision, most of the open-ended responses fell into three categories: (1) women felt safer collecting their sample at home for fear of being exposed to COVID, (2) their clinic was closed due to shut-down measures or they had difficulty getting an appointment, and (3) they felt more comfortable/less embarrassed collecting their sample at home. The majority of women who used the kit (87.7%) said the kit was easy to use and almost all (96.3%) stated they would be willing to use an HPV self-sample kit again.

Conclusions: Preliminary results from our survey suggest that the COVID-19 pandemic influenced women an urban safety-net healthcare system to participate in an at-home HPV self-test trial. Overall, the kit had high acceptability among trial participants.

Poster

Dr Kerr Colm
colmkerr@gmail.com - Ireland

Dr Kerr Colm
colmkerr@gmail.com - Ireland

#3783

ANAL CYTOLOGY AND MOLECULAR FINDINGS FROM HIV POSITIVE AND NEGATIVE MSM IN IRELAND

28 - Anal neoplasia

Christine W., Joseph H., Helen K., Emer C., Stephen R., Padmaja N., Corinna S., John O., Cara M., Colm B

Background/Objectives: Almost 90% of anal cancers globally are attributable to HPV infection. While rare amongst the general population, men who have sex with men (MSM), particularly HIV positive MSM, are far more at risk for the development of anal cancer. This research aimed to describe the anal cytology and molecular findings from HIV positive and negative MSM in Ireland

Methods: 252 MSM participants were recruited to this study. Anal swabs were collected from participants and placed in PreservCyt. hrHPV DNA testing was performed using the Cobas HPV test on the Cobas 4800 Platform (Roche diagnostics). HPV mRNA testing was performed using the Aptima HPV assay (Hologic). Samples also underwent anal pap staining for cytological analysis. Samples that tested positive for the presence of hrHPV DNA underwent further analysis using P16/Ki67 dual staining. 148 participants underwent repeat testing at 1 year.

Results: 234 participants had valid baseline samples for HPV16 DNA analysis. 52 (22%) tested positive for HPV16 DNA. 232 had valid baseline samples for HPV18 DNA analysis of which 22 (9%) tested positive. 238 participants had valid baseline samples for other hrHPV DNA analysis of which 139 (58%) tested positive. HIV positivity was statistically significantly associated with the presence of other hrHPV DNA on multivariate analysis (OR 2.87, 95% CI 1.34 - 6.14, $p=0.007$). Overall, 240 participants had valid baseline samples for any hrHPV DNA analysis, of which 153 (64%) tested positive for at least one hrHPV type. HIV positivity was statistically significantly associated with the presence of any hrHPV DNA on multivariate analysis (OR 2.71, 95% CI 1.29 - 5.70, $p=0.008$). 252 participants had valid baseline samples for hrHPV mRNA analysis, of which 101 (40%) tested positive. Current smoking was statistically significantly associated with hrHPV mRNA positivity on multivariate analysis (OR 2.02, 95% CI 1.11 - 3.68, $p=0.022$). 245 samples were valid for cytological analysis. 110 (45%) were reported as normal, 81 (33%) were reported as AIN1 (LSIL), 46 (19%) were reported as AIN2 (HSIL) and 8 (3%) were reported as AIN3 (HSIL). Receptive anal intercourse (RAI) was statistically significantly associated with the presence of HSIL (OR 8.81, 95% CI 1.91 - 40.65, $p=0.005$). 145 returned valid P16-Ki67 dual staining results. 50 (34%) of these samples were positive for P16/Ki-67 dual staining. Being originally from Ireland was significantly associated with lower rates of dual staining positivity on multivariate analysis (OR 0.25, 95% CI 0.10 - 0.60, $p=0.002$). At 1 year follow up, hr HPV mRNA persistence was seen in 60% of participants (29/48). HPV16 DNA persistence was seen in 77% (17/22), HPV18 DNA persistence was seen in 44% (4/9), and other hrHPV DNA in 77% (58/75). Overall, persistence of any hrHPV DNA was seen in 79% (63/80). HSIL persistence was present in 22% (7/32), with AIN3 persis

Conclusions: This study demonstrates the molecular burden of hrHPV infection and persistence of hrHPV infection in this high risk group in an Irish context for the first time. It also demonstrates the burden of cytological HPV disease in this high risk cohort. In light of recent data showing the benefit of treating anal cancer precursor lesions in HIV positive MSM, this study supports calls for the consideration of the currently unmet need in Ireland for anal cancer screening in this high risk population.

Oral presentation

Dr Mesquita Pinto Ana Rita
anaritapinto@hotmail.com - Portugal

Sousa Rita
ritamsousacf@gmail.com - Portugal

#3784

LASER VAPORIZATION FOR TREATMENT OF LOW-GRADE CERVICAL LESIONS: A SINGLE CENTRE STUDY

25 - Colposcopy

Mesquita Pinto R, Silva S, Carvalho C, Raposo S, Sousa R

Background/Objectives: Low grade intraepithelial lesions (LSIL) of the cervix are histological manifestations of HPV infection, with a very high regression rate, specially in young women. Management of this patients is an issue of debate, being an option for persistent lesions preceded by minor findings, destructive treatments. The aim of this study was to evaluate the success of laser vaporization for low grade cervical lesions.

Methods: Retrospective study of the 104 women with biopsy-proven CIN1 who underwent laser vaporization between January and December 2020 in Instituto Português de Oncologia de Coimbra, Portugal. The regression and persistence of the lesion were evaluated.

Results: Most of them, 54 (52%) women returned to the regular screening program. Thirty eight (36,5%) women remained in the colposcopy clinic for follow-up, with one case of excisional treatment, due to long-term lesion persistence, after two vaporizations. Twelve of the patients missed the follow-up appointment (11,5%). Eleven patients were vaccinated and 23% smokers.

Conclusions: Most of low grade cervical lesions were successfully treated with laser vaporization (52%), presenting a negative HPV test at 6 months-follow-up. The need of a posterior excisional procedure was of one case (< 1%). CO2 laser vaporization, a simple procedure, performed in outpatient clinics, may be a successful option for low grade cervical lesions.

Poster

Mr Dafa'alla Tarig
tarig.dafaalla@genefirst.com - United Kingdom

Mr Dafa'alla Tarig
tarig.dafaalla@genefirst.com - United Kingdom

#3803

VALIDITY OF A NEW URINE COLLECTION MEDIA IN STABILISING RNA FOR HPV-MRNA DETECTION

13 - Self-sampling

Horne R, Wu W, Fu G

Background/Objectives: Numerous recent studies have recommended the use of urine samples as an alternative to vaginal swabs and cervical samples for HPV DNA detection in regular screening programmes^{1,2}. Urine sample collection is simple and non-invasive when compared to alternative methods that are currently in use.

Methods: In this study, we evaluated two urine preservation media for the detection of HPV mRNA. One of the media, UCM, is included in the Colli-Pee urinary self-sampling device developed for HPV DNA detection (Novosanis). The second medium, UAS, is included in Colli-Pee urine collection device under development for HPV mRNA detection. To evaluate the performance of the two media, a spike RNA from cervical adenocarcinoma (HeLa-S3, ThermoFisher) or combined in vitro transcripts of IC, HPV16, HPV18 and HPV45, prediluted in normal saline (0.9%), was used in the ratios recommended by the manufacturer for each medium. The in vitro transcripts used contain template DNA in addition to the transcribed RNA. Test samples, each containing 20 ng of either HeLa RNA or the in vitro transcribed RNA, were prepared from each medium. To test the capability of these media in preserving RNA and DNA, the test samples were incubated (in triplicate) at different temperatures and for different lengths of time. These include incubations at room temperature (RT) for 48 hr, at RT for 8 days, or at 37°C for 8 hr followed by an overnight incubation at RT. The integrity of RNA and DNA in the samples was assessed by real-time RT-PCR using Papilloplex®HR-HPV mRNA Detection Kit and the obtained Ct-values for the different targets were statistically compared for significant differences.

Results: The UAS medium was found to be good at preserving both RNA and DNA as no significant change in the detected Ct-value of all tested targets in HeLa-S3 cells and in vitro transcribed targets between the fresh samples and those incubated for 48 hr at RT, 8 days at RT or 8 hr at 37°C. The UCM medium which was designed to preserve DNA was found to be good at preserving DNA under all the tested conditions where no significant loss in the amount of detected DNA was observed in any of the tested in vitro transcription samples. This medium, however, was found not suitable for preserving RNA since no RNA could be detected even the samples incubated for 48 hr at RT.

Conclusions: UAS medium is capable of stabilising both RNA and DNA and is potentially suitable for collection of urine samples required for HPV mRNA/DNA detection. UCM medium is suitable for preserving urine samples for DNA detection.

References: 1. Choi, Y. S., Jin, H. & Lee, K. E. Usefulness Analysis of Urine Samples for Early Screening of Human Papilloma Virus Infection. J. Cancer Prev. 24, 240 (2019). 2. Sargent, A., Fletcher, S., Bray, K., Kitchener, H. C. & Crosbie, E. J. Cross-sectional study of HPV testing in self-sampled urine and comparison with matched vaginal and cervical samples in women attending colposcopy for the management of abnormal cervical screening. BMJ Open 9, 1-7 (2019).

Oral presentation

Dr Di Lorenzo Antonio
antoniodilorenzo95@gmail.com - Italy

Di Lorenzo Antonio
antoniodilorenzo95@gmail.com - Italy

#3808

Surveillance of adverse events following HPV9 vaccine in a Region which implemented UMV: data from Puglia (Italy)

06 - HPV prophylactic vaccines

Berardi P, Di Lorenzo A

Background/Objectives: Human Papillomavirus (HPV) is responsible for epithelial lesions and cancers in both males and females. The latest licensed HPV vaccine is Gardasil-9, a 9-valent vaccine effective both against high-risk HPV types and those responsible for non-cancerous lesions. The Italian Vaccination Schedule for Life of National Vaccine Prevention Plan 2017-2019 recommends 2 or 3 doses of HPV vaccine, according to the vaccine and the patient's age; the plan's targets are both males and females aged 12 and older and women over 25 years of age not yet immunized. Vaccination is also offered actively and free-of-charge to high-risk groups, such as MSM or women with HPV-related lesions. The surveillance of the safety profile of recently introduced vaccines is a key strategy for the success of immunization campaigns; in Italy, surveillance of Adverse Events Following Immunization (AEFIs) is carried out by the Italian Drug Authority (AIFA) in collaboration with Regional Health Authorities. This report describes the adverse events following Gardasil-9 administration reported in Puglia, a region in the South of Italy, from January 2018 to November 2021.

Methods: This is a retrospective observational study. Data about Gardasil-9 AEFIs were collected from AIFA's database. The overall number of Gardasil-9® doses administered during the study period was extrapolated from the regional online immunization database (GIAVA). AEFIs were classified as serious or non-serious accordingly to World Health Organization (WHO) guidelines, and serious ones underwent causality assessment, as recommended by WHO, in order to determine their causal association with the vaccination.

Results: During the study period, 266,647 Gardasil-9 doses were administered in Puglia and 22 AEFIs were reported, with a reporting rate (RR) of 8.25/105 doses. 15 out of 22 AEFIs regarded subjects under 18 years of age, and the male/female ratio was 0.833 (10 males vs. 12 females). Neurological symptoms were the most reported (7/22, RR: 2.62/105 doses), followed by local events of pain, swelling or erythema (6/22, RR: 2.25/105 doses) and allergic reactions (6/22, RR: 2.25/105 doses). 5 AEFIs (22.7%, RR: 1.87/105 doses) were serious, and 2 of these led to hospitalization. One case of permanent impairment occurred. Following causality assessment, only 2 out of 5 serious AEFIs were deemed consistently associated with the vaccination. One of them consisted in a severe skin rash localized to the torso followed by glottis oedema on the next day. The second one consisted in fainting, occurred one hour after the vaccine's administration. The RR for vaccine-related AEFIs was therefore 0.750/105 doses. 15 out of 22 AEFIs (68.2%) healed completely, while for 3 of them (13.6%) only partial healing occurred. 2 out of 22 patients (9.10%) were still suffering from the AEFIs, while for 2 the outcome is still not known.

Conclusions: Our data are similar to pre-licensure evidence as far as the AEFIs' nature is concerned. The RR though is far lower than the ones described in clinical trials; this may be due to the different data collection method: our study used passive surveillance, while pre-marketing studies generally employ active call. While allowing a closer surveillance of rare AEFIs, in fact, passive surveillance on large populations determines an increased risk of under-reporting, especially for mild AEFIs. Gardasil-9's safety profile seems favourable, with a low rate of serious adverse events and an advantageous risk/benefits ratio.

References: 1.Luria L, Cardoza-Favarato G. Human Papillomavirus. 2021 Jan 24. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. PMID: 28846281. 2.Stillo M, Carrillo Santistevé P, Lopalco PL. Safety of human papillomavirus vaccines: a review. *Expert Opin Drug Saf.* 2015;14(5):697-712. doi:10.1517/14740338.2015.1013532 3.Schmiedeskamp MR, Kockler DR. Human papillomavirus vaccines. *Ann Pharmacother.* 2006;40(7-8):1344-1352. doi:10.1345/aph.1G723 4.Cheng L, Wang Y, Du J. Human Papillomavirus Vaccines: An Updated Review. *Vaccines (Basel).* 2020;8(3):391. Published 2020 Jul 16. doi:10.3390/vaccines8030391 5.Petrosky E, Bocchini JA Jr, Hariri S, et al. Use of 9-valent human papillomavirus (HPV) vaccine: updated HPV vaccination recommendations of the advisory committee on immunization practices. *MMWR Morb Mortal Wkly Rep.* 2015;64(11):300-304. 6.Conferenza permanente per i rapporti tra lo stato le regioni e le province autonome di Trento e Bolzano, Intesa, ai sensi dell'articolo 8, comma 6, della legge 5 giugno 2003, n. 131, tra il Governo, le regioni e le province autonome di Trento e Bolzano sul documento recante "Piano nazionale prevenzione vaccinale 2017-2019" (Rep. atti n. 10/CSR) (17A01195). Available online at <https://www.trovanorme.salute.gov.it/norme/dettaglioAtto?id=58185&completo=true> last accessed 11 October 2021 at 16:40. 7.REGIONE PUGLIA. CALENDARIO VACCINALE PER LA VITA 2018 - REGIONE PUGLIA; 2019 Feb 07. Available online at <https://www.sanita.puglia.it/documents/20182/26673928/Calendario+vaccinale+per+la+vita+2018/f36429b9-5f76-44c2-b495-d5b92289> last accessed on 11 October 2021, at 18:55. 8.World Health Organization (WHO). Causality assessment of an adverse event following immunization (AEFI): user manual for the revised WHO classification, 2nd ed., 2019 update. Available online at <https://www.who.int/publications/i/item/causality-assessment-aefi-user-manu>

Oral presentation

Smith Jennifer
JenniferS@unc.edu - United States

Fan Lee
fan_lee@med.unc.edu -

#3917

Prevalence of high-risk HPV by RNA assay in home self-collected samples among underscreened people in North Carolina

11 - Screening for women difficult to reach

Fan L, Bukowski A, P. Spees L, B. Wheeler S, T. Brewer N, Sanusi B, G. Hudgens M, Jackson S, Barclay L, Carter A, H. Tang J, S. Smith J.

Background/Objectives: Low-income and uninsured people with a cervix (PWC) are at highest risk of being underscreened for cervical cancer. We evaluated the prevalence of high-risk HPV (hrHPV) on home self-collected samples, as well as rates of in-clinic follow-up and risk factors associated with hrHPV positivity in this at-risk population

Methods: My-Body-My-Test-3 was conducted between 2016-2019 in North Carolina among individuals aged 25-64 years, overdue for cervical cancer screening, and with incomes of <250% of the U.S. Federal Poverty Level. Our analytic sample included participants randomized to the self-collection arm who returned self-collected cervico-vaginal brush samples for HPV testing (n=329). Samples were tested for 14 hrHPV types by an HPV RNA assay and further genotyped for HPV 16 and HPV 18/45. We examined behavioral risk factors for hrHPV positivity using logistic regression and between-subjects t-tests.

Results: HrHPV RNA prevalence was 16% (n=52/329) in self-collected samples. Of the hrHPV-positive participants, 24 (46%) presented for in-clinic cervical cancer screening, compared to 56 (20%) of hrHPV-negative participants. Those with ≥ 2 sexual partners in the past year were twice as likely to be hrHPV positive in adjusted analyses (adjusted OR=2.00 (95% CI: 1.03-3.88)). HrHPV-positive and negative participants had similar attitudes towards screening, with the exception of hrHPV-positive participants who reported a lower perceived risk of cervical cancer than those who were hrHPV-negative ($p<.05$).

Conclusions: The hrHPV RNA prevalence was similar to findings in other underscreened PWC in the US. Efforts to reach underscreened PWC is critical for cervical cancer prevention. Future studies aimed at home self-collection should address methods of increasing clinic attendance and completion of treatment among those with HPV-positive results.

Oral presentation

Smith Jennifer
JenniferS@unc.edu - United States

Smith Jennifer
JenniferS@unc.edu - United States

#3918

EXTENDED HUMAN PAPILLOMAVIRUS GENOTYPING TO PREDICT PROGRESSION TO HIGH-GRADE CERVICAL PRECANCER: A PROSPECTIVE COHORT STUDY IN THE SOUTHEASTERN UNITED STATES

03 - Epidemiology and natural history

Smith J, Bukowski A, Hoyo C, Hudgens M, Brewster W, Valea F, Bentley R, Vidal A, Maguire R, Schmitt J, Murphy S, North K

Background/Objectives: High-risk human papillomavirus (hrHPV) testing is recommended for primary cervical cancer screening and is used alongside cytology to triage screening abnormalities to colposcopy. Most screening-based hrHPV tests involve pooled detection of any hrHPV type or of HPV 16/18 only. Extended HPV genotyping, particularly of non-16/18 hrHPV types, may improve risk stratification for millions of cervical abnormalities detected each year, but HPV genotype-specific progression risks are not well characterized. HPV genotype-specific incidence of high-grade cervical intraepithelial neoplasia or more severe (CIN2+) was examined among individuals with normal or low-grade (CIN1) histology following an abnormal screening cytology result.

Methods: A US-based prospective, multiracial cohort of 343 colposcopy referral patients with normal (n=226) or CIN1 (n=117) histology was assessed. Baseline cervical samples underwent HPV DNA genotyping. Participants were followed for up to five years. Genotype-specific CIN2+ incidence rates (IR) were estimated with accelerated failure time models for interval-censored data. Cumulative CIN2+ risk over five years was estimated non-parametrically and compared for HPV risk groups (HPV 16/31/18, else HPV 33/58/52/45, else HPV 39/68/35/51/59/56/66) and age groups (ages 21-24, 25-29, 30+).

Results: At enrollment, median participant age was 30.1 years; 67.1% were hrHPV-positive. During follow-up (median 24.3 months), 24 participants progressed to CIN2+ (7.0%; 6.2% among normal histology and 8.6% among CIN1). CIN2+ IR among hrHPV-positive participants was 3.2/1,000 person-months. CIN2+ IRs were highest for HPV 16 (IR 8.3; 95% CI 4.1, 16.6), HPV 33 (IR 7.8; 95% CI 2.0, 31.3), and HPV 58 (IR 4.9; 95% CI 1.2, 19.6). IRs were highest for HPV 16, 68, and 33 among normal histology and for HPV 33, 31, 16, and 58 among CIN1. Five-year CIN2+ risks were 0.27 for HPV 16/31/18, 0.11 for HPV 33/58/52/45, and 0.13 for HPV 39/68/35/51/59/56/66 (p=0.04). CIN2+ risk did not differ by age group.

Conclusions: In addition to the established HPV type 16, HPV 33 and 58 were consistently predictive of progression to CIN2+ over a five-year period in this cohort. These findings also support the utility of HPV risk groups to stratify women with non-16/18 hrHPV positivity who exhibit differential risks of progression. Additionally, HPV risk groups appear to predict progression regardless of age; thus, although hrHPV testing is not recommended for initial screening of women under 30 years of age, it may be a risk-stratification tool for these younger women, once a cervical abnormality is found. In conclusion, extended genotyping of non-16/18 hrHPV may be useful to identify individuals with normal or low-grade histology results who are at increased risk of progression to CIN2+, thereby improving the efficacy of cervical cancer screening.

Oral presentation

Smith Jennifer
JenniferS@unc.edu - United States

Pretsch Peyton
ppretsch@live.unc.edu - United States

#3931

EFFECT OF HPV SELF-COLLECTION ON CERVICAL CANCER SCREENING COMPLETION AMONG LOW-INCOME, UNDER-SCREENED PEOPLE WITH A CERVIX IN THE US: MY BODY, MY TEST 3 STUDY

11 - Screening for women difficult to reach

Pretsch P, Spees L, Rohner E, Wheeler S, Hudgens M, Sanusi B, Des Marais A, Jackson S, Barclay L, Carter A, Brewer N, Smith J

Background/Objectives: Most cervical cancer cases in the US occur among under-screened people with a cervix. Mailed, at-home HPV self-collection has the potential to reduce barriers and increase screening coverage among under-screened people with a cervix. We conducted a randomized controlled trial to assess the effectiveness of mailing HPV self-collection kits to increase completion of cervical cancer screening among low-income, under-screened people with a cervix.

Methods: My Body My Test-3 study participants came from 22 counties in North Carolina, US. Eligibility included being aged 25-64 years, uninsured or enrolled in Medicaid or Medicare, reporting an income of $\leq 250\%$ of US Federal Poverty Level, and overdue for screening per national guidelines (Pap >4 years or high-risk HPV test >6 years ago). Participants randomized to the intervention arm received an HPV self-collection kit and received scheduling assistance to obtain a free screening appointment. In the control arm, participants received scheduling assistance alone. Self-collection kits included illustrated instructions and a Viba self-collection brush (Rovers). Cervico-vaginal samples were tested for high-risk HPV RNA (Aptima, Hologic). The primary trial outcome was completion of screening within 6 months of study enrollment: defined as testing HPV-negative on self-collected samples (intervention arm) or attending in-clinic screening. Our primary analysis applied an intention-to-treat (ITT) principle to compare screening completion between study arms.

Results: We randomized 665 participants to intervention and control arms in a 2:1 ratio. Median age was 42 (range 25-61) years and median time since last screening was 5 (range 4-25) years. Screening completion was higher in the intervention arm than in the control arm (72% vs. 37%; relative risk [RR]=1.93, 95% confidence interval [CI]: 1.62-2.32) in primary ITT analyses. Among 329 intervention participants who returned a kit with valid test results, 16% were HPV positive. Among all 665 participants, 3.5% had abnormal cytology (4.1% in intervention arm, 2.2% in control), and 1.4% were referred for colposcopy (1.6% in intervention, 0.9% in control). Screening detected CIN2+ in 0.5% of the intervention and none of the control participants.

Conclusions: Mailed HPV self-collection kits with scheduling assistance led to more cervical cancer screening completion among low-income, under-screened people with a cervix as compared to scheduling assistance alone. At-home HPV self-collection testing has the potential to increase screening completion among infrequently screened people with a cervix in the US.

Poster

Duvlis Sotirija
sotirijaduvlis@yahoo.com - Macedonia

Duvlis Sotirija
sotirija.duvlis@ugd.edu.mk - Macedonia

#3932

HPV E6/E7mRNA association with Interleukin - 10 592C/A variant in group of Macedonian women

02 - Viral and molecular biology

Duvlis S., Osmani D.

Background/Objectives: Interleukin 10 (IL-10) is an immunosuppressive cytokine and its genetic variant could have an indirect impact on viral biology and HPV E6/E7 mRNA expression as well. In the study, we evaluate the association between IL10 -592 C/A polymorphism and HPV E6/E7 mRNA expression in a group of women from R North Macedonia.

Methods: Using commercial tests we analyzed 272 women's cervical samples for HPV E6/E7 mRNA and HPV DNA presence respectively. The cases were stratified into three groups: double-positive (n=108, positive for both tests), negative (n=51, negative for HPV E6/E7 mRNA and HPV DNA positive), and the control group (n=113, negative for both tests). The IL10-592 C/A polymorphism was analyzed using polymerase chain reaction-restriction fragment length polymorphism.

Results: The CC genotype and the C allele frequencies of IL10-592C/A were significantly higher in double-positive (59.3% and 78.2%) compared to negative group (39.2% and 65.7%), ($p=0.01$, $CI=0.44;0.22-0.87$ - dominant model; and $p=0.01$, $CI=0.53; 0.3-0.8$) respectively and compared to negative and control groups together

Conclusions: The CC genotype and C allele of IL10-592 showed to be associated with HPV E6/E7 mRNA but not with HPV DNA positivity, which could mean this polymorphism could affect the course of the infection only after HPV onset and it is not associated with susceptibility to HPV.

Oral presentation

Smith Jennifer
JenniferS@unc.edu - United States

Lee Fan
fan_lee@med.unc.edu -

#3933

SELF-COLLECTION TEST PERFORMANCE FOR HIGH-RISK HPV RNA DETECTION AMONG HIV-POSITIVE AND HIV-NEGATIVE WOMEN ENGAGED IN FEMALE SEX WORK IN KENYA

34 - Sexually transmitted diseases and HIV infection

Lee F, Islam J, Mutua M, Kabare E, Manguro G, Waweru W, Mandaliya K, Shafi J, Mcclelland S, Smith J

Background/Objectives: In low and middle-income countries, WHO recommends primary HPV testing for cervical cancer screening where available. Self-collection is also recommended given its potential to overcome screening barriers. We examine self-collected high-risk HPV RNA testing for the detection of histologically confirmed high-grade cervical intraepithelial neoplasia (CIN2/3) among women engaged in female sex work (FSW) in Mombasa, Kenya. We report hrHPV prevalence among HIV-positive and HIV-negative women, and determine predictors of baseline hrHPV infection and CIN2/3 outcomes.

Methods: 400 Kenyan women (193 HIV-positive, 206 HIV-negative) who were ≥ 18 years old, non-pregnant and without a history of treatment for cervical precancer participated (2013-2018). Baseline demographics were obtained at study enrollment. Screening procedures included self-collection of cervico-vaginal via a Viba cytobrush (Rovers) and provider-collection of cervical samples, inspection with acetic acid (VIA) and Pap smear screening. APTIMA HPV Assay (Hologic Corporation) was used to detect E6/E7 oncogene mRNA of 14 hrHPV types. Abnormal results from any screening method and 10% random sample of participants with negative results on all tests underwent colposcopy exam. Cervical biopsies were obtained from lesions, and if no lesions were identified, taken at 12 and 6 o'clock at the squamo-columnar junction. Sensitivity and specificity analyses were used to evaluate test performance for CIN2+ detection. Risk factors for hrHPV positivity and CIN2,3 disease were determined via multi-variable logistic regression.

Results: 400 Kenyan women (193 HIV-positive, 206 HIV-negative) who were ≥ 18 years old, non-pregnant and without a history of treatment for cervical precancer participated (2013-2018). Baseline demographics were obtained at study enrollment. Screening procedures included self-collection of cervico-vaginal via a Viba cytobrush (Rovers) and provider-collection of cervical samples, inspection with acetic acid (VIA) and Pap smear screening. APTIMA HPV Assay (Hologic Corporation) was used to detect E6/E7 oncogene mRNA of 14 hrHPV types. Abnormal results from any screening method and 10% random sample of participants with negative results on all tests underwent colposcopy exam. Cervical biopsies were obtained from lesions, and if no lesions were identified, taken at 12 and 6 o'clock at the squamo-columnar junction. Sensitivity and specificity analyses were used to evaluate test performance for CIN2+ detection. Risk factors for hrHPV positivity and CIN2,3 disease were determined via multi-variable logistic regression.

Conclusions: Self-collection for hrHPV testing performed similarly to provider-collection in this high-risk cohort. The relatively higher sensitivity of mRNA-based hrHPV testing compared to cytology for the detection of clinically relevant cervical disease is important in this high-risk cohort.

Oral presentation

Dr Lefevre Caroline
caroline.lefeuvre@chu-angers.fr - France

Lefevre Caroline
caroline.lefeuvre@chu-angers.fr - France

#3938

URINARY HPV DNA TESTING AS A TOOL FOR CERVICAL CANCER SCREENING IN FRENCH WOMEN

13 - Self-sampling

Lefevre C, Pivert A, Le Duc Banaszuk A, Ducancelle A

Background/Objectives: Despite population-based campaigns including invitation letters sent to women recommending them to contact a physician to take a cervical sample, cervical cancer screening coverage remains moderate (~60%) in France. We conducted the CapU1 study in France (Maine-et-Loire department) on women who did not undergo a Pap-smear during the last three years. This study demonstrated that urine self-sampling increases patient compliance and can be relevant to detect high grade cervical lesions (1). Because of these promising results and in order to extend the screening coverage, we led a third study named CapU3 (2). This study sent approximately 13 000 urine collection kits to women living in the Department of Maine-et-Loire, aged 35-65 years, who did not have screening record since seven years ago or longer. The participation rate was 15.4% and good gynecological follow-up after a positive urinary HPV testing was observed (92.1 %). However, the CapU3 trial did not include a comparison arm impeding to demonstrate evidence of comparative efficacy. Since 2019, the French health authority specifies that vaginal self-sampling is possible for women who are not regularly screened. The collection of a urine specimen is less invasive for women reluctant to undergo gynaecological sampling (cervical sampling or vaginal self-sampling) and may generate higher participation rates. Consequently, the aim of the CapU4 study is to evaluate the effectiveness of two experimental invitation strategies (urine or vaginal self-sampling) to reach under-screened populations and compare them to the current invitation strategy in two rural medically under-served departments, in France, with low screening coverage.

Methods: The CapU4 study is a randomized controlled trial with three arms. The target population (15 000 women) includes women aged 30-65 years, living in two rural departments with low medical density, who had no screening test recorded since more than four years and who did not respond to an invitation letter within twelve months before. A team of psychologists will investigate attitudes and experiences by semi-structured/focus-group interviews with voluntary CapU4 participants and with health professionals.

Results: This study, funded from the French National Institute, will begin in February 2022.

Conclusions: The Capu4 study will for the first time compare directly two experimental invitation strategies (offering self-sampling of vaginal specimens or collection of first-void urine) with the current procedure including traditional reminder letters. CapU4 will identify effective strategies to reach women not responding to current screening invitations and will generate information about acceptance of self-sampling among women and health professionals.

References: 1. Ducancelle A, Reiser J, Pivert A, Le Guillou-Guillemette H, Le Duc-Banaszuk AS, Lunel-Fabiani F. Home-based urinary HPV DNA testing in women who do not attend cervical cancer screening clinics. J Infect. 2015;71(3):377-84. 2. Lefevre C, Pivert A, Guillou-Guillemette HL, Lunel-Fabiani F, Veillon P, Le Duc-Banaszuk A-S, et al. Urinary HPV DNA testing as a tool for cervical cancer screening in women who are reluctant to have a Pap smear in France. J Infect. 2020;81(2):248-54.

Oral presentation

Prof. Dr Wieland Andreas
andreas.wieland@osumc.edu - United States

Wieland Andreas
andreas.wieland@osumc.edu - United States

#3940

Defining HPV-specific B cell responses in head and neck cancer patients

30 - HPV and oropharynx / Head and neck cancer

Wieland A

Background/Objectives: Tumours can harbour significant numbers of B cells and plasma cells, however, little is known about the antigen specificity of intra-tumoral B cells.

Methods: Using a wide range of methodologies including ELISPOT, flow cytometry, scRNAseq, and multiplex IHC, we characterized human papillomavirus (HPV)-specific B cell responses in the tumor microenvironment (TME) of HPV+ head and neck cancer patients.

Results: We show that HPV-specific B cell responses are detectable in HPV-positive head and neck cancers, with active production of HPV-specific IgG antibodies in situ. HPV-specific antibody secreting cells (ASCs) were readily detectable (~0.7-25% of IgG+ ASCs) in the TME with minimal bystander recruitment of influenza-specific cells, suggesting a localised and antigen-specific ASC response. HPV-specific ASC responses, which correlated with plasma IgG titres, were directed against the HPV proteins E2, E6, and E7, with E2-specific responses tending to be the most dominant. Using intra-tumoral B cells and plasma cells, we generated several HPV-specific human monoclonal antibodies, which exhibited a high degree of somatic hypermutation, consistent with chronic antigen exposure. scRNAseq analyses revealed the presence of activated B cells, germinal centre B cells, and ASCs within the TME. ASCs and B cells were preferentially localised in the tumour stroma, with well-formed clusters of activated B cells indicating ongoing germinal centre reactions.

Conclusions: Overall, we show that antigen-specific activated and germinal centre B cells as well as plasma cells can be found in the TME. Our findings provide a better understanding of humoral immune responses in human cancer and suggest that tumour-infiltrating B cells could be harnessed for the development of novel therapeutic agents.

Oral presentation

Mrs Marini Bruna
b.marini@ulissebiomed.com - Italy

Marini Bruna
b.marini@ulissebiomed.com - Italy

#3942

Clinical validation of full genotyping HPV Selfy assay according to the international guidelines for HPV test requirements for cervical cancer screening on clinician collected and self-collected samples.

14 - Genotyping

Marini B, Avian A, Clemente N, Mauro E, Isidoro E, Di Napoli M, Dudine S, Del Fabro A, Morini S, Perin T, Giudici F, Cammisuli T, Foschi N, Mocenigo M, Montrone M, Modena C, Polenghi M, Puzzi L, Valenti G, Sola R, Zanolli S, Vogrig E, Riva E, Ciccozzi M, Angeletti S, Castriciano S, Pachetti M, Zella D, Petti M, Centonze S, Gerin D, Banks L, Canzonieri V, Sopracordevole F, Zanconati F, Ippodrino R

Background/Objectives: The study aims at evaluating the diagnostic performance of HPV Selfy, a PCR-based DNA test for the detection and full genotyping of 14 high-risk Human Papillomavirus, in a single reaction, on clinician-collected cervical specimen and self-collected vaginal samples in a primary cervical cancer screening population, according to international guidelines.

Methods: In 2018 and 2019 women consecutively referring to Azienda Sanitaria Integrata Giuliano Isontina (ASUGI - Trieste, Italy) and to CRO - National Cancer Institute (Aviano, Italy) for the Italian cervical cancer screening program were invited to participate to the study. Eligible women were asked to perform both a vaginal self-sampling analyzed with HPV Selfy (Ulisse BioMed) and a clinician-collected cervical brush for Pap smear, analyzed both with the Hybrid Capture® 2 DNA test (HC2) (Qiagen) and HPV Selfy. As required by guidelines, a non-inferiority test was conducted to compare the clinical performance of HPV Selfy with the reference test. Meijer's guidelines for validation of HPV test for primary cervical cancer screening, and VALHUDES protocol for validation of the assay on self-collected samples.

Results: The study demonstrates that HPV Selfy clinical sensitivity and specificity are non-inferior to those of HC2. In particular, by analysis of a total of 889 cervical liquid-based cytology specimens from a cervical cancer screening population, of which 98 were from women with histologically diagnosed high grade cervical lesion (CIN2+), HPV Selfy showed relative sensitivity and specificity for CIN2+ of 0.98 and 1.00 respectively (non-inferiority score test: $P=0.01747$ and $P=0.00414$, respectively). In addition, an adequate intra- and inter-laboratory reproducibility has been reached by the test. Moreover, we demonstrated that the performance of HPV Selfy on self-collected vaginal swabs was non-inferior to the performance obtained on clinician-collected cervical brushes (0.92 relative sensitivity and 0.97 relative specificity). Finally, exploiting the genotyping capability of HPV Selfy test, we described HPV types prevalence in the study population. The most frequent high-risk HPV types detected in the cytology-negative screening population were HPV31 and HPV58, followed by HPV59. On the other hand, in the CIN2+ population, HPV16 was the predominant strain followed by HPV31 and HPV58.

Conclusions: In conclusion, this study demonstrates that the full genotyping test HPV Selfy is clinically non-inferior to HC2. The clinical performance and reproducibility of the assay meet the international criteria for validation of a high-risk HPV DNA test for cervical cancer screening purposes according to Meijer's guidelines. Moreover, we demonstrated that HPV Selfy is also validated on self-collected samples for primary screening purposes, satisfying the requirements indicated by VALHUDES protocol. The study confirmed that self-sampling could be an attractive solution to increase women's participation to cervical cancer screening programs. Self-sampling could be particularly relevant nowadays, when COVID-19 pandemic deeply impacted the effectiveness of hospital-based organized screening programs. In addition, HPV Selfy provides also full genotyping information in the same PCR reaction, that can be valuable for future triage strategies development since significant differences have been observed in the oncogenicity among different high risk HPV genotypes.

Oral presentation

Prof. Cuzick Jack
j.cuzick@qmul.ac.uk - United Kingdom

Cuzick Jack
j.cuzick@qmul.ac.uk - United Kingdom

#3943

Role of HPV Genotyping in Refining Cervical Screening

10 - HPV screening

Background/Objectives: See accompanying email for this

Methods: as above

Results: as above

Conclusions: as above

Oral presentation

Dr Puram Sidharth
sidpuram@wustl.edu - United States

Puram Sidharth
sidpuram@wustl.edu - United States

#3949

Cellular states are coupled to genomic and viral heterogeneity in HPV-related oropharyngeal carcinoma

30 - HPV and oropharynx / Head and neck cancer

Puram S

Background/Objectives: Define the heterogeneous cell states present in HPV-related oropharyngeal cancer and determine their impact on disease pathogenesis.

Methods: Single cell RNA-sequencing of HPV-related and HPV-negative oropharyngeal tumors in a prospectively collected cohort (n=15).

Results: We uncovered a high level of cellular diversity within and between tumors. First, we detect diverse chromosomal aberrations within individual tumors, suggesting remarkable genomic instability and enabling the identification of malignant cells even at pathologically-negative margins. Second, we uncover diversity with respect to HNSCC subtypes and other cellular states such as the cell cycle, senescence, and epithelial-mesenchymal transitions. Third, we find diversity in the expression of viral genes within HPV-positive tumors. These findings suggest that expression diversity in HPV oropharyngeal tumor must be taken into account during diagnosis and treatment of HPV-positive tumors, with important ramifications for uncovering patients at risk for poor outcomes among an otherwise largely favorable prognosis.

Conclusions: In summary, our comprehensive single cell analysis of oropharyngeal tumors reveals unappreciated diversity both in host and in viral genes, with critical implications for our understanding of virally-mediated tumors.

Oral presentation

Dr Robbins Hilary
RobbinsH@iarc.fr - France

Robbins Hilary
robbinsh@iarc.fr - France

#3952

Absolute risk of oropharyngeal cancer after an HPV16-E6 serology test and implications for screening

30 - HPV and oropharynx / Head and neck cancer

Robbins H

Background/Objectives: The incidence of HPV-associated oropharyngeal cancer (OPC) is rising and earlier detection could be beneficial. The HPV16-E6 blood-based biomarker has high sensitivity and specificity for HPV-driven OPC, but the absolute risk of OPC following a positive or negative HPV16-E6 test is unknown.

Methods: We constructed an OPC risk prediction model that integrates data for (1) relative odds of OPC for HPV16-E6 serostatus, cigarette smoking, and sex from the HPV Cancer Cohort Consortium (HPVC3); (2) US population risk factor data from the National Health Interview Survey, and (3) US population OPC incidence and mortality.

Results: The 9 HPVC3 cohorts included 365 OPC cases with up to 10 years between blood draw and diagnosis and 5,794 controls. The estimated 10-year OPC risk for HPV16-E6 seropositive males at age 50 years was 16.7% (95%CI=10.5%-29.4%) and at age 60 was 25.0% (95%CI=15.5%-44.6%). Corresponding 5-year risk estimates were 7.0% and 13.3%, respectively. For HPV16-E6 seropositive females, 10-year risk estimates were 4.8% (95%CI=1.4%-11.4%) at age 50 and 7.5% (95%CI=2.3%-17.4%) at age 60, and 5-year risk estimates were 2.0% and 3.8%, respectively. Over 30 years, after a seropositive result at age 50, an estimated 48.1% of men and 16.6% of women would develop OPC. In contrast, 10-year risks among HPV16-E6 seronegative people were very low, ranging from 0.01% to 0.23% depending on age, sex, and smoking status.

Conclusions: We estimate that a substantial proportion of HPV16-E6 seropositive individuals will develop OPC, with 10-year risks of 17-25% for males and 5-8% for females aged 50-60 years in the US. The high risk of OPC among HPV16-E6 seropositive individuals may warrant periodic, minimally invasive surveillance, particularly for males, in regions where incidence rates reach levels that warrant such intervention. Establishing an appropriate protocol for this surveillance is an important direction for future research. This could involve evaluation of blood-based biomarkers and imaging techniques.

Oral presentation

Prof. Dr Steenbergen Renske
r.steenbergen@amsterdamumc.nl - Netherlands

Steenbergen Renske
r.steenbergen@amsterdamumc.nl - Netherlands

#3954

Methylation analysis of anal cancer and pre-cancerous lesions- implications for cancer risk stratification of anal HSIL

28 - Anal neoplasia

Background/Objectives: High-grade anal intraepithelial neoplasia (HGAIN; AIN2/3) is highly prevalent in HIV+ men, but only a minority will progress towards cancer. Since the risk of progression cannot be established, current screening options result in substantial over-referral and overtreatment. Therefore risk stratification of HGAIN using biomarkers is needed.

Methods: Host cell DNA methylation markers for detection of HGAIN and anal cancer were evaluated in a cross-sectional series of 148 anal tissue samples of HIV+ men using quantitative methylation-specific-PCR. The best performing markers were validated in a large, independent cross-sectional series of 345 HIV+ samples and also evaluated in a cross-sectional series of 176 samples of HIV-negative women and men. Accuracy for detection of AIN3 and cancer was determined by univariable and multivariable logistic regression analysis, followed by leave-one-out-cross-validation. The association with cancer progression was assessed in a longitudinal series of ten anal cancer cases with preceding HGAIN at similar anatomic locations.

Results: In all series of anal biopsies tested methylation marker levels increased with increasing severity of disease ($p < 0.05$). A high resemblance was found between all patient groups. Histopathologically similar HGAIN revealed a heterogeneous methylation pattern, a subset resembling cancer. In HIV+ MSM a three-marker panel ASCL1, SST, ZNF582 was most accurate (AUC=0.89) in distinguishing AIN3 and cancer from controls, not missing cancers. AIN3+ detection using these markers in HIV-negative women and men was comparable (AUC=0.85). In the longitudinal series, all HGAIN preceding cancer displayed high methylation levels similar to cancers.

Conclusions: We identified and validated methylation markers for the detection of anal (pre-)cancer in HIV+MSM. Accurate AIN3+ detection was confirmed in HIV-negative patients. High methylation levels were associated with progression to cancer. Therefore, these methylation markers provide a promising cancer risk stratification tool for all risk groups of anal cancer. Methylation testing can identify HGAIN in need of treatment and prevent overtreatment of HGAIN with a low cancer risk. A prospective clinical validation study has recently been initiated.

References: van der Zee, RP, Richel, O, van Noesel, CJM, Novianti, PW, Ciocanea-Teodorescu, I, van Splunter, AP, Duin, S, van den Berk, GEL, Meijer, CJLM, Quint, WGV, de Vries, HJC, Prins, JM & Steenbergen, RDM 2019, 'Host Cell Deoxyribonucleic Acid Methylation Markers for the Detection of High-grade Anal Intraepithelial Neoplasia and Anal Cancer', *Clinical Infectious Diseases*, vol. 68, no. 7, pp. 1110-1117 van der Zee, RP, Richel, O, van Noesel, CJM, Ciocănea-Teodorescu, I, van Splunter, AP, Ter Braak, TJ, Nathan, M, Cuming, T, Sheaff, M, Kreuter, A, Meijer, CJLM, Quint, WGV, de Vries, HJC, Prins, JM & Steenbergen, RDM 2021, 'Cancer risk stratification of anal intraepithelial neoplasia in HIV-positive men by validated methylation markers associated with progression to cancer', *Clinical infectious Diseases*. 72, 12, p. 2154-2163 van der Zee, RP, van Noesel, CJM, Martin, I, Ter Braak, TJ, Heideman, DAM, de Vries, HJC, Prins, JM & Steenbergen, RDM 2021, 'DNA methylation markers have universal prognostic value for anal cancer risk in HIV-negative and HIV-positive individuals', *Molecular oncology*. Nov;15(11):3024-3036. van der Zee, R. P., Meijer, C. J. L. M., Cuming, T., Kreuter, A., van de Sandt, M. M., Quint, W. G. V., de Vries, H. J. C., Prins, J. M. & Steenbergen, R. D. M. 2021, Characterisation of anal intraepithelial neoplasia and anal cancer in HIV-positive men by immunohistochemical markers p16, Ki-67, HPV-E4 and DNA methylation markers, *Int J. of Cancer*. 149, 10, p. 1833-1844

Oral presentation

Iftner Thomas
thomas.iftner@med.uni-tuebingen.de - Germany

Iftner Thomas
thomas.iftner@med.uni-tuebingen.de - Germany

#3955

HPV-TESTANFORDERUNGEN FÜR DAS ORGANISIERTE GEBÄRMUTTERHALSKREBS-SCREENING PROGRAMM IN DEUTSCHLAND

10 - HPV screening

Iftner T

Background/Objectives: Die Richtlinie für das organisierte Gebärmutterhalskrebs-Screening-Programm (oKFE) in Deutschland bietet Frauen ab einem Alter von 35 Jahren alle drei Jahre einen HPV-Test in Kombination mit einer Zytologie an. Frauen ab 20 Jahren bekommen weiterhin jährliche PAP-Abstriche und zwischen 30-35 Jahren einen HPV-Test als Triage. Ein HPV-DNA-Test im oKFE sollte nur die 13 HPV-Typen nachweisen, die von der IARC/WHO als krebserregend (class I/IIa) für den Menschen eingestuft wurden. Die klinische Sensitivität und Spezifität eines HPV-Tests für den Nachweis von CIN2+ darf nicht unter 95% bzw. 98% bereits etablierter und in RCTs verwendeter HPV-Testsysteme wie dem HC2-Test liegen. Außerdem sollte die Positivitäts-Rate bei Frauen mit unauffälliger Zytologie nicht höher sein als die des HC2-Tests. Die Reproduzierbarkeit innerhalb eines Labors und zwischen verschiedenen Laboren mit unterschiedlichen Geräten und Mitarbeitern sollte mindestens 90% betragen. Da die oKFE Screening-Intervalle von 3 Jahren vorsieht, sollte sich die longitudinale kumulative Inzidenzrate von CIN2+ und der negative Vorhersagewert (NPV; für mindestens 3 Jahre) nicht signifikant von dem Goldstandard HC2 unterscheiden. Darüber hinaus sollte der gewünschte Test in Populations-basierten Studien überzeugende Leistungen gezeigt haben, vom Labor und den überweisenden Gynäkologen akzeptiert werden und kosteneffektiv sein.

Methods: n.a.

Results: n.a.

Conclusions: n.a.

Oral presentation

Prof. Wilson Allan
allan.wilson@lanarkshire.scot.nhs.uk - United
Kingdom

Wilson Allan
allan.wilson@lanarkshire.scot.nhs.uk - United
Kingdom

#3956

HPV primary screening: the Scottish approach

10 - HPV screening

Wilson A

Background/Objectives: This presentation will describe the introduction of HPV primary screening to the screening programme in Scotland, our unique experience to date and some early data on the effectiveness of the new primary test within a Scottish context.

Methods: The unique elements of the introduction of HPV primary screening will be described focussing on the IT challenges, procurement of the HPV platforms and the impact of the pandemic on the "go live" date and early months of the new service. Early data on the performance of the primary HPV test and the cytology triage will be presented. The programme uses focussed co-testing in a series of pathways designed to use cytology and HPV testing where it is deemed clinically appropriate

Results: The data so far has been difficult to compare with cytology as the primary test due to the disruption to the programme caused by the pandemic. However, early data suggests an increased detection of disease and less impact on colposcopy than the earlier modelling indicated. The re-designed laboratory service appears to be working well and data on the performance of both labs in HPV testing and cytology triage will be presented.

Conclusions: Despite years of planning the impact of the pandemic provided a significant challenge to the introduction of HPV primary screening in the Scottish screening programme. The new service adapted well in the face of the issues the pandemic presented and due to the commitment of the staff across the programme the re-designed programme is now well established and has 2 years of data to analyse. The Scottish IT system is one of the strengths of the screening programme and has underpinned the move to HPV primary. The introduction of HPV primary screening in Scotland has several unique aspects and these will be described during this presentation.

Oral presentation

Van Der Burg Sjoerd
s.h.van_der_burg@lumc.nl - Netherlands

Sjoerd Van Der Burg
shvdburg@lumc.nl - Netherlands

#3957

HPV-specific T cells sustain a chemokine-driven spatial organization of intratumoral immune microaggregates required for HPV16+ OPSCC long survival

30 - HPV and oropharynx / Head and neck cancer

Background/Objectives: The composition of the tumor immune microenvironment (TIME) associated with good prognosis generally also predicts the success of immunotherapy, and both entail the presence of pre-existing tumor-specific T cells. Here, the blueprint of the TIME associated with such an ongoing tumor-specific T cell response was dissected in a unique prospective oropharyngeal squamous cell carcinoma (OPSCC) cohort, in which tumor-specific tumor-infiltrating T cells were detected (IR+) or not (IR-).

Methods: A comprehensive multi-modal, high-dimensional strategy was applied to dissect the TIME of treatment naive IR+ and IR- OPSCC tissue, including bulk RNA sequencing by NanoString, imaging mass cytometry (Hyperion) for phenotyping and spatial interaction analyses of immune cells, and combined single-cell gene expression profiling and T cell receptor sequencing (scRNAseq) to characterize the transcriptional states of clonally expanded tumor-infiltrating T cells.

Results: IR+ patients had an excellent survival during >10 years follow-up. The tumors of IR+ patients expressed higher levels of genes strongly related to IFN γ signaling, T-cell activation, T-cell receptor signaling, and mononuclear cell differentiation as well as genes involved in several immune signaling pathways, than IR- patients. The top differently overexpressed genes included CXCL12 and LTB, involved in ectopic lymphoid structure development. Moreover, scRNAseq not only revealed that CD4+ T cells were the main producers of LTB, but also identified a subset of clonally expanded CD8+ T cells, dominantly present in IR+ tumors, which secreted the T cell and dendritic cell (DC) attracting chemokine CCL4. Indeed, immune cell infiltration in IR+ tumors is stronger, highly coordinated and has a distinct spatial phenotypic signature, characterized by intratumoral microaggregates of CD8+CD103+ and CD4+ T cells with DCs. In contrast, the IR- TIME comprised spatial interactions between lymphocytes and various immunosuppressive myeloid cell populations. The impact of these chemokines on local immunity and clinical outcome was confirmed in an independent The Cancer Genome Atlas OPSCC cohort.

Conclusions: The production of lymphoid cell attracting and organizing chemokines by tumor-specific T cells in IR+ tumors constitutes a positive feedback loop to sustain the formation of the DC-T-cell microaggregates and identifies patients with excellent survival after standard therapy and potentially responsive to neoadjuvant immunotherapy.

References: 1 Z. Abdulrahman, S.J. Santegoets, G. Sturm, P. Charoentong, M.E. IJsselsteijn, A. Somarakis, T. Höllt, F. Finotello, Z. Trajanoski, S.L. van Egmond, D.A.M. Mustafa, M.J.P. Welters, N.F.C.C. de Miranda, S.H. van der Burg (2022) Chemokine-driven spatial organization of immune cell microaggregates marks oropharyngeal squamous cell carcinomas containing tumor-specific T cells. J. Immunother. Cancer, accepted (IF 13.751)

Oral presentation

Prof. Dr Steenbergen Renske
r.steenbergen@amsterdamumc.nl - Netherlands

Steenbergen Renske
r.steenbergen@amsterdamumc.nl - Netherlands

#3959

Epigenetics and methylation markers for CIN3+ detection

15 - Molecular markers

Background/Objectives: Disruption of DNA methylation patterns is one of the hallmarks of cancer, including high-risk human papillomavirus (hrHPV)-driven anogenital cancers. Several hyper-methylated cellular genes have been found associated with carcinogenesis of the uterine cervix or other anogenital sites. We assessed the value of host cell DNA methylation markers to serve a triage markers in primary HPV screening and to enable the detection and risk assessment of HPV-induced precancerous lesions.

Methods: Using targeted and genome wide approaches various host cell genes that become methylated during HPV-induced carcinogenesis have been defined. Selected methylation markers were tested and validated in HPV-positive cervical scrapes and self-samples for CIN2/3 and cancer detection.

Results: Methylation markers detect virtually all cervical cancers. In CIN2/3 with varying cancer risks, distinct methylation patterns were seen. Our data indicate that so-called advanced CIN lesions display a cancer-like methylation-high pattern and have a high short-term risk of progression to cancer.

Conclusions: In conclusion, methylation marker analysis provides an attractive triage tool for hrHPV-positive women, with the advantage of applicability to self-collected specimens allowing for full molecular cervical screening. Methylation markers lead to detection of advanced HPV-induced precancerous lesions in need of treatment and can prevent overtreatment of lesions with a low cancer risk.

Oral presentation

Dr Pere Hélène
helene.pere@aphp.fr - France

Péré Hélène
helene.pere@aphp.fr - France

#3960

How droplet digital PCR improves HPV Ct DNA detection?

30 - HPV and oropharynx / Head and neck cancer

Background/Objectives: HPV circulating (Ct) DNA can thus be used as a biomarker that is specific to HPV-positive malignant cells. Indeed, HPV life cycle is strictly epithelial (no viremia) and several studies have shown that HPV DNA is absent in the blood of women with cervical HPV infection, whereas it is detectable in patients with cervical invasive lesion. This DNA is released by cancer cells, as confirmed by sequencing studies that have demonstrated that the HPV type detected in the plasma was identical to that found in the cervical tumor from the same patients. Numerous recent studies have shown that at the time of diagnosis, the great majority of patients affected by an HPV-positive cancer (regardless of the anatomic site - eg. oropharyngeal, cervical or anal) have detectable level of HPV DNA in their plasma whereas controls (healthy individuals or non HPV-related cancers) do not or exceptionally.

Methods: New ultrasensitive technology such as droplet-based digital PCR (ddPCR) or NGS have shown their performance to monitor HPVctDNA. ddPCR allows to generate quantitative and accurate data without standard curves and also present higher sensitivity compared to conventional quantitative PCR (qPCR). Indeed, this method is based on the realization of millions of single-molecule PCRs in parallel in independent compartment (here droplets of an emulsion) and consequently avoids the bias seen in conventional PCR.

Results: The resulting amplification products thus reflect more closely the original composition of nucleic acid mixtures than conventional PCR. In addition to a higher sensitivity over qPCR, ddPCR also demonstrated substantially higher resistance to a variety of inhibitors that could be contained within biological samples and improved comparison between results in different laboratory thanks to the high reproducibility of the obtained data. The increased tolerance of ddPCR to PCR inhibitory molecules makes it an attractive alternative to qPCR for medical applications including viral diagnostics and cancer research. Finally, ddPCR offers an optimized approach for the sensitive detection and quantification of low-target-abundance biological samples.

Conclusions: Thanks to all its benefits, ddPCR represents a technique of choice for monitoring HPV ctDNA in HPV driven cancer patients.

Oral presentation

Heideman Daniëlle
dam.heideman@amsterdamumc.nl - Netherlands

Heideman Daniëlle
dam.heideman@amsterdamumc.nl - Netherlands

#3961

Introduction methylation assays for clinical use

18 - Methylation

Heideman D

Background/Objectives: DNA methylation is an emerging biomarker for the management of HPV-positive women in cervical screening and for risk stratification of women diagnosed with premalignant disease. Aberrant DNA methylation of human genes and HPV genome has been found associated with carcinogenesis of the uterine cervix. CIN2/3 lesions represent heterogeneous disease entities that harbor lower and higher risks for cancer progression. The ability to define lesions with potential short-term risk of progression to cancer by a tumor-like, or high-risk methylation pattern is gaining increased attention. Detection of methylated DNA can be performed by molecular methods on different sample types, including cervical tissue, cervical scrapes, self-collected vaginal samples and urine. Methylation assays could be valuable for enhanced screening, targeted prevention and to reduce overtreatment. This presentation will give an introduction to the clinical applications of methylation assays that will be highlighted in this session.

Methods:

Results:

Conclusions:

References:

Oral presentation

Ms Verhoef Lisanne
l.verhoef@amsterdamumc.nl - Netherlands

Verhoef Lisanne
l.verhoef@amsterdamumc.nl - Netherlands

#3962

Cellular methylation markers in the CIN triage of HPV-vaccinated women

18 - Methylation

Verhoef L

Background/Objectives: HPV vaccination programs will impact the choice of optimal screening strategies. Owing to the reduced incidence of cervical (pre)cancer in vaccinated women, the predictive value of a positive screening result may decrease and the proportion of false-positive results, and associated unnecessary referrals and costs, could increase, if screening programs are not adapted to vaccinated women. Host cell DNA methylation markers (e.g., FAM19A4, miR124-2) provide a molecular means to identify women with cervical cancer and advanced high-grade CIN. Host cell DNA methylation analysis has the ability to discriminate CIN2/3 lesions with a high short-term risk of progression to cancer by a tumor-like, or high-risk methylation pattern. This presentation will discuss the potential of host cell methylation markers in the context of HPV-vaccinated women.

Methods:

Results:

Conclusions:

References:

Oral presentation

Dr Bleeker Maaïke
mcg.bleeker@amsterdamumc.nl - Netherlands

Bleeker Maaïke
mcg.bleeker@amsterdamumc.nl - Netherlands

#3963

Cancer risk and biomarkers in high-grade vulvar intraepithelial neoplasia

15 - Molecular markers

Background/Objectives: Cancer risk and biomarkers in high-grade vulvar intraepithelial neoplasia High-grade vulvar intraepithelial neoplasia (HG-VIN), the precursor of vulvar carcinoma, can be divided into HPV-associated HSIL and HPV-independent and lichen sclerosus (LS)-associated dVIN. HSIL is much more commonly diagnosed than dVIN and the cancer risk is relatively low. In contrast, timely and accurate diagnosis of dVIN is difficult, while the cancer risk is high. The use of immunohistochemical biomarkers and HPV assays can improve the identification of HSIL and dVIN. Host cell DNA methylation markers can identify vulvar lesions with a high cancer risk, which are promising for refining diagnostics and tailored management in affected women.

Methods:

Results:

Conclusions:

References:

Oral presentation

Lei Jiayao
jiayao.lei@ki.se - Sweden

Lei Jiayao
jiayao.lei@ki.se - Sweden

#3964

COUNTRY-SPECIFIC UPDATES ON HPV VACCINE EFFECTIVENESS AGAINST CERVICAL CANCER: SWEDEN

06 - HPV prophylactic vaccines

Lei J, Sparén P

Background/Objectives: Several studies have demonstrated the population effectiveness of quadrivalent HPV (qHPV) vaccination against genital warts, cervical precancerous lesions and invasive cervical cancer (ICC). We aimed to re-examine the effectiveness of qHPV vaccination against ICC with a longer follow-up period.

Methods: We conducted an updated analysis using the same methods as in our previous paper. We investigated the effectiveness of HPV vaccination against ICC in a national cohort study of 1,775,503 women aged 10-31 during 2006-2018 (we aim to further extend the follow-up to 2020 for the actual presentation, data application pending) utilizing Swedish registries. Girls and women that had at least one dose of qHPV vaccination were considered as vaccinated, and HPV vaccination was considered as a time-varying exposure. We assessed the association between HPV vaccination and the risk of ICC, controlling for age at follow-up, calendar year, county of residence, and parental characteristics, including education, household income, mother's country of birth, and maternal disease history. Incidence rate ratios (IRR) of ICC, with 95% confidence intervals (CI), were estimated in Poisson regression models.

Results: A total of 572,180 women (26.9%) had at least one dose of HPV vaccination. During the study period 28 HPV vaccinated women acquired ICC before age 32, while the corresponding number for unvaccinated women was 701, yielding an incidence rate of 0.90 and 6.19 per 100 000 person-year, respectively. The IRR for HPV vaccinated vs. unvaccinated women was 0.53, (95% CI 0.36 - 0.77) after adjustment of age and 0.38, (95% CI 0.26 - 0.57) after adjusting for other covariates. The IRR for women vaccinated before age 17 and age 17-30 were 0.23, (95% CI 0.10 - 0.53) and 0.45, (95% CI 0.29 - 0.70), respectively, after adjusting for all covariates.

Conclusions: In this updated analysis with extended follow-up period, qHPV vaccination was associated with a substantially reduced risk of invasive cervical cancer at the population level.

Oral presentation

Mr Forslund Ola
ola.forslund@med.lu.se - Sweden

Forslund Ola
ola.forslund@med.lu.se - Sweden

#3965

Self-sampling within routine cervical cancer screening in Region of Skåne, Sweden

10 - HPV screening

Forslund O, Borgfeldt C

Background/Objectives: Region Skåne is one of 21 regions in Sweden. The region has about 1.4 million inhabitants, and cervical cancer screening is organized by Public Health. About 400 000 women are in the screening ages between 23-70 years and all samples are analyzed at one laboratory. Year 2017 primary HPV screening (mRNA assay) was introduced for women 30-70 years old. Recently in September year 2021, vaginal self-sampling was commenced as the first sample option for women aged 23-70 years. The aim of the presentation is to give an overview of the characteristics and performance of the newly introduced self-sampling program.

Methods: Organization: The organization of the self-sample screening program will be described via interviews of key informant personnel at Laboratory Medicine Skåne. The qualitative performance of the self-sample program will be analyzed by the use of data from the cytology/pathology database. HPV testing: Self-sampling kits (Multitest Swab Specimen Collection kit, Hologic) are sent by mail to the women. The kit contains information along with the necessary equipment. The self-sample is returned in a prepaid envelope to the Department of Pathology in Lund. Self-samples are heated at 90°C for 1h and 15 min, and cooled to room temperature before being analysed by Aptima HPV mRNA. Self-samples with invalid results from the HPV mRNA assay are re-tested in duplicate and scored as valid if at least one of the results are valid. Self-sample-HPV-positive women are then invited for a follow-up within 3 months, where liquid based cytology (LBC) cervical samples are collected by midwives. Renewed HPV analysis are performed and reflex cytology are made from HPV positive samples. Further gynaecological follow-up due to cytology of >ASCUS+ is done at outpatient colposcopy clinics, where appropriate histological specimens are taken and sent for pathological assessment at Department of Clinical Pathology in Lund. Non-attendees for the follow-up procedure with a midwife are sent a reminder by regular mail. Estimated costs for the self-sampling program will be compared to that of the previous primary HPV-screening approach of LBC samples. We will include costs from the laboratory and midwives, such as cost for self-samples kits sent out by post, HPV-analyses, follow ups at midwives of self-sample HPV-positive women with LBC-sampling and renewed HPV testing and reflex cytology.

Results: At the conference an overview of the organization of self-sample screening program will be described. In addition, the outcome of self-sampling program will be given as well as the estimated cost of the program.

Conclusions: Will be given at the conference.

Oral presentation

Dr Dick Stéfanie
s.dick@amsterdamumc.nl - Netherlands

Dick Stéfanie
s.dick@amsterdamumc.nl - Netherlands

#3966

Use of methylation markers as guidance for CIN2/3 management

15 - Molecular markers

Dick S

Background/Objectives: Many CIN2/3 lesions, especially in young women of reproductive age, show spontaneous regression. Since surgical treatment is associated with cervical morbidity and preterm birth, overtreatment should be avoided. A test that can predict clinical regression of CIN2/3 lesions may help to prevent overtreatment and is urgently needed. Current histopathological grading of CIN by pathologists cannot discriminate between regressive and non-regressive CIN2 and CIN3. Several biomarkers have been evaluated for their prognostic value such as HPV viral load and immunohistochemical staining of p16INK4A and Ki-67, but none of these were able to predict regression or progression of CIN. Recent studies have shown that DNA methylation levels of host cell genes and viral genes increase with increasing severity of CIN grade and are extremely high in cervical cancer. DNA methylation has shown a high sensitivity for cervical cancer and advanced CIN2/3. In addition, a well-studied methylation marker panel consisting of host cell genes FAM19A4 and miR124-2 showed a similar reassurance against CIN3 and cervical cancer compared with cytology up to 14 years of follow-up. Based on these results, it is assumed that DNAmethylation detects CIN2/3 lesions at highest risk of progression to cervical cancer. To date, the prognostic value of two methylation biomarker panels have been evaluated as a predictor of clinical regression in women with untreated CIN2 or CIN3. The S5 classifier, which evaluates methylation of EPB41L3 and the late (L1 and/or L2) regions of HPV16, HPV18, HPV31, and HPV33, was able to predict regression or progression of CIN2 in women <27 years. Recently, the QIASure Methylation Test®, which evaluates methylation of FAM19A4/miR124-2, was studied in a prospective follow-up study of women aged 18 to 55 years with untreated CIN2 or CIN3. Women with a negative FAM19A4/miR124-2 result on the baseline clinician-collected cervical sample showed more clinical regression (74.7%) than women with a positive methylation result (51.4%, $p=0.013$). Similar findings were found for baseline self-collected samples. The use of methylation markers as guidance for CIN2/3 management will be further discussed in this presentation.

Methods: --

Results: --

Conclusions: --

Oral presentation

Dr Jeannot Emmanuelle
emmanuelle.jeannot@curie.fr - France

Jeannot Emmanuelle
emmanuelle.jeannot@curie.fr - France

#3967

HPV CTDNA AS A MARKER FOR EARLY DETECTION OF RELAPSE IN CERVICAL AND ANAL CANCERS

15 - Molecular markers

Background/Objectives: Almost all cervical cancers (CC) are caused by human papillomavirus (HPV) which may be also responsible for other anogenital cancers, such as anal squamous cell carcinoma (ASCC). Chemoradiotherapy is the current standard care for locally advanced cervical or anal cancer but patients with advanced stage are at high risk for relapse. Circulating HPV DNA (HPV ctDNA) may serve as a residual tumor marker at the end of chemo-radiation or may predict relapse during the follow-up period.

Methods: We analyzed serum samples from patients with HPV16- or HPV18-related CC or ASCC. Samples were collected before and after treatment. We used digital droplet PCR (ddPCR) to detect circulating HPV DNA and to assess its prognostic impact as well as its prediction value of relapse.

Results: Circulating HPV DNA (HPV ctDNA) was detected in 63% (59/94) of patients with CC, before treatment. HPV ctDNA detection in serum sample was associated with high FIGO stage ($p=0.02$) and para-aortic lymph node involvement ($p=0.01$). In the cohort of patients with ASCC, HPV ctDNA was detected in 88% (29/33) of the patients, HPV ctDNA levels were also associated with tumor stage ($p=0.008$) and lymph node involvement ($p=0.03$). Complete clearance of HPV ctDNA by the end of treatment was significantly associated with a longer PFS ($p<0.0001$) in both cohorts. In the cervical cohort, analysis of additional serum samples taken during follow-up showed that patients with persistent HPV ctDNA in serum relapsed with a median time of 10 months (range, 2-15) from HPV ctDNA detection.

Conclusions: Circulating HPV DNA may be a useful marker to identify patients at risk of relapse of CC or ASCC.

Oral presentation

Dr Wei Feixue
weif@iarc.fr - France

Wei Feixue
weif@iarc.fr - France

#3968

Incidence and clearance of anal human papillomavirus infection in 13,928 individuals, according to HIV status, sex, and male sexuality: a collaborative pooled analysis of 26 longitudinal studies

03 - Epidemiology and natural history

Wei F, Georges D, Alberts C, Clifford G, Anal Hpv Study Group T

Background/Objectives: Understanding natural history of anal high-risk human papillomavirus (hrHPV) is key for designing anal cancer prevention programs, but has not been systematically characterised.

Methods: We reanalysed data from 26 studies with 13,928 individuals in six risk groups defined by HIV status, sex, and male sexuality: men who have sex with men (MSM) living with HIV (LWH) [MSMLWH], HIV-negative MSM, women LWH (WLWH), HIV-negative women, men who have sex with women (MSW) LWH (MSWLWH) and HIV-negative MSW. Markov models were used to estimate incidence and clearance of 13 hrHPV types and their determinants.

Results: HPV16 showed highest incidence/clearance-ratio of hrHPV types. MSMLWH had highest hrHPV incidence (e.g. 14.7% newly HPV16-infected within 2 years), then HIV-negative MSM (7.1%), WLWH (6.5%), HIV-negative women (2.9%), MSWLWH (1.3%), and HIV-negative MSW (0.6%). Determinants of HPV16 incidence included HIV-positivity and number of sexual partners for MSM and women, as well as anal sex for MSM but not for women. Anal HPV16 clearance was significantly lower for people LWH (PLWH), and significantly lower for prevalent than incident infections. Among MSM, increasing age was a strong predictor of persistence of prevalent, but not incident, HPV16 infections.

Conclusions: This robust and unifying pooled analysis of anal hrHPV natural history will underpin designing and predictions of impact of HPV-vaccination and HPV-based screening programs on anal cancer prevention, particularly in MSM and PLWH. Importantly, it establishes the evidence for the concept of differential carcinogenic potential of longstanding versus more recent anal hrHPV infection.

Oral presentation

Dr Bouchard Céline
celinebouchard2@gmail.com - Canada

Bouchard Céline
celinebouchard2@gmail.com -

#3970

Management and follow-up of women with CIN3-ADC in situ : The Colposcopic Approach

25 - Colposcopy

Background/Objectives: Describe the colposcopic approach of CIN3 / AIS in order to increase the expertise of the clinician.

Methods: Review of recent literature (publications and textbooks) will be provided. Colposcopic images from a selection of author's clinical experience will be presented.

Results: Colposcopic grading of CIN3/AIS lesions includes description of lesions with a focus on the characteristics such as location, contour, margin, and color. Revision of these features will be supported by clinical images. Reid's index and future use of Artificial Intelligence (AI) can help clinicians to achieve an appropriate identification of significant lesions in order to proceed with a biopsy at the appropriate area. AIS represents a diagnostic challenge for the clinicians and different colposcopic characteristics will be revised. Revision of the 3 main colposcopic expressions of AIS will be demonstrated with clinical images.

Conclusions: Colposcopy represents the optimal technique to identify CIN3 / AIS permitting conservative treatment of precancerous lesions, thus eliminating progression to cervical cancer. Increasing the quality of colposcopy training is still relevant to attain the goal of eliminating cervical cancer.

References: Modern Colposcopy, Textbook and Atlas, Third Edition, Mayeaux, Jr. E.J. et Cox JT Zhao Y et al. The Performance of Artificial Intelligence in Cervical Colposcopy: A Retrospective Data Analysis. J. of Oncol. 2022, Article ID 4370851, 6 pages. <https://doi.org/10.1155/2022/4370851> Perkins RB et al. 2019 ASCCP Risk-Based Management Consensus Guidelines for Abnormal Cervical Cancer Screening Tests and Cancer Precursors. J. Lower Gen. Tract Dis. 2020;24(2):102-131

Oral presentation

Mr Latsuzbaia Ardashesl
ardasheslatsuzbaia@gmail.com - Luxembourg

Latsuzbaia Ardashesl
ardashesl.latsuzbaia@sciensano.be - Belgium

#3971

NEW FINDINGS FROM THE VALHUDES STUDY: CLINICAL PERFORMANCE OF RIATOL QPCR HPV ASSAY ON SELF-COLLECTED VAGINAL AND URINE SAMPLES

13 - Self-sampling

Latsuzbaia A, Van Keer S, Vanden Broeck D, De Sutter P, Donders G, Doyen J, Tjalma W, Weyers S, Vorsters A, Arbyn M

Background/Objectives: The VALHUDES framework was established to evaluate the relative clinical accuracy of HPV testing to detect cervical intraepithelial neoplasia grade 2 or higher (CIN2+) on self-samples versus matched clinician-taken cervical samples. In this study we evaluated the clinical performance of Riatol qPCR HPV assay.

Methods: 523 women referred to colposcopy due to previous HPV infection or cervical abnormality were recruited at five Belgian colposcopy centres (NCT03064087). First-void urine was collected at home with Colli-Pee device containing 7 ml conservation medium (Novosanis) the day before the colposcopy clinic. Vaginal self-samples were collected with Evalyn Brush (Rovers Medical Devices) [n=247] or Qvintip (Aprovix) [n=276] and matched cervical samples were taken by gynaecologists with a Cervex-Brush (Rovers Medical Devices). Both, vaginal and cervical samples were suspended in 20ml PreservCyt LBC vials (Hologic Inc.). Colposcopy and histological assessment of biopsies determined the outcome, defined as CIN2+. The Riatol qPCR HPV assay was used for HPV detection considering 0.0001 HPV copies/cell as positivity threshold.

Results: 494 sample triplets (median age: 40 years, range 19-71) with valid HPV test and biopsy/colposcopy results were included in the study. Riatol qPCR HPV assay on urine was as sensitive and specific as on cervical samples (relative sensitivity=1.00 [95%CI 0.97-1.03]; relative specificity=0.98[95%CI 0.89-1.08]). Likewise, the relative accuracy on vaginal samples taken with Evalyn vs on cervical samples did not differ from unity (relative sensitivity=1.00 [95%CI 0.97-1.03]; relative specificity=1.06 [95%CI 0.93-1.20]). However, the accuracy on vaginal samples taken with Qvintip versus on cervical samples were lower, although not significantly (relative sensitivity=0.91 [95%CI 0.82-1.01]; relative specificity 0.89 [95%CI 0.80-1.00]). Sensitivity on vaginal samples collected with Evalyn Brush was higher than on samples collected with Qvintip (ratio=0.90 [95%CI 0.78-1.04]). We observed a significant decrease in β -globin DNA concentration by age in vaginal samples, and a significant increase in cervical samples, no relation in urine.

Conclusions: Clinical sensitivity and specificity of the Riatol qPCR HPV assay on self-collected vaginal and first-void urine samples were not significantly different from clinician-taken cervical samples. However, accuracy tended to be lower in vaginal specimens taken with Qvintip.

Oral presentation

Arroyo Mühr Laila Sara
Sara.Arroyo.Muhr@ki.se - Sweden

Laila Sara Arroyo Mühr
sara.arroyo.muhr@ki.se - Sweden

#3972

DEEP-DIVING FOR HPV IN CERVICAL CANCERS USING NEXT GENERATION SEQUENCING

02 - Viral and molecular biology

Laila Sara A, Camilla L, Jiangrong W, Agustin U, Joakim D, Karin S

Background/Objectives: Organized cervical HPV-based screening and HPV vaccination are two key factors for eliminating cervical cancer. Therefore, understanding in detail why some cancers may be HPV-negative is key. HPV testing is generally performed with PCR-based methods, which "only" detect HPV types they are designed to target. If there is a non-targeted HPV type in the specimen, a negative result will be reported. This may also happen if the HPV is mutated, or if a region of the viral DNA is fragmented/integrated in the human genome. Therefore, standard methods may result in false screening negatives. Next generation sequencing (NGS) moves beyond basic HPV detection to detect novel HPV types, HPV mutations and the whole virome. NGS further analyzes the tumor microenvironment; bacteria, fungi and others that can influence cancer development and prognosis. We present a summary of our work on NGS and how this has enabled us to further understand cervical cancer pathology.

Methods: We analyzed no less than 1300 cervical specimens from population-based studies nested in the Swedish organized cervical screening program. These samples derived from women with normal cervix, precancerous lesions and cervical cancers and we used NGS to investigate i) known and novel HPV types, ii) HPV mutations, iii) viral genomic fragmentation and iv) the meta-transcriptome present. NGS for HPV DNA and/or HPV RNA was performed on Illumina platforms NextSeq and NovaSeq using extensions of validated laboratory and bioinformatical analysis protocols.

Results: The majority (86%) of cervical cancers included tested positive for HPV with standard methods. NGS found HPV in a further 169/392 HPV negative cancers, translating into 7% final HPV negativity in cervical cancers. This final set of cancers is strongly correlated to poor prognosis in the patient, compared to patients with HPV-positive tumors. The extended set of HPV types identified by NGS could clinically be achieved through wide-spectrum PCR-assays; no new HPV types nor mutations in known HPV viral sequences were found so as to support potential failure of traditional testing methods. In the same set of cancers, we investigated the meta-transcriptome but did not detect any significant difference in such between HPV positive and negative cancers. Finally, detection of HPV RNA was a clear marker for ongoing high-risk HPV infection but did not provide a clear insight into risk for incident CIN2+.

Conclusions: Deep diving for HPV using NGS improves our knowledge of HPV biology in cervical cancer. Further studies should elucidate pathological processes underlying HPV negativity in cervical lesions.

Oral presentation

Dr Markowitz Lauri
lem2@cdc.gov - United States

Markowitz Lauri
lem2@cdc.gov - United States

#3973

HPV VACCINATION IN THE UNITED STATES: EFFECTS OF THE COVID-19 PANDEMIC, 2020-22

40 - Public health

Markowitz L, Stokley S, Pingali C, Herring B, Santoli J

Background/Objectives: The United States declared a national emergency in response to the COVID-19 pandemic on March 13, 2020. Stay-at-home orders and staffing shortages resulted in disruptions in preventive health care services. In the United States, routine HPV vaccination is recommended at age 11-12 years with catch-up through age 26. Most HPV vaccinations are provided in primary care settings and about 50% of adolescents are eligible for public purchased vaccines. We reviewed data to inform impact of the COVID-19 pandemic on HPV vaccination.

Methods: We present data on public purchased vaccine orders through the Vaccines for Children Program (VFC) for fiscal years (FY) 2019 through 2022 (as of February 13, 2022) and compare vaccine orders for FY2020-2022 (pandemic period) to FY2019 (pre-pandemic period). We also present 2019 and 2020 vaccination coverage data from the National Immunization Survey (NIS)-Teen.(1) NIS-Teen includes adolescents aged 13-17 years; the survey assesses vaccinations received throughout life, not limited to those administered in the survey year. Immunization data in NIS-Teen are from provider verified records.

Results: Public purchase HPV vaccine orders significantly decreased soon after the national emergency for COVID-19 was declared. During FY2020, HPV vaccine orders decreased 25% (1,026,593 doses) compared to FY2019. HPV vaccine orders improved in FY2021 but continued to be lower than in FY2019 (decrease of 9%; 391,101 doses). As of February 13, 2022, HPV vaccine orders were still lower than during the same time period in FY2019. HPV vaccination coverage (≥ 1 dose) among 13-17-year-olds in NIS-Teen 2019 and 2020, was 71.5% and 75.1%, respectively; up-to-date coverage was 54.2% and 58.6%. In 2020, most adolescents (73.6%) in NIS-Teen started the vaccine series before the pandemic began. Among those unvaccinated before March of the respective survey year, HPV vaccination initiation by April was lower in the 2020 survey than in the 2019 survey. The difference was largest in August/September (4.9 percentage points lower in 2020) but narrowed in subsequent months and was no longer significant by the end of November.

Conclusions: Public HPV vaccine orders through VFC in the United States significantly decreased in FY2020 and continue to be lower than pre-pandemic levels. Although national survey data show that coverage among 13-17-year-olds increased slightly in 2020 compared with 2019, most teens included in NIS-Teen 2020 started the HPV vaccination series before the pandemic began. While HPV vaccination coverage continues to improve, the HPV vaccination program is threatened by disruption of routine medical services due to the COVID-19 pandemic. Efforts are needed to increase coverage and to catch-up those who missed vaccinations over the past two years.

References: 1. Pingali C, Yankey D, Elam-Evans LD, et al. National, Regional, State, and Selected Local Area Vaccination Coverage Among Adolescents Aged 13-17 Years - United States, 2020. MMWR 2021;70:1183-1190.

Oral presentation

Prof. Hanley Sharon
sjbh1810@med.hokudai.ac.jp - Japan

Sharon Janet Hanley
sjbh1810@med.hokudai.ac.jp - Japan

#3974

HOW THE COVID-19 PANDEMIC SAVED JAPAN'S HPV VACCINATION PROGRAM

06 - HPV prophylactic vaccines

Sharon Janet H

Background/Objectives: Sensational media reporting involving the measles, mumps and rubella (MMR) vaccine in the 1990s and the human papillomavirus (HPV) vaccine in 2013 has resulted in a high degree of vaccine hesitancy in Japan. For the former, a successful class-action lawsuit was brought against the Ministry of Health Labour and Welfare (MHLW), by a small number of parents who claimed their children had experienced aseptic meningitis after MMR vaccination, for the latter several class-action lawsuits against both the MHLW and the manufacturers of the HPV vaccines due to purported adverse events are ongoing. In both instances, no evidence of a causal link was or has been presented.

Methods: Covid-19 vaccine rollout in Japan lagged behind Europe and the US due to the necessity of implementing clinical trials in Japanese populations before a vaccine can be licensed in Japan. The government also wanted to wait and observe for any safety issues in countries that had rushed to roll out Covid-19 vaccination programs before going ahead with its own program. Japan did not begin vaccinating the general public until late April 2021. However, in June 2021 with <40% of the population vaccinated and the Olympics quickly approaching, a daily target of 1,000,000 doses was set. From July 2021, vaccination of children 12-15 years also commenced. By the middle of August Japan recorded a record high of 26,000 Covid-19 cases due to the Delta variant and most of the country was under a State of Emergency. By November, Covid-19 vaccination coverage was around 80% with 70% coverage in those 12-19 years. Daily Covid-cases had fallen sharply to around 200 and on November 7th no deaths were recorded for the first time in 15 months. More importantly, no major safety issues with the vaccine had been reported in adolescents.

Results: On November 12th the Vaccines Adverse Reactions Review Committee concluded there was no reason to continue with the withdrawal of proactive recommendations for the HPV vaccine and on November 26th, the government formally announced proactive recommendations for the HPV vaccine would begin from April 2022, almost 9 years after the suspension. A catch-up program for girls born between 1997 and 2005 was also included.

Conclusions: While control measures such as wearing facemasks, physical distancing and vaccination have become political in many countries. In Japan this was never the case. Both mask wearing and vaccination was regarded as 'social courtesy' and a means to ending the burden the virus was putting on the economy and health services. Peer pressure is high in Japan, it influenced uptake of the Covid-19 vaccine; hopefully it will influence the recovery of the HPV vaccination program too.

Oral presentation

Dr Iacobone Anna Daniela
annadaniela.iacobone@ieo.it - Italy

Iacobone Anna Daniela
annadaniela.iacobone@ieo.it - Italy

#3975

TYPE SPECIFIC PERSISTENCE IN THE TEST OF CURE POPULATION AND IMPLICATIONS FOR MANAGEMENT.

14 - Genotyping

Iacobone A, Bottari F

Background/Objectives: Women diagnosed with cervical intraepithelial neoplasia grade 2 or worse (CIN2+) and treated by excisional procedures remain at high risk for recurrence over time. "Treatment failure" has been reported in up to 23% of women within two years after treatment. The aim of this study was to investigate the impact of HPV same genotype persistence on CIN2+ recurrence. As secondary outcomes, we evaluated the impact of multiple genotypes and of specific HR genotype, especially HPV 16/18, on recurrence risk.

Methods: A prospective cohort observational study was carried out at the European Institute of Oncology, Milan, Italy, from December 2006 to December 2014. HPV test was performed at the time of treatment ("baseline") and as "test-of-cure" at first follow-up visit planned at 6 ± 3 months after treatment. A ThinPrep PreservCyt (Hologic, Inc, Bedford, MA, USA) cervical sample was collected in all patients to perform Hybrid Capture 2 HR-HPV Test (HC2; Qiagen, Gaithersburg, MD, USA) and Linear Array HPV Genotyping Test (Roche Diagnostics, Pleasanton, CA, USA), in case of positive HC2 test. Two-year cumulative incidences for relapse were estimated and compared by the Gray's test.

Results: A total of 408 women surgically treated by excisional procedure for pre-neoplastic and neoplastic cervical lesions were enrolled. Overall, 96 (23.5%) patients were persistent for at least one genotype at three to nine months from baseline and 21 (5.1%) patients relapsed. Whereas histological grade, glandular crypt involvement, and margin status are not significantly related with treatment failure, HPV same genotype persistence has 30-fold increased odds of developing relapse ($p < 0.001$). The two-year cumulative relapse incidence was higher in HPV persistent patients compared to not-persistent (CIF = 27.6%, 95% CI: 16.2-40.2% versus CIF = 1.7%, 95% CI: 0.3-5.8%, $p < 0.001$), in women with persistent multiple infections (CIF = 27.2%, 95% CI: 7.3-52.3%, $p < 0.001$), and with the persistence of at least one genotype between 16 and 18, irrespective of the presence of other HR genotypes (CIF = 32.7%, 95% CI: 17.9-48.3%, $p < 0.001$), but not significantly different from women positive for single infections or any other HR genotype, but not for 16 and 18.

Conclusions: The risk of CIN2+ recurrence should not be underestimated when same HPV genotype infection persists after treatment, even in women with negative resection margins. HPV genotyping as "test-of-cure" enables a personalized risk-based management, by identifying women at higher risk of relapse who need intensive follow-up and avoiding risk of over-treatment in women with new HPV genotype infection after surgery.

References: Iacobone AD, Radice D, Sandri MT, et al. Human Papillomavirus Same Genotype Persistence and Risk of Cervical Intraepithelial Neoplasia2+ Recurrence. *Cancers (Basel)*. 2021;13:3664. Acknowledgments: This work was partially supported by the Italian Ministry of Health with Ricerca Corrente and 5×1000 funds.

Oral presentation

Dr Oštrbenk Valencak Anja
Anja.Ostrbenk@mf.uni-lj.si - Slovenia

Oštrbenk Valenčak Anja
anja.ostrbenk@mf.uni-lj.si - Slovenia

#3976

Commercial HPV tests: what's on the market and what's not (and what we desperately need)?

01 - HPV disease and COVID-19

Oštrbenk Valenčak A, Poljak M

Background/Objectives: Molecular tests for detection of human papillomaviruses (HPV) play a crucial role in the prevention of cervical cancer, including announced WHO's elimination efforts. HPV testing is a recommended approach for cervical cancer screening of women over 30 and for management of those with precancerous cervical lesions. In addition, they are widely used in epidemiological studies, HPV surveillance and vaccination impact monitoring. We present here the 2020 update of the inventory of commercial molecular HPV tests available on the market and outline implications for future research and policies.

Methods: Data from our internal files were retrieved and reviewed and a detailed search using Medline/Pubmed, Web of Science, Scopus, Google Scholar, Google and Bing, without language or period restrictions, was performed in September 2019 and again in January 2020.

Results: We have identified at least 254 distinct HPV tests (and 425 test variants) on the global market in 2020, which represent significant increase in comparison to 2015. Unfortunately, this increase came mostly in terms of quantity and at the detriment of quality. Current situation is quite alarming since 60% of the HPV tests on the global market are without a single peer-reviewed publication, 82% lack any published analytical and/or clinical evaluation, and over 90% are not evaluated in line with consensus requirements that ensure safe use in clinical settings.

Conclusions: Although COVID-19 pandemic had negative impact on cervical cancer control it might also generated opportunities for more efficient prevention. However, significant challenges in post COVID-19 era remains for both the HPV scientific community and the manufacturers of HPV tests.

Oral presentation

Dr Deshmukh Ashish
ashish.a.deshmukh@uth.tmc.edu - United States

Deshmukh Ashish
ashish.a.deshmukh@uth.tmc.edu -

#3977

Should screening for and treating anal HSIL be included in standard of care guidelines? What additional information is needed?

36 - Economics and modelling

Background/Objectives: New data from the Anal Cancer HSIL Outcomes Research (ANCHOR) study (indicating that anal HSIL treatment results in a 57% reduction in anal cancer incidence compared to active monitoring among HIV-infected persons with HSIL) provides a foundational benchmark to answer several unaddressed questions, including whether screening for and treating anal HSIL could be included in the standard of care guidelines and what additional data are critical to improve our understanding of population-level harms and benefits, necessary to inform optimal screening and treatment algorithms.

Methods: This talk will be guided by our ongoing efforts to synthesize a comprehensive mathematical modeling framework (Simulation Model of Anal Cancer: SMAC) of HIV, HPV, and anal precancer/cancer natural history that aims to determine effective and cost-effective age-specific anal cancer screening (for prevention and early detection) and anal HSIL treatment algorithms.

Results: Designed using empirical data and integrating anal HSIL treatment efficacy (57% reduction in cancer incidence) into the mathematical modeling (SMAC) framework, this talk aims to cover several pragmatic questions, including (1) downstream (i.e., extrapolation beyond-trial follow-up) implications of active monitoring versus anal HSIL treatment (in comparison with the disease natural history) and (2) key real-world attributes related to human behavior and access to care (e.g., treatment uptake and compliance) that will impact anal cancer prevention and/or early detection and further translation of these outcomes into survival benefits. In addition, the talk will also cover key questions, including (3) important disease natural history (e.g., progression/regression) and anal HSIL treatment-related aspects that will likely drive outcomes of screening and treatment, in terms of harms/benefits, stage-shift protection, quality of life, and downstream impact on survival, and (4) current uncertainties and additional data that is necessary to inform age-specific risk-based screening algorithms guided by the principle of equal risk/equal management.

Conclusions: Evidence-based optimal anal cancer screening guidelines are needed, informed by the determination of optimal use of screening (in terms of screening initiation/frequency/termination) for and treatment of anal HSIL. It is also crucial that currently scarce screening resources be directed to key risk groups to optimize screening use and ensure risk-targeted equitable prevention/early detection. This talk suggests a framework for integrating anal HSIL treatment efficacy into the anal cancer control continuum and highlights current uncertainties and the importance of addressing them to improve anal cancer prevention/early detection among key risk groups.

Oral presentation

Gillis Jennifer L.
jennifer.gillis@cw.bc.ca - Canada

Gillis Jennifer
jennifer.gillis@cw.bc.ca - Canada

#3978

Disparities in uptake of anal cancer screening

28 - Anal neoplasia

Gillis J

Background/Objectives: Implementation of anal cancer screening has the potential to significantly reduce the burden of anal cancer among people living with HIV. It is imperative that facilitators to screening are identified to ensure high levels of participation and equitable access to care. Using data from an ethno-racially diverse cohort of people living with HIV, our objective was to assess HPV-related knowledge, past participation in and beliefs regarding anal cancer screening and identify factors associated with willingness to participate in screening.

Methods: Results will be presented from cross-sectional questionnaires that were developed using the Theory of Planned Behavior to examine factors such as knowledge and beliefs that influence participation in HPV prevention strategies. The questionnaires were administered to men and women living with HIV in a multi-site HIV clinical cohort in Canada. These results will be presented in concert with emerging literature to discuss potential disparities in anal cancer screening participation as international guidelines are being considered.

Results: Racialized men were less likely to report undergoing anal cancer screening, regardless of sexual orientation, such that even among gay, bisexual and other men who have sex with men who are prioritized for preventive screening, there were racial inequities. This disparity in uptake of anal cancer screening occurred despite universal access to medically necessary services in our setting. However, the vast majority of men were willing to undergo screening in the future. Men generally had positive beliefs regarding the screening process, which were associated with increasing willingness to be screened, as posited by the Theory of Planned Behavior. Heterosexual men and racialized men reported lower willingness to be screened, though willingness was still very high among these men. Results on knowledge and willingness to undergo anal cancer screening among women living with HIV will also be discussed.

Conclusions: Observed disparities in knowledge, uptake, and willingness to undergo screening are notable, suggesting future disparities in anal cancer screening and outcomes. Accordingly, we believe these data are timely to inform development and implementation of anal cancer screening guidelines and community-led initiatives to encourage equitable attendance.

References: PLEASE NOTE: This abstract is submitted in reference to an invited speaker presentation -Scientific Session 4: Anal Cancer Screening / Chaired by A. Nyitray (USA), L. Abramowitz (France) / April 10 / 13:30 - 15:00 / Title: Disparities in uptake of anal cancer screening ACKNOWLEDGEMENTS: These data arise from a large anal cancer screening study among men living with HIV in Canada, the HPV-SAVE study (PIs: Troy Grennan & Irving Salit) and a study among women living with HIV (PIs: Ann Burchell & Jennifer Gillis). I thank all co-investigators of these studies (HPV-SAVE: Ron Rosenes, Jason Brunetta, Owen McEwen, Ann Burchell, Rupert Kaul, Paul MacPherson, Janet Raboud, Anita Rachlis, Alberto Severini, Darrell Tan, Jill Tinmouth, Daniel Grace, Joel Palefsky; HPV Prevention in WLHIV: Joanne Lindsay, Mona Loutfy, Claire Kendall, Janet Raboud, Anita Rachlis, Beth Rachlis, Anita Benoit, Mark Yudin, Gina Ogilvie, Abigail Kroch, Ramandip Grewal, Anna Yeung). Altogether, we gratefully acknowledge and thank the Ontario HIV Treatment Network (OHTN) Cohort Study participants, OHTN clinical and research staff, and Governance Committee members.

Oral presentation

Prof. Darragh Teresa
teresa.darragh@ucsf.edu - United States

Darragh Teresa
teresa.darragh@ucsf.edu - United States

#3979

New Pathogenetic Classification for Invasive Adenocarcinoma of the Endocervix: Implications for the Clinician

26 - Cervical neoplasia

Darragh T

Background/Objectives: In 2020, WHO updated the classification of endocervical adenocarcinoma. This new categorization was first championed by the International Endocervical Adenocarcinoma Criteria and Classification (IECC) in 2018.

Methods: Histologically, tumors are separated into the HPV-associated and HPV-independent categories based primarily on descriptive morphological criterion on routine H&E stains. The histologic subcategories of HPV-associated and HPV-independent cervical adenocarcinomas will be briefly discussed.

Results: Most cervical adenocarcinomas are HPV-associated and linked to HPV types 18, 16, and 45. Approximately 15% of cervical adenocarcinomas are HPV-independent. Clinically, stage is the most important prognostic variable; HPV-associated adenocarcinomas have a significantly better survival rate than HPV-independent adenocarcinomas of similar stage. Pattern of invasion is associated with risk for nodal metastasis.

Conclusions: This updated classification system and its clinical implications will be discussed.

References: WHO Classification of Tumours Editorial Board. Female genital tumours. Lyon (France): International Agency for Research on Cancer; 2020. (WHO classification of tumours series, 5th ed.: vol. 4). <https://publications.iarc.fr/592>.

Oral presentation

Dr Pimenoff Ville
ville.pimenoff@ki.se - Sweden

Pimenoff Ville
ville.pimenoff@ki.se - Sweden

#3980

Gender-neutral HPV vaccination: rapid decrease of high oncogenicity vaccine-targeted human papillomaviruses and an ecological response of less oncogenic types.

06 - HPV prophylactic vaccines

Pimenoff V, Gray P, Louvanto K, Eriksson T, Lagheden C, Söderlund-strand A, Dillner J, Lehtinen M

Background/Objectives: Human papillomavirus (HPV) vaccination programs target either girls only or both genders. Real life effectiveness and long-term effects of such programs can only be assessed with long-term follow-up of randomized trials.

Methods: We performed a long-term follow-up of 33 communities randomized to either gender-neutral HPV16/18 vaccination, girls-only HPV16/18 vaccination and hepatitis B-virus (HBV) vaccination of boys, and control vaccination. In the 1992-94 birth cohorts, 8,618/31,117 eligible boys and 15,615/30,139 eligible girls were vaccinated. Follow-up visits for cervico-vaginal sampling at ages 18 and 22 years were attended by 8,782 and 4,273 participants, respectively. Difference in prevalences of HPV types 6/11/16/18/31/33/35/39/45/51/52/56/58/59/66/68 was assessed and further modeled from the observed data.

Results: Rapid decrease of high oncogenicity vaccine-targeted HPV types 16/18/31/45 and significant increase of low oncogenicity HPV52 and 66 was observed for the gender-neutral vaccination communities eight years post-vaccination [likelihood ratio test (LRT) = 172.7, $P=0.001$] compared to the control communities, and also compared to girls only vaccination (LRT = 30.45, $P=0.031$). Compared to girls-only vaccination, gender-neutral vaccination associated with increased diversity ($P=0.00038$) of the circulating HPV types four to eight years post-vaccination. Simulations confirmed these diversity patterns eight years post-vaccination.

Conclusions: The gender-neutral vaccination has a faster decrease of high oncogenicity vaccine-targeted HPV types and a subsequent increase of low oncogenicity HPV types 52 and 66. The HPV types increasing after elimination of vaccine-targeted types have low oncogenicity and should thus not interfere the WHO milestone to eliminate cervical cancer. However, the post-vaccination diversity of remaining HPV types will affect future cervical cancer screening programs.

References: Lehtinen, M. et al. Gender-neutral vaccination provides improved control of human papillomavirus types 18/31/33/35 through herd immunity: Results of a community randomized trial (III). *Int. J. cancer* (2018) doi:10.1002/ijc.31618. Lehtinen, M., Baussano, I., Paavonen, J., Vänskä, S. & Dillner, J. Eradication of human papillomavirus and elimination of HPV-related diseases - Scientific basis for global public health policies. *Expert Rev. Vaccines* (2019). Vänskä, S. et al. Vaccination With Moderate Coverage Eradicates Oncogenic Human Papillomaviruses If a Gender-Neutral Strategy Is Applied. *J. Infect. Dis.* (2020) doi:10.1093/infdis/jiaa099.

Oral presentation

Dr Wang Jiangrong
jiangrong.wang@ki.se - Sweden

Wang Jiangrong
jiangrong.wang@ki.se - Sweden

#3981

OPTIMAL MANAGEMENT OF OLDER WOMEN TREATED FOR CIN: BALANCING THE RISKS AND BENEFITS

24 - Risk management

Wang J, Andrae B, Sahlgren H, Sparén P

Background/Objectives: Women having been treated for CIN are still confronted with increased risk of cervical cancer, particularly at older ages exiting routine screening program¹. The management for this group is challenging: morphological assessment and HPV testing performance may be hindered due to atrophy of cervix, and colposcopy evaluation can be unsatisfied due to the complication of cervical stenosis after CIN treatment. We aim to identify the challenges and discuss potential superior managements.

Methods: We used the Swedish National Cervical Screening Registry to examine the positivity proportion of HPV and cytology among women exiting screening program who had had history of CIN2+ (great majority of them were treated). When assessing HPV with partial genotyping result, 455 tests at ages 65-70 in Stockholm region during 2013-2020 with history of CIN2+ were included. When assessing cytology positivity, 20,635 tests at ages 65-70 in entire Sweden during 1981-2020 with history of CIN2+ were included.

Results: Among women aged 65-70 who had had history of CIN2+, 20% were tested positive for high-risk HPV, and 22% of the positive tests were HPV 16 or 18. The corresponding HPV positivity in general Swedish women population at ages 56-64 was around 8%, and 25% of the positive tests were HPV 16 or 18. On the other hand, only 9% of women aged 65-70 with history of CIN2+ were tested positive for cytology (ASCUS+). In a catch-up screening project, 566 women aged 65-70 who had had high-grade abnormality since age 50 without sufficient follow-up was targeted for self-sampling of HPV. Among 8 women who participated, tested high-risk HPV positive and participated in follow-up colposcopy, 4 had cervical stenosis thus were insufficient for histopathological diagnosis.

Conclusions: Our data implies that, HPV-primary testing may outperform cytology among older women exiting screening program with history of CIN2+. However, triage tests of either cytology or HPV genotyping are expected to be suboptimal. Moreover, colposcopy and biopsy for histopathology assessment are particularly challenging due to high prevalence of cervical stenosis. We are quantifying the effectiveness of current management strategies, hoping to facilitate the discussion about whether more aggressive treatment, e.g. hysterectomy, is necessary for certain groups. We also plan to explore whether other biomarkers, e.g. DNA methylation, may perform better on risk stratification for older women with CIN history.

References: 1.Strander B, Hällgren J, Sparén P. Effect of ageing on cervical or vaginal cancer in Swedish women previously treated for cervical intraepithelial neoplasia grade 3: population based cohort study of long term incidence and mortality. *BMJ*. 2014 Jan 14;348:f7361. doi: 10.1136/bmj.f7361. PMID: 24423603; PMCID: PMC3898577.

Oral presentation

Dr Jentschke Matthias
Jentschke.Matthias@mh-hannover.de - Germany

Jentschke Matthias
jentschke.matthias@mh-hannover.de - Germany

#3982

Assessing value of vaccine in women with cone

06 - HPV prophylactic vaccines

Jentschke M

Background/Objectives: HPV vaccination is the essential element in primary cervical cancer prevention. There is a growing body of evidence showing the value of HPV vaccination in the context of cervical excisional surgery.

Methods: We conducted a meta-analysis analysing the effect of pre- or post-conization vaccination (bi- or quadrivalent vaccine) against HPV. In the original meta-analysis from 2020 three retrospective and three prospective studies, three post-hoc analyses of RCTs and one cancer registry study were included after a systematic review of literature. Random-effect models were prepared to evaluate the influence of vaccination on recurrent CIN 2+.

Results: Overall, 21,059 patients (3,939 vaccinations vs. 17,150 controls) took part in the studies included in the meta-analysis. We observed a significant risk reduction for the development of new high-grade intraepithelial lesions after HPV vaccination (relative risk (RR) 0.41; 95% CI [0.27; 0.64]), independent from HPV type. There was a high heterogeneity between studies leading to a low level of evidence. The number of women needed to vaccinate before or after conization to prevent one case of recurrent CIN 2+ (NNV) is 45.5.

Conclusions: Our meta-analysis showed a significant risk reduction of developing recurrent cervical intraepithelial neoplasia after surgical excision and HPV vaccination compared to surgical excision only.

Oral presentation

Dr Jentschke Matthias
Jentschke.Matthias@mh-hannover.de - Germany

Jentschke Matthias
Jentschke.Matthias@mh-hannover.de - Germany

#3983

Kann das Einladungsmodell die Teilnehmerzahl erhöhen

11 - Screening for women difficult to reach

Jentschke M

Background/Objectives: 2020 wurde in Deutschland die neu organisierte Zervixkarzinomvorsorge mit Co-Testing (HPV-Test + Zytologie) in 3-Jahres-Intervallen für alle Frauen ab 35 Jahren gestartet. In der Vergangenheit lag die Teilnahme rate bei etwa 70 % in einem Zeitrahmen von drei Jahren. Die Nicht-Teilnehmerinnen haben ein höheres Risiko, an invasiven Gebärmutterhalskrebs zu entwickeln. Die Bereitschaft zur Teilnahme kann durch Einladungsschreiben oder das Angebot von Selbsttests für die häusliche Anwendung verbessert werden.

Methods: In der Hannoverschen Self-Collection Studie (HaSCo-Studie) evaluieren wir einen systematischen Ansatz zur Selbstuntersuchung auf humane Papillomaviren (HPV) bei Personen, die nicht am regulären Screeningprogramm teilnehmen. 20.000 Frauen im Alter von 30 bis 65 Jahren aus der Region Hannover, Niedersachsen, wurden nach dem Zufallsprinzip (Stichprobe des Melderegisters) rekrutiert. 10.000 Frauen erhielten direkt einen HPV-Selbsttest nach Hause geschickt, die anderen 10.000 ein Informationsschreiben mit der Möglichkeit zur Teilnahme an der Studie (Opt-out- bzw. Opt-in-Strategie). Es wurde eine Schichtung nach Alter (7 Kohorten) und Wohnort (Stadt vs. Land) vorgenommen.

Results: Zum Eurogin Kongress werden erste Daten vorliegen, wie gut sich Frauen auf diese Weise für einen HPV-Selbsttest motivieren lassen. Diese Daten können mit bereits vorhandenen Studien zum Einladungsmodell, z.B. der MARZY-Studie verglichen werden.

Conclusions: Um die Inzidenz des invasiven Zervixkarzinoms in Deutschland weiter zu senken, ist es essentiell, gezielte Strategien für Nicht-Teilnehmerinnen an der regulären Vorsorge zu entwickeln. Neben Einladungsverfahren ist die häusliche Selbstuntersuchung auf HPV eine vielversprechende Option. Die HaSCo-Studie wird von der Deutschen Krebshilfe unterstützt.

Oral presentation

Dr Arbyn Marc
Marc.Arbyn@sciensano.be - Belgium

Arbyn Marc
marc.arbyn@sciensano.be - Belgium

#3984

VALcolp: A protocol for clinical validation of portable colposcopy devices

25 - Colposcopy

Arbyn M, Jaafar I, Vergauwen G, Almonte M

Background/Objectives: Technical advances in medical imaging and artificial intelligence-based interpretation of digitized images have generated new optical devices that allow assisted visual assessment of the cervical surface applicable in remote low-resource areas as well as adjunctive sophisticated tools (e.g., spectroscopy, other optical processes) that may improve the quality of current colposcopy practice in well-equipped gynaecologic centres in high-income countries. However, their clinical accuracy for cervical precancer is poorly documented in peer-reviewed scientific literature and there is an urgent need for robust clinical validation of these new devices. VALcolp is a framework for validation and comparison of colposcopy devices involving a group of centres-of-excellence with world-widely recognized expertise in the field of management of cervical precancer. VALcolp aims to evaluate the relative sensitivity and specificity of new visual devices (simplified or advanced) to detect cervical precancerous lesions compared to high-quality standard colposcopy using histological assessment of biopsies as reference.

Methods: The core VALCOLP protocol follows principles of diagnostic test accuracy where the index test (new colposcopy device) and the comparator test (standard colposcopy) are applied to 500 women enrolled at colposcopy clinics and who subsequently have biopsies taken. The histological assessment of biopsies will be used as reference standard. All tests are applied at the same colposcopy visit and, where applicable, the result of the firstly performed test will be blinded to the assessor prior to performing the second test. Local adaptation of the generic protocol according to context and capacity may be considered.

Results: VALcolp will generate robust test accuracy data for multiple new colposcopy devices. VALcolp completed with literature data will yield a rich database that may be used to define internationally agreed criteria for validation of new generation colposcopy devices.

Conclusions: ValCOLP is an ambitious concept that finally will inform stakeholders which colposcopy devices may be considered as clinically validated.

References: 1. Arbyn M, Depuydt C, Benoy I, et al. VALGENT: a protocol for clinical validation of human papillomavirus assays. J Clin Virol 2016; 76 (Suppl 1): S14-S21. 2. Arbyn M, Peeters E, Benoy I, et al. VALHUDES: a protocol for VALidation of HUMAN papillomavirus assays and collection DEvices for HPV testing on Self-samples and urine samples. J Clin Virol 2018; 117: 52-6.

Oral presentation

Prof. Hanley Sharon
sjbh1810@med.hokudai.ac.jp - Japan

Hanley Sharon
sjbh1810@med.hokudai.ac.jp - Japan

#3985

HOW TO BECOME 'AT RISK' DUE TO AN ABANDONED VACCINATION POLICY: THE JAPANESE EXPERIENCE

06 - HPV prophylactic vaccines

Hanley S

Background/Objectives: The Japanese Government provided emergency public funding for the HPV vaccine from November 2010 for girls 12-16yrs born after 1994. The HPV vaccine was formally included in the national immunization programme for girls 12-16yrs from April 2013. In June 2013, after unconfirmed reports of adverse events following immunisation appeared in the media, the Japanese government suspended proactive recommendations for the vaccine. Despite a large body of evidence supporting the safety of HPV vaccination, the suspension of proactive recommendation continued until April 2022, almost 9 years. During this time coverage declined rapidly from >70% to <1%, even though the vaccine was still included in the Japanese national immunisation programme and provided free of charge for girls in the target age group. This talk will focus on how an abandoned vaccination policy has increased Japanese women's risk of cervical disease.

Methods: Results of modeling and real-world data will be reported.

Results: Using the Policy-1 Cervix model, Simms et al (2020) estimated an additional 25,000 cervical cancer cases and >5,000 additional deaths among females born between 1994 to 2007 could occur over their lifetime compared to if coverage had remained at 70%. They also estimated that If coverage of < 1% continued, around 60,000 preventable cases and 10,000 preventable deaths due to cervical cancer could occur in Japan over the next 50 years, compared to if uptake had remained at 70%. However, they also found that if 70% coverage could be restored in 12 year olds in 2020 along with 50% catch-up coverage in girls 13-20 years using the 9 valent vaccine, 70%-80% of these cases and deaths could be prevented. Another study by Yagi et al (2020) estimated that the risk of cervical cancer was reduced by 41% in girls born in 1998 compared to those born in 1993. However, they found that the risk for girls born after 2000 was similar to those born in 1993. For real-world data, Sekine et al (2021) reported a 2.1% prevalence of HPV16/18 in women born in 2000 in Niigata prefecture compared to a 0.3% prevalence in women born between 1994 and 1999. Finally, Yagi et al (2022) reported a 5.04% prevalence of cytological abnormalities in Japanese women born in 2000, compared to a prevalence of 3.76% in women born between 1994 and 1999.

Conclusions: Both modeling and real-world data has shown that women born after 2000 are at increased risk of cervical cancer due to Japan's abandoned HPV vaccination policy, compared to those born between 1994 and 1999. The extent to which this damage can be mitigated depends on how quickly HPV vaccination coverage returns to former levels after reintroduction of proactive recommendations for the vaccine in April 2022 for girls 12-16yrs and those born between 1997 and 2005.

Oral presentation

Dr Klein Sebastien
sebastian.klein@uk-koeln.de - Germany

Klein Sebastian
sebastian.klein@uk-koeln.de - Germany

#3986

Deep learning in histological samples of HPV associated oropharyngeal tumors

30 - HPV and oropharynx / Head and neck cancer

Klein S., Klussmann J

Background/Objectives:

Human papillomavirus (HPV) in oropharyngeal squamous cell carcinoma (OPSCC) is tumorigenic and has been associated with a favorable prognosis. Deep learning, has evolved as a powerful tool to predict molecular- and cellular alterations of medical images of various sources.

Methods:

We generated a deep learning-based HPV prediction score (HPV-ps) on regular H&E stains and assessed its performance to predict HPV-association.

Results: Although pathologists were able to diagnose HPV-association from H&E stained slides, (AUC=0.74, median of four observers) the inter-rater reliability was minimal (Light's Kappa=0.37; $p=0.129$), as compared to AUC=0.8 using the HPV-ps within two independent cohorts ($n=273$). The HPV-ps identified individuals with a favorable prognosis in a total of 594 patients from three cohorts (Giessen, OPSCC, HR=0.55, $p<0.0001$; Cologne, OPSCC, HR=0.44, $p=0.0027$; TCGA, non-OPSCC Head and Neck, HR=0.69, $p=0.0073$). Interestingly, the HPV-ps further stratified patients when combined with p16-status (Giessen, HR=0.06, $p<0.0001$; Cologne, HR=0.3, $p=0.046$).

Conclusions: Detection of HPV-association in OPSCC using deep learning with help of regular H&E stains may either be used as a single biomarker-or in combination with p16-status-to identify OPSCC patients with a favorable prognosis potentially outperforming combined HPV-DNA/p16-status as a biomarker for patient stratification. We anticipate that prediction of HPV association by deep learning from histology of routine histological samples of OPSCC tumors may be a viable biomarker in the near future.

References: Sebastian Klein, Alexander Quaas, Jennifer Quantius, Heike Löser, Jörn Meinel, Martin Peifer, Steffen Wagner, Stefan Gattenlöhner, Claus Wittekindt, Magnus von Knebel Doeberitz, Elena-Sophie Prigge, Christine Langer, Ka-Won Noh, Margaret Maltseva, Hans Christian Reinhardt, Reinhard Büttner, Jens Peter Klussmann, Nora Wuerdemann; Deep Learning Predicts HPV Association in Oropharyngeal Squamous Cell Carcinomas and Identifies Patients with a Favorable Prognosis Using Regular H&E Stains. Clin Cancer Res 15 February 2021; 27 (4): 1131-1138. <https://doi.org/10.1158/1078-0432.CCR-20-3596>

Oral presentation

Dr Faden Daniel
Daniel_Faden@meei.harvard.edu - United States

Faden Daniel
Daniel_Faden@meei.harvard.edu - United States

#3987

Circulating Tumor HPV DNA in the Diagnostic and Surgical Settings

30 - HPV and oropharynx / Head and neck cancer

Faden D

Background/Objectives: HPV-associated head and neck squamous cell carcinoma (HPV+HNSCC) is the most common HPV-associated malignancy in the United States and continues to increase in incidence. New ultrasensitive methods for detecting, diagnosing, and monitoring HPV+HNSCC are needed. Here, we will review emerging data regarding the use of circulating tumor HPV DNA as both a tool at the time of diagnosis and following surgery.

Methods: We conducted a prospective observational study in 140 subjects (70 cases and 70 controls) to determine if: 1. a noninvasive diagnostic approach for HPV+HNSCC would have improved diagnostic accuracy, lower cost, and shorter diagnostic interval compared with standard approaches, and 2. the clearance kinetics of ctHPVDNA are associated with postoperative disease status. Blood was collected, processed for circulating tumor HPV DNA (ctHPVDNA), and analyzed with custom ddPCR assays for HPV genotypes 16, 18, 33, 35, and 45. Diagnostic performance, cost, and diagnostic interval were calculated for standard clinical workup and compared with a noninvasive approach using ctHPVDNA combined with cross-sectional imaging and physical examination findings. In 33 patients with HPV+OPSCC undergoing surgery, blood was collected before surgery, postoperative days 1 (POD 1), 7, and 30 and with follow-up. A subcohort of 12 patients underwent frequent blood collections in the first 24 hours after surgery to define early clearance kinetics.

Results: Sensitivity and specificity of ctHPVDNA for detecting HPV+HNSCC were 98.4% and 98.6%, respectively. Sensitivity and specificity of a composite noninvasive diagnostic using ctHPVDNA and imaging/physical examination were 95.1% and 98.6%, respectively. Diagnostic accuracy of this noninvasive approach was significantly higher than standard of care (Youden index 0.937 vs. 0.707, $P = 0.0006$). Costs of noninvasive diagnostic were 36% to 38% less than standard clinical workup and the median diagnostic interval was 26 days less. Following surgery, in patients without pathologic risk factors for recurrence who were observed after surgery, ctHPVDNA rapidly decreased to <1 copy/mL by POD 1 ($n = 8/8$). In patients with risk factors for macroscopic residual disease, ctHPVDNA was markedly elevated on POD 1 (>350 copies/mL) and remained elevated until adjuvant treatment ($n = 3/3$). Patients with intermediate POD 1 ctHPVDNA levels (1.2-58.4 copies/mL) all possessed pathologic risk factors for microscopic residual disease ($n = 9/9$). POD 1 ctHPVDNA levels were higher in patients with known adverse pathologic risk factors such as extranodal extension >1 mm ($P = .0481$) and with increasing lymph nodes involved ($P = .0453$) and were further associated with adjuvant treatment received ($P = .0076$). One of 33 patients had a recurrence that was detected by ctHPVDNA 2 months earlier than clinical detection.

Conclusions: A noninvasive diagnostic approach for HPV+HNSCC demonstrated improved accuracy, reduced cost, and a shorter time to diagnosis compared with standard clinical workup and could be a viable alternative in the future. POD 1 ctHPVDNA levels are associated with the risk of residual disease in patients with HPV+OPSCC undergoing curative intent surgery and thus could be used as a personalized biomarker for selecting adjuvant treatment in the future.

Oral presentation

Prof. Hanley Sharon
sjbh1810@med.hokudai.ac.jp - Japan

Hanley Sharon
sjbh1810@med.hokudai.ac.jp - Japan

#3991

HPV-Impfung und Screening in Japan

40 - Public health

Hanley S

Background/Objectives: Die Inzidenzrate für Gebärmutterhalskrebs in Japan beträgt 15,3 pro 100.000 Frauen. Das ist fast zweimal so hoch wie in anderen Hocheinkommensländern im asiatisch-pazifischen Raum und ebenfalls höher als in vielen Mittel- und Niedrigeinkommensländern. Besonders besorgniserregend ist dabei der starke Anstieg der Fallzahlen bei Frauen im gebärfähigen Alter. Der Hauptgrund dafür ist die geringe Vorsorge durch HPV-Impfungen sowie Gebärmutterhalskrebs-Screenings.

Methods: Die aktuellen Zervixkarzinom-Vorsorgerichtlinien in Japan empfehlen eine Vorsorgeuntersuchung alle zwei Jahre mittels Pap-Abstrich für Frauen ab 20 Jahren. Eine Altershöchstgrenze existiert zur Zeit nicht. Es gibt kein organisiertes Screeningprogramm wie z.B. im Grossbritannien, allerdings haben Frauen diverse Möglichkeiten ein Screening vornehmen zu lassen. Sie können sich im Rahmen eines Gesundheits-Check-ups am Arbeitsplatz, kommunaler Screeningprogramme oder beim Gynäkologen untersuchen lassen. In der Realität kommen Frauen mit der Zervix-Vorsorgeuntersuchung das erste Mal bei einer Schwangerschaft in Berührung und es passiert nicht selten, dass dann bei der Feststellung ebendieser auch ein Gebärmutterhalskrebs entdeckt wird. 2020 wurden die Vorsorgerichtlinien zwar überarbeitet, aber es gilt weiterhin die Empfehlung alle zwei Jahre einen Pap-Abstrich durchzuführen, allerdings nur bis zu einem Alter von 65 Jahren. Ebenfalls wird Frauen im Alter zwischen 30-60 alle fünf Jahre die HPV-DNA-Diagnostik im Primärscreening empfohlen. Die Notwendigkeit einer HPV-DNA-Diagnose in Verbindung mit einer zytologischen Untersuchung in der Altersgruppe 30-60 wurde mit der Kategorie C als niedrig eingestuft - die Verfahren werden demnach nicht zur populationsbasierten Krebsvorsorge empfohlen. Ebenfalls wird von HPV-Selbsttests abgeraten. Momentan liegt die Vorsorgerate bei 30-40%.

Results: In Japan wurde der zweiwertige HPV-Impfstoff im Oktober 2009 lizenziert, der vierwertige HPV-Impfstoff im Juli 2011. Kurzzeitig, von 2010 bis 2012 wurden für den Einsatz des Impfstoffes bei Mädchen von 12-16 Jahren öffentliche Mittel zur Verfügung gestellt und ab April 2013 wurde der HPV-Impfstoff in das nationale Immunisierungsprogramm für Mädchen der o.g. Altersgruppe aufgenommen. Die anfängliche Impfabdeckung war mit 70-80% sehr hoch. Am 14. Juni 2013 jedoch, zwei Monate nach der formellen Aufnahme in das nationale Immunisierungsprogramm Japans wurden die aktiven Empfehlungen und Aufrufe zur HPV-Impfung im Zuge sensationsträchtiger Berichterstattungen in den Medien über unerwünschte Nebenwirkungen, die bisher in keinen kausalen Zusammenhang mit der Impfung gebracht werden konnten, zurückgezogen. Die Impfung blieb zwar weiterhin kostenlos für Mädchen von 12 bis 16 Jahren, aber die Kommunen und Ärzte durften die Impfung nicht mehr aktiv empfehlen. In der Folge fiel die Impfrate auf unter 1% und ist seither unverändert.

Conclusions: Im November 2021 kündigte das japanische Ministerium für Gesundheit, Arbeit und Wohlstand an, dass es wieder aktive Empfehlungen für den HPV-Impfstoff aussprechen werde, fast 9 Jahre nach der ursprünglichen Rücknahme von Empfehlungen. Ebenfalls angekündigt wurde, dass Nachholimpfungen angeboten würden für Frauen, die zwischen 1997 und 2005 geboren sind und sich bisher wegen Sicherheitsbedenken nicht haben impfen lassen. Aktuell schließt das japanische HPV-Impfprogramm sowohl die Verwendung des 9-fach Impfstoffs als auch Jungen von der Impfung aus.

Oral presentation

Dr Lewis James
james.lewis@vumc.org - United States

Lewis James
james.lewis@vumc.org - United States

#3992

Pathology-Based HPV Testing in Head and Neck Carcinomas - The Big Questions

09 - HPV testing

Lewis J

Background/Objectives: Human papillomavirus status (HPV) has become critical for oropharyngeal squamous cell carcinoma (SCC) patients. As the field is evolving, the type of HPV testing, whether with p16 immunohistochemistry alone, HPV-specific testing alone, or a combination of tests, is in flux. Further, cytology aspirate specimens present unique challenges for HPV testing and circulating high risk HPV DNA testing is emerging which may supplant tissue and fine needle aspirate specimen testing completely. Many very big questions with pathology-based HPV testing remain that will be explored in this talk.

Methods: Based on years of experience and as a co-chair of the College of American Pathologists' expert panel of HPV testing in head and neck carcinomas, the presenter will address critical questions in daily practice for oropharyngeal SCC patients.

Results: The four major questions revolve around the "messy middle", or the 5-10% of patients with mismatch of p16 and HPV-specific testing results, around the optimal testing for cytopathology fine needle aspirate specimens from cervical nodal metastases, around how pathology-based testing will evolve if circulating HPV DNA blood testing continues to improve and be used, and around whether (and where) testing of non-oropharyngeal SCCs will finally arrive in practice.

Conclusions: The field is moving rapidly and HPV testing has to evolve to match clinical needs and to apply the best testing for patient risk stratification and prognostication. Availability of more sophisticated and accurate HPV and surrogate marker testing will help to better tailor prognostication and individual patient treatment in oropharyngeal SCC.

Oral presentation

Gheit Tarik
gheitt@iarc.fr - France

Gheit Tarik
gheitt@iarc.fr - France

#3993

Validation of new rapid robust affordable HPV test based on CRISPR technology

09 - HPV testing

Gheit T

Background/Objectives: Approximately 90% of cervical cancer cases and deaths occur in low- and middle-income countries (LMICs). Human papillomavirus (HPV) nucleic acid testing for cervical cancer screening which is well established in developed countries is more effective in preventing cervical cancer than cytology-based methods. The development of HPV tests that are affordable and do not require laboratory infrastructures is essential for LMICs.

Methods: The recent outbreak of SARS-CoV-2 paved the way for the development of new molecular tools for such methodologies for viral detection. The combination of an isothermal amplification with CRISPR-based technology for high-risk HPV detection may represent an interesting opportunity for a more efficient screening in LMICs.

Results: An overview of the emerging CRISPR-based HPV testing will be provided. Several studies showed that the use of recombinase polymerase amplification (RPA) combined with CRISPR-Cas technology allowed specific and sensitive detection of individual or multiple high-risk HPV types in less than one hour of time. Our laboratory at the International Agency for Research on Cancer (WHO/IARC) is also evaluating this technology combined with E7 primers from a well validated multiplex HPV assay, the E7-MPG.

Conclusions: There is still a need to clinically validate the use of this new technology for cervical screening and assess its applicability in low resource settings.

Oral presentation

Dr Windon Melina
mbenso15@jhmi.edu - United States

Windon Melina
mwindon@jhmi.edu - United States

#3995

Patient Preferences and Decision-Making among Patients with HPV-Associated Cancers

30 - HPV and oropharynx / Head and neck cancer

Windon M

Background/Objectives: The era of human papillomavirus (HPV) - associated oropharyngeal cancers has brought with it a treatment paradigm shift toward deintensification for patients at lower risk of recurrence with an eye toward preserving long-term quality of life. Patients with early stage oropharyngeal cancer (OPC) are a population with unique needs among head and neck cancer patients. Often, OPC patients have a choice of primary treatment between upfront surgery and upfront radiation therapy, regimens which quote equal survival, thus representing a situation of clinical equipoise and an opportunity for patient-centered counseling. This presentation will review the current understanding of patient preference in head and neck cancer treatment decision-making and will identify strategies for addressing the literature gaps and for conducting patient preference research within head and neck surgical oncology.

Methods:

Results:

Conclusions:

References:

Oral presentation

Dr Rettig Eleni
emrettig@bwh.harvard.edu - United States

Rettig Eleni
emrettig@bwh.harvard.edu - United States

#3996

Detection of Circulating Tumor Human Papillomavirus DNA Before Diagnosis of HPV-Positive Head and Neck Cancer

15 - Molecular markers

Rettig E, Faden D, Sandhu S, Wong K, Faquin W, Warinner C, Stephens P, Kumar S, Kuperwasser C, Richmon J, Uppaluri R, Varvares M, Sethi R, Hanna G, Sroussi H

Background/Objectives: Human papillomavirus (HPV), most commonly HPV16, causes a growing subset of head and neck cancers (HNCs), including the majority of oropharynx cancers in many developed countries. Circulating biomarkers for HPV-positive HNC may allow for earlier diagnosis, with potential to decrease morbidity and mortality. The objective of this study was to determine whether circulating tumor HPV DNA is detectable in a greater proportion of archived plasma samples from individuals later diagnosed with HPV-positive HNCs than from cancer-free controls.

Methods: Cases were participants in a hospital-based research biobank with archived plasma collected at least 6 months before HNC diagnosis, and available archival tumor tissue for HPV testing. Controls were biobank participants without cancer or HPV-related diagnoses, matched 10:1 to cases by sex, race, age and year of plasma collection. Plasma and tumor tissue were evaluated for HPV DNA using a previously validated digital droplet PCR-based assay that quantifies tumor-tissue-modified viral (TTMV) HPV DNA.

Results: Twelve HNC patients with median age of 68.5 years (range, 51-87 years) were included. Ten (83.3%) had HPV16 DNA-positive tumors. ctHPV16DNA was detected in pre-diagnostic plasma from 3 of 10 (30%) patients with HPV16-positive tumors, including 3 of 7 (43%) patients with HPV16-positive oropharynx tumors. The timing of the plasma collection was 19, 34 and 43 months before cancer diagnosis. None of the 100 matched controls had detectable ctHPV16DNA.

Conclusions: This is the first report that ctHPV16 DNA is detectable up to several years before diagnosis of HPV16-positive HNC for a subset of patients. Further investigation of ctHPV16DNA as a biomarker for early diagnosis of HPV16-positive HNC is warranted.

Oral presentation

Prof. Bonanni Paolo
paolo.bonanni@unifi.it - Italy

Bonanni Paolo
paolo.bonanni@unifi.it - Italy

#3997

HPV vaccination and COVID, Europe

06 - HPV prophylactic vaccines

Background/Objectives: Human papillomavirus (HPV) is the commonest sexually transmitted virus worldwide, with first infection typically occurring soon after sexual debut. HPV-related diseases cause substantial morbidity and mortality globally [i]. Overall, 2.2 million new cancer cases were attributable to infections in 2018, representing 13% of all cancer cases (excluding non-melanoma skin cancers), with HPV counting for 690 000 new cases (ASIR 8.0 cases per 100 000 person-years), predominantly cervix uteri carcinoma 1. In Europe, cervical cancer ranks as the 9th most frequent cancer among women and the 2nd most common female cancer deaths in women aged 15 to 44 years [ii]. Cervical cancer is preventable and curable, as long as it is detected early and managed effectively.

Methods: In May 2018, the WHO Director-General announced a global call for action to eliminate cervical cancer. In August 2020 the World Health Assembly adopted the Global Strategy for cervical cancer elimination with the "90-70-90 Approach". This global strategy was promptly implemented by the European CanCer Organisation in December 2019 and by the European Union with Europe's Beating Cancer Plan of February 3, 2021, to accelerate the elimination of cervical cancer [ii], [iii].

Results: HPV vaccination is most effective if provided to both sexes. However, most countries in Europe do not currently vaccinate boys. HPV vaccination uptake also remains low in some countries. As of January 2020, in the World Health Organization European Region (WHO/ER) national recommendations for HPVv exist in 46/53 (87%) countries, of which 38 (83%), 2 (4%), and 6 (13%) countries provided full, partial, or no funding, respectively, for the primary cohort [i]. Fully or partially funded HPVv was provided for girls only in 25/53 (47%) countries and for both boys and girls in 15/53 (28%) countries and 30/53 (57%) WHO/ER countries reported VCRs. Monitored VCRs ranged from 4.3% to 99%. Of the 25 countries reporting a monitored VCR following at least two doses of HPVv, 24 (80%) pertained to a primary cohort of girls and 10/24 (42%) reported a high VCR (i.e. $\geq 71\%$), 6/24 (25%) reported a moderate VCR (i.e. 51-70%), 6/24 (25%) reported a low VCR (i.e. 31-50%), and 3/24 (13%) had a very low VCR ($\leq 30\%$). Only Portugal exceeded its target for the primary cohort. Just four countries with a GNV program reported a monitored VCR for boys, which in all cases was categorized as either low (Austria) or very low (Czech Republic, Italy, and Switzerland). The COVID-19 pandemic has significantly disrupted HPV vaccination and screening programmes across Europe.

Conclusions: The Region as a whole experienced a smaller decline for routine immunization coverages during 2020 than all other regions of the world, but this decrease masks large variations among countries, with larger declines in some countries, and a lack of reported data from some countries where COVID-19 disruptions have also affected capacity to collect and report routine immunization coverage data [ii]. Additional efforts are required to ensure HPVv NIPs are fully funded and high VCRs maintained, contrasting discrepancies in vaccination coverage to enable elimination of all HPV-related cancers. Now is the time for decisive action to create an HPV-cancer-free future for men and women across Europe.

References: [1] Catherine de Martel, Damien Georges, Freddie Bray, Jacques Ferlay, Gary M Clifford. Global burden of cancer attributable to infections in 2018: a worldwide incidence analysis. *Lancet Glob Health* 2020;8: e180-90. [1] Cervical Cancer Prevention Policy Atlas 2020. 22/01/2020. Available at: Cervical Cancer Prevention Policy Atlas 2020 | EPF (epfweb.org). [1] World Health Organization. Cervical Cancer Elimination Initiative. Available at: Cervical Cancer Elimination Initiative (who.int) [1] Peter Baker, Daniel Kelly, Rui Medeiros, Mike Morrissey, Richard Price. Eliminating HPV-caused cancers in Europe: Achieving the possible. *Journal of Cancer Policy* 28 (2021) 100280. [1] European Public Health Alliance. Europe's Beating Cancer Plan: the path towards cervical cancer elimination. Jul 7, 2021. Available at: Europe's Beating Cancer Plan: the path towards cervical cancer elimination - EPHA. [1] Paolo Bonanni, Pascaline Faivre, Pier Luigi Lopalco, Elmar A. Joura, Tobias Bergroth, Stefan Varga, Nathalie Gemayel & Rosybel Drury (2020): The status of human papillomavirus vaccination recommendation, funding, and coverage in WHO Europe countries (2018-2019), *Expert Review of Vaccines*, DOI: 10.1080/14760584.2020.1858057. [1] World Health Organization/Europe. Varied impact of COVID-19 on routine immunization in the European Region. 16-07-2021. Available at: WHO/Europe | Vaccines and immunization - Varied impact of COVID-19 on routine immunization in the European Region. [1] European Parliament resolution of 16 February 2022 on strengthening Europe in the fight against cancer - towards a comprehensive and coordinated strategy. Texts adopted - Strengthening Europe in the fight against cancer - Wednesday, 16 February 2022 (europa.eu).

Oral presentation

Daniels Vincent
vincent_daniels@merck.com - United States

Vincent Daniels
vincent_daniels@merck.com - United States

#3998

ESTIMATION OF HPV VACCINE MODEL PARAMETERS AND UNCERTAINTY FROM INTERIM ANALYSIS OF KEN-SHE TRIAL

36 - Economics and modelling

Vincent D, Kunal S, Cody P, Ya-ting C

Background/Objectives: Modeling the impact of off-label single dose (OLSD) vaccination programs requires parameters characterizing the effect of the vaccine on the target population. These parameters and their associated uncertainty are ideally derived from vaccine randomized controlled clinical trial (RCT) data analyzed in the context of a specific vaccination model. The KEN-SHE trial¹ in Kenya is one of the first RCTs specifically designed to study the efficacy of OLSD on prevention of HPV infections among adolescents and young women.

Methods: We employed a Markov chain Monte Carlo (MCMC) method to generate a probability distribution of model parameter values using the pooled HPV 16/18/31/33/45/52/58 KEN-SHE and control arm interim analysis of 18-month data². The resulting parameter values can be used to characterize effectiveness in models and provide an uncertainty distribution for probabilistic sensitivity analyses (PSA) of vaccine model outcomes. We first fit a 2-parameter, constant rate compartmental model with vaccinated-partially protected, vaccinated susceptible, and infected compartments. We estimate a joint distribution of the model's constant rates: "degree of protection", ψ , defined as the fraction of exposed vaccinated people that become infected (breakthrough infection); and "duration of protection", $\tau_{1/2}$, which is inversely proportional to the rate at which vaccinated people become completely susceptible - specifically, the time it would take for $\frac{1}{2}$ the vaccinated population to lose protection. We also explore sensitivity to the vaccine model structure assumptions.

Results: For the simple 2-parameter vaccine model the "most probable" parameter pair is $(\psi, \tau_{1/2}) = (93.0\%, 8.7 \text{ years})$ with 95% CI for ψ and $\tau_{1/2}$ of 74.2% to 99.3% and 1.5 years to 68 years, respectively. The parameters are correlated such that lower degree of protection is more likely associated with longer duration and vice versa. Similar results are obtained for other vaccines models.

Conclusions: Our results show that while the KEN-SHE provides evidence of high OLSD efficacy at 18 months, there is substantial uncertainty in the long-term effectiveness. In addition, the trial data provides a scientifically rigorous foundation for estimating model parameter values and uncertainty distributions that are critical inputs for health and economic models used to inform vaccination policy decisions. As more efficacy data on OLSD from RCTs become available in the coming years improved estimates (and more certainty) of these important model parameter inputs will be possible.

References: 1. Barnabas, R. V., et al. (2021). "Single-dose HPV vaccination efficacy among adolescent girls and young women in Kenya (the KEN SHE Study): study protocol for a randomized controlled trial." *Trials* 22(1): 661. TRIAL REGISTRATION: ClinicalTrials.gov NCT03675256 . Registered on September 18, 2018. 2. Barnabas, R. V, Brown, E. R., Onono, M. A., Bukusi, E. A., Winer, R. L., Galloway, D. A., ... Morrison, S. (2021). Efficacy of single-dose HPV vaccination among young African women. <https://doi.org/10.21203/RS.3.RS-1090565/V1> (not peer-reviewed)

Oral presentation

Dr Bonawitz Rachel
rachel.bonawitz@merck.com - United States

Bonawitz Rachel
rachel.bonawitz@merck.com - United States

#3999

IMMUNOGENICITY OF 2-DOSE REGIMENS OF 9 VALENT HUMAN PAPILLOMAVIRUS (9vHPV) VACCINE WITH EXTENDED DOSING INTERVALS: INTERIM RESULTS OF PROTOCOL V503-069

06 - HPV prophylactic vaccines

Bonawitz R

Background/Objectives: HPV vaccines are licensed as 2-dose regimen (administered 6 to 12 months apart) in 9-14-year-olds. Immunogenicity of dosing intervals >12 months is not well-characterized. Protocol V503-069 [NCT04708041] is an international, open-label study to evaluate the safety and immunogenicity of extended dosing intervals described previously (1). This interim analysis characterizes the immunogenicity and safety of the 9vHPV vaccine in boys and girls 10-15 years of age who received one dose of commercial 9vHPV vaccine dose (Dose 1) at least 1 year prior to study start and received a second dose (Dose 2) on Day 1 of the study.

Methods: Serum anti-HPV6/11/16/18/31/33/45/52/58 antibodies were evaluated by competitive Luminex Immunoassay at Day 1 and 4 weeks postdose 2. Anti-HPV GMTs at 4 weeks postdose 2 were evaluated with simple linear regression modeling, accounting for time interval between administration of Dose 1 and Dose 2. The impact of time interval between doses on the expected GMT was estimated as a change (increase or decrease) in the unadjusted GMT for every unit increase in time between doses. Participants were followed for injection-site and systemic adverse events (AEs) 15 days after each vaccination, and serious AEs for duration of participation.

Results: 146 participants enrolled and received 9vHPV vaccine on Day 1. The majority (96.6%) completed the trial 4 weeks postdose 2. Time between doses varied (median = 2.1 years; range 1.0 to 4.4 years). The geometric mean titers (GMTs) in mMU/mL at Day 1 represent GMTs corresponding to varying lengths of time postdose 1, ranging from 16.3 (HPV 45) to 104.6 (HPV 16). The GMTs at Month 1 represent GMTs at 4 weeks postdose 2 and ranged from 507.1 (HPV 45) to 22,237.2 (HPV 16). The data support a fold-change of 1.0 in unadjusted GMTs (i.e., unadjusted GMT x 1.0) for every 6 months increase in time interval between doses. No serious AEs were observed among participants.

Conclusions: This interim analysis is the first to describe immunogenicity and safety of extended interval dosing regimens of 9vHPV vaccine in adolescents. With a minimum interval of 1 year between the 2 doses of 9vHPV vaccine, the GMT at 4 weeks following the second dose of 9vHPV vaccine was not impacted by the varying time interval between doses (1 year to 4.4 years). The study is ongoing with additional analyses in cohorts of 9-14 year-old boys and girls who receive 2 doses at various intervals to demonstrate non-inferior immunogenicity of extended intervals compared with 16-26 year-old women who receive 3 doses.

References: 1. Teppler H et al. Design of a Phase III immunogenicity and safety study evaluating two-dose regimens of 9-valent human papillomavirus (9vHPV) vaccine with extended dosing intervals. *Contemp Clin Trials*: 2021 Jun;105:106403.

Oral presentation

Dr Ursu Ramona Gabriela
ramonagabrielaursu@yahoo.com - Romania

Dr Ursu Ramona Gabriela
ramonagabrielaursu@yahoo.com - Romania

#4000

The value of p16 and HPV DNA in non-tonsillar, non-base of tongue oropharyngeal cancer

30 - HPV and oropharynx / Head and neck cancer

Ursu R., Hammarstedt L, Holzhauser S, Zupancic M, Kapoulitsa F, Ramqvist T, Haegglblom L, Nasman A, Iancu L, Damian C, Marklund L, Dalianis T

Background/Objectives: Background: Oropharyngeal squamous cell carcinoma (OPSCC) is dominated by tonsillar and tongue base carcinomas (TSCC/BOTSCC), but there are carcinomas at other sites, such as uvula/soft palate/ pharyngeal wall here defined as other OPSCC. Human papillomavirus (HPV) positive TSCC/BOTSCC have favorable outcome, and the TNM-classification separates OPSCC into HPV mediated (p16INK4a overexpressing, p16+) and HPV unrelated OPSCC (p16INK4a non-overexpressing, p16-) cancer, but the prognostic role of p16+ in other OPSCC is unclear. Aims/Objectives: This study therefore aimed to further investigate the prognostic role of p16 +, presence of HPV DNA, or both combined in other OPSCC.

Methods: Material and methods: 195 other OPSCC, from patients diagnosed 2000-2018 were tested for p16, and/or presence of HPV DNA and the data correlated to outcome.

Results: Results: Neither overall survival, nor disease free survival correlated to presence of p16+ or HPV DNA in other OPSCC. p16+ and HPV DNA presence were correlated ($p<.0001$), but the sensitivity of p16 as a surrogate marker for presence of HPV DNA was low (49%).

Conclusions: Conclusions and significance: The data suggest that p16+ (and p16+/HPV DNA) positive other OPSCC should be analyzed cautiously and possibly separately from the HPV mediated OPSCC staging group.

References: Funding: The main author R.G.U. is supported by "Grigore T Popa" University of Medicine and Pharmacy, Iași, research contract 6983 from 21.04.2020.

Oral presentation

Mrs Kjaer Susanne
susanne@cancer.dk - Denmark

Kjaer Susanne K
susanne@cancer.dk - Denmark

#4001

HPV vaccine effectiveness against cervical cancer in Denmark

06 - HPV prophylactic vaccines

Background/Objectives: The total burden of cancers attributable to HPV is substantial and among HPV-related cancers, cervical cancer has the strongest association with HPV, as the development of virtually all cases is dependent on persistent infection with oncogenic HPV types. Therefore, vaccination against HPV is such an important primary prevention tool against cervical cancer.

Methods: From nationwide registries, information on HPV vaccination and cervical cancer diagnoses were retrieved. Incidence rates of cervical cancer in women vaccinated against HPV were compared to those in unvaccinated women, estimating incidence rate ratios (IRRs) with 95% confidence intervals (CIs) using Poisson regression with HPV vaccination treated as a time-varying variable and stratified by age at vaccination. We adjusted for attained age, education, and ethnicity.

Results: For women vaccinated at ages 16 years and younger or 17-19 years, the IRRs of cervical cancer were 0.14 (95% CI ¼ 0.04 to 0.53) and 0.32 (95% CI ¼ 0.08 to 1.28), respectively, compared with unvaccinated women. In women aged 20-30 years at vaccination, the incidence rate was higher than among unvaccinated women (IRR¼1.19, 95% CI ¼ 0.80 to 1.79) but slightly decreased with increasing buffer period (IRR¼0.85, 95% CI ¼ 0.55 to 1.32, with 4-year buffer period).

Conclusions: HPV vaccine effectiveness against cervical cancer at the population level is high among girls vaccinated at a young age.

Oral presentation

Ms Snoek Barbara
bcsnoek@hotmail.com - Netherlands

Snoek Barbara
bcsnoek@hotmail.com - Netherlands

#4002

Genome-wide microRNA analysis to inform biomarker design

15 - Molecular markers

Snoek B, Steenbergen R, Heideman D, Meijer C

Background/Objectives: MicroRNAs (miRNAs) are a class of small non-coding RNAs that have emerged as key regulators of gene expression. They are often found deregulated in cervical cancer and precancerous lesions and are increasingly recognized as promising biomarkers for cervical cancer diagnostics. Alongside altered expression in cervical (pre)cancer, miRNAs meet most of the requirements of being an ideal biomarker, such as their presence in small amounts of cervical specimens (i.e. tissues, scrapes, self-samples) and biofluids (i.e. blood, urine) allowing for relatively non-invasive sample collection. In addition, miRNAs are remarkably stable and can be assayed by a range of different methods. Next-generation sequencing is the preferred method for accurate and sensitive miRNA profiling, and as such numerous small RNA sequencing technologies have been developed. This has resulted in an explosion of biomarker studies and a comprehensive list of promising miRNAs for (cervical) cancer diagnostics.

Methods: ..

Results: ..

Conclusions: This presentation will give an overview of our expertise on genome-wide miRNA biomarker discovery for early detection of cervical cancer and summarizes recent findings on HPV-positive self-samples, HPV-immortalized cells and exosomes.

Oral presentation

Dr Nyitray Alan
anyitray@mcw.edu - United States

Nyitray Alan
anyitray@mcw.edu -

#4003

Suitability of clinician-performed Digital Anal Rectal Exams for detection of HPV-associated palpable lesions

16 - Screening methods

Nyitray A

Background/Objectives: The Digital Anal Rectal Examination (DARE) was fully described for the first time in 2018 by the International Anal Neoplasia Society. In July 2021, it was recommended by the United States Centers for Disease Control and Prevention for anal cancer screening in persons with HIV (PWH) and in HIV-negative gay and bisexual men. For people in areas with no high-resolution anoscopy infrastructure, DARE may be the only option for detection of anal cancer. Yet to date, there have been few reported data on the suitability and performance of DARE for anal cancer screening. Our objective was to describe the use of DARE in the Prevent Anal Cancer (PAC) studies at baseline.

Methods: Gay and bisexual cis-gendered men who were PWH or HIV-negative, and transgender persons who have sex with men, aged ≥ 25 years, were recruited into the PAC studies in the US from Chicago, Houston, and Milwaukee. Data from the first 560 participants to receive a DARE were analyzed. The performance of 16 PAC Study clinicians were described including DARE adequacy, length of time for DARE, and referral patterns. Lesion findings by DARE were stratified by HIV status and included lesion size, type, and location.

Results: Clinicians were 6 medical doctors (MD), 6 advanced practice nurses (APN), and 4 registered nurses (RN). Median age of enrolled participants was 42 years (range 25-80 years) and 31.0% were PWH. Eleven percent of participants reported a DARE in the prior year. Clinicians reported adequate examination of the perianus and anal canal in 556/560 participants. Median time to complete DARE was 1 minute and did not differ by presence of an abnormality. A total of 285 anal canal and perianal lesions among 211 (37.7%) participants were annotated with a higher proportion of lesions among PWH. Thirty persons had lesions at both anal canal and perianus. Of 79 participants with masses, focal areas of thickening or granularity, or linear lesions in the anal canal, 21 (26.5%) were labeled as "suspicious" or as condyloma. Of 162 participants with perianal lesions, 14 (8.6%) were labeled as "suspicious" or as condyloma. The median size of anal canal and perianal lesions was 3 mm (range 1-40 mm) and 3 mm (range 1-30 mm), respectively, with lesion size largest in Milwaukee. A total of 7.1% of participants were referred for additional care with PWH more likely to get referred ($p=0.02$). Compared to MDs, RNs were more likely to report anal canal and perianal abnormalities ($p=0.003$ and $p=0.002$, respectively) and more likely to make a referral ($p=0.04$). In a subset of 125 participants, $< 5\%$ reported the DARE as "quite a bit" or "very" embarrassing, uncomfortable, or anxiety producing.

Conclusions: Clinicians were able to recognize and describe anal canal and perianal lesions efficiently. The proportion of participants referred was low; however, RNs documented more lesions and referred participants to additional care more often than physicians. Referral patterns and variance in lesion size indicate the importance of DARE-specific training for HCPs. NCT03489707 and NCT04090060.

References: Hillman RJ, Berry-Lawhorn JM, Ong JJ, et al. International Anal Neoplasia Society guidelines for the practice of digital anal rectal examination. J Low Genit Tract Dis 2019;23:138-46. doi: 10.1097/lgt.0000000000000458

Poster

Dr Prieske Katharina
k.prieske@uke.de - Germany

Dr Prieske Katharina
k.prieske@uke.de - Germany

#4004

Transcriptome analysis in vulvar squamous cell cancer

15 - Molecular markers

Alawi M, Jaeger A, Wankner M, Eylmann K, Reuter S, Lebok P, Burandt E, Blessin N, Schmalfeldt B, Oliveira- Ferrer L, Joosse S, Woelber L

Background/Objectives: To date, therapeutic strategies in vulvar squamous cell carcinoma (VSCC) are lacking molecular pathological information and targeted therapy hasn't been approved in the treatment of VSCC, yet. Two etiological pathways are widely accepted: HPV induced vs. HPV independent, associated with chronic skin disease, often harboring TP53 mutations (mut). The aim of this analysis was to analyze RNA expression patterns for subtype stratification on VSCC samples that can be integrated in to the previously performed whole exome sequencing data for detection of prognostic markers and potential therapeutic targets.

Methods: We performed multiplex gene expression analysis (NanoString) with 770 genes, in 24 prior next generation sequenced samples. Integrative data analysis was performed

Results: 98 genes were differentially expressed in TP53mut vs. HPV+ VSCC, in the TP53mut cohort, 56 genes were upregulated and 42 were downregulated in comparison to HPV+ tumors. In the TP53mut subgroup Dickkopf-1 (DKK1: log2 FC: 3.69) and cell cycle regulators CCND1 (log2FC: 2.68), CCND2 (log2FC: 1.82), CCNA1 (log2FC: 3.22) and CDK6 (log2FC: 2.25) were observed to be highly upregulated compared to HPV+ VSCC. Within the TP53mutgroup a distinct cluster was identified which correlated to a significant worse overall survival ($p=0.017$). In the HPV+ subgroup genes involved in G1/S transition like CDKN2A, CCNE2, CDKN1C, CDKN2C, as well as S/G2 and G2/M phase regulators CDK2, Wee1, CDC25C, CDC7 and mini chromosome maintenance complex genes (MCM2/4/5/7) were particularly often upregulated. Also, genes involved in DNA repair like BRCA2, BRIP1, FANCA, FANCC, RFC4, PCNA, POLE2, RFC3, and EZH2 were upregulated compared to TP53mut tumors. None of the HPV+ tumors showed upregulation of cell cycle regulator cyclin D1, this were exclusively upregulated in TP53mut tumors.

Conclusions: RNA expression profiles show distinct regulation of selective candidate genes with regard to the known VSCC subtypes and potentially enable further subclassification in the TP53mut group. For HPV+ VSCC upregulation of cell cycle regulators was predominantly shown.

Oral presentation

Dr Van Trommel Nienke
n.v.trommel@nki.nl - Netherlands

Van Trommel Nienke
n.v.trommel@nki.nl - Netherlands

#4005

Detection of Cervical Cancer and Cervical Intraepithelial Neoplasia by hrHPV Analysis and DNA Methylation Analysis in Urine and Preferences for Cervical Cancer Screening Methods

18 - Methylation

Van Trommel N, Van Den Helder R, Schaafsma M, Van Splunter A, Mom C, Tjong M, Martin I, Rosier-van Dunné F, Van Der Avoort I, Renske S, Bleeker M

Background/Objectives: Offering urine as sampling method for cervical cancer screening may lead to increased attendance rates. Therefore, we aimed to determine the detection rate of cervical cancer and high-grade cervical intraepithelial neoplasia (CIN2/3) in urine and cervical (self)samples by analyzing the presence of high-risk human papillomavirus (hrHPV) and the levels of DNA methylation markers. Secondly, we evaluated preferences for different sample types for future cervical cancer screening in patients planned for a large loop excision of the transformation zone (LLETZ).

Methods: A total of 587 urine samples, cervicovaginal self-samples and clinician-taken cervical scrapes were included from 113 women with cervical cancer, 92 women with CIN2/3, and 64 controls. Samples were tested for hrHPV DNA and five methylation markers (ASCL1, GHSR, LHX8, SST, ZIC1). Using univariate and multivariate logistic regression and leave-one-out cross-validation, the methylation marker performance for detection of CIN3 and cervical cancer (CIN3+) in urine was determined. The agreement between samples was calculated using Cohen's kappa statistics and the Spearman correlation coefficients. In addition, 140 women, who were planned for LLETZ, filled in a questionnaire about their experiences with urine collection and cervicovaginal self-sampling and about their preferences for the sampling method of future cervical cancer screening.

Results: HrHPV presence was high in all sample types, 79% to 92%. All five methylation marker levels rose significantly with increasing severity of disease. The optimal marker panel (ASCL1/LHX8) resulted in an AUC of 0.84 for CIN3+ detection in urine, corresponding to an 86% sensitivity at a 70% predefined specificity. At this threshold 96% (109/113) of cervical cancers, 68% (46/64) of CIN3 and 58% (14/24) of CIN2 were detected. A strong agreement for HPV16/18 genotyping and a fair to strong correlation for methylation markers was observed between paired samples. Regarding patient's experiences with sample collection, patients considered urine collection compared to cervicovaginal self-samples as more acceptable ($p < 0.001$) and to provide more reliable results ($p < 0.001$). For future cervical cancer screening the majority of patients preferred urine collection (28%). Other preferred sampling types were: clinician-taken cervical scrape (23%); equal preference for urine collection, clinician-taken cervical scrape and cervicovaginal self-sampling equally (21%); cervicovaginal self-sampling (14%); equal preference for urine collection and cervicovaginal self-sampling (9%); equal preference for urine collection and clinician-taken cervical scrape (4%); missing (1%).

Conclusions: HrHPV DNA and DNA methylation testing in urine offers a promising solution to detect cervical cancer and CIN2/3 lesions. The majority of the patients preferred urine collection for future cervical cancer screening. Urine collection followed by molecular analysis may provide a solution to reach non-attendees of cervical cancer screening.

References: van den Helder, R., Steenbergen, R. D., van Splunter, A. P., Mom, C. H., Tjong, M. Y., Martin, I., Rosier-van Dunné, F. M. F., van der Avoort, I. A. M., Bleeker, M. C. G., & van Trommel, N. E. (2022). HPV AND DNA METHYLATION TESTING IN URINE FOR CERVICAL INTRAEPITHELIAL NEOPLASIA AND CERVICAL CANCER DETECTION. *Clinical cancer research: an official journal of the American Association for Cancer Research*. Schaafsma, M., van den Helder, R., Bleeker, M. C. G., Rosier-van Dunné, F. M. F., van der Avoort, I. A. M., Steenbergen, R. D., & van Trommel, N. E. (2022). Experiences and preferences towards collecting a urine and cervicovaginal self-sample among women attending a colposcopy clinic. *Preventive medicine reports*, 101749.

